



universität
wien

DISSERTATION

Titel der Dissertation

Projective Wellorders and the Nonstationary Ideal

verfasst von

Stefan Hoffelner

angestrebter akademischer Grad

Doktor der Naturwissenschaften (Dr.rer.nat)

Wien, im Juli 2016

Studienkennzahl lt. Studienblatt: A 791 405
Studienrichtung lt. Studienblatt: Mathematik
Betreuer: Sy-David Friedman

Abstract

We show that, under the assumption of the existence of $M_1^\#$, there exists a model on which the restricted nonstationary ideal $\text{NS}_{\omega_1} \upharpoonright A$ is \aleph_2 -saturated, for A a stationary co-stationary subset of ω_1 , while the full nonstationary ideal NS_{ω_1} can be made Δ_1 definable with ω_1 as a parameter. Further we show, again under the assumption of the existence of $M_1^\#$ that there is a model of set theory such that NS_{ω_1} is \aleph_2 -saturated and such that there is lightface Σ_4^1 -definable well-order on the reals. This result is optimal in the presence of a measurable cardinal.

Abstract

Unter der Annahme der Existenz von $M_1^\#$ wird ein mengentheoretisches Modell von ZFC konstruiert in dem das nonstationäre Ideal NS_{ω_1} auf ω_1 saturiert ist und in dem eine Σ_4^1 -definierbare Wohlordnung auf den reellen Zahlen möglich ist. Dieses Resultat ist optimal sobald eine messbare Kardinalzahl in dem Universum angenommen wird. Desweiteren wird, wieder unter der Annahme der Existenz von $M_1^\#$ ein mengentheoretisches Modell von ZFC konstruiert in welchem das auf eine beliebige, vorher fixierte stationäre, co-stationäre Teilmenge $A \subset \omega_1$ eingeschränkte nonstationäre Ideal $\text{NS}_{\omega_1} \upharpoonright A$ saturiert ist, während NS_{ω_1} selbst Δ_1 -definierbar ist mit ω_1 als einzigem Parameter.

Introduction

This thesis deals with two fundamental notions in set theory: definability and the saturation of the nonstationary ideal. The significance of the first for set theoretic investigations was already known to the early descriptive set theorists of the Russian and Polish schools who realized that the continuum hypothesis has a positive answer when looking at analytic (i.e. easily describable) sets only. Twenty years later K. Gödel showed how immensely powerful the notion of definability can be when used to investigate the set theoretic universe itself via his introduction of the constructible universe L . The quest of finding L -like universes which at the same time allow large cardinal properties is still one of the central open problems in set theory and is just another testimony of the importance of the concept.

The investigation of the saturation of the nonstationary ideal on ω_1 , NS_{ω_1} has a very interesting history as well. K. Kunen showed that given a huge cardinal there is a model in which NS_{ω_1} is \aleph_2 -saturated, a result which caused a considerable amount of attention as supercompact cardinals served as an informal upper bound for all natural consequences of large cardinal axioms, and hugeness is considerably stronger. Using a generic ultrapower argument he even argued that hugeness is the exact consistency strength of the assertion “ NS_{ω_1} is saturated”. Taking stationary subsets as conditions in a poset and picking a generic filter one arrives at a V -ultrafilter, which can be used to form the ultrapower $\text{Ult}(V, G)$ in $V[G]$. The resulting embedding $j : V \rightarrow M$ has ω_1 as critical point and $M^{<j(\omega_1)} \subset M$ does hold, which resembles the definition of hugeness. This intuition however turned out to be flawed as by the work of M. Foreman, M. Magidor and S. Shelah the forcing axiom MM (whose consistency follows from a supercompact cardinal) implies that NS_{ω_1} is \aleph_2 -saturated, which was later improved by S. Shelah who showed that already a Woodin cardinal suffices to construct a model in which NS_{ω_1} is saturated.

A natural question to ask is whether the saturation of NS_{ω_1} is consistent with CH . The original proof of Shelah for the saturation of NS_{ω_1} does not indicate an answer to this question. H. Woodin however, in [21], proved that if there is a measurable cardinal and if NS_{ω_1} is saturated then CH fails, in fact there exists a definable counterexample to CH . This impressive result indicates that there might be a surprising connection between the statements of CH and NS_{ω_1} being saturated.

Definability enters the picture in the following way. By a result of G. Hjorth (see [7]), a Σ_3^1 -definable wellorder of the reals together with the assumption that every real has a sharp implies CH . An investigation of the possibility of a set theoretic universe where NS_{ω_1} is saturated and the reals admit a projectively definable wellorder will therefore help to clarify the actual connection between CH and the saturation of NS_{ω_1} . One of the two main theorems of this thesis is the following:

Theorem 1. *Assume that $M_1^\#$ exists. Then there is a set theoretic universe in which NS_{ω_1} is saturated and there is a (lightface) Σ_4^1 -definable wellorder on the reals.*

Very roughly speaking its proof takes Shelah’s argument for making NS_{ω_1} saturated and adds forcings which will code reals by triples of ordinals, using a technique developed by A. Caicedo and B. Velickovic in [2]. Twisting their original definition makes a further localization forcing possible which codes all the relevant information for the wellorder in one real in such a way that any suitable, countable model can work with it, thus witnessing the relation.

This result is also interesting from a second perspective. It is often a challenging task to consider set theoretic universes with certain features, usually obtained assuming some large cardinal, and additionally equip them with definability properties. One can find a lot of literature devoted to this type of investigations such as [6], [3], and [1]. Though very diverse in their goals and means these results have in common that the ground model from which they start is L or some slight extension of it. This approach will not work when investigating possible models for “ NS_{ω_1} saturated and definable wellorders on the reals”. By a result of R. Jensen and J. Steel the statement “ NS_{ω_1} is saturated” is equiconsistent with the existence of a Woodin cardinal, thus one is compelled to work in inner models which will differ from L quite substantially in certain ways and render many of the usual arguments harder if not impossible. So this thesis can be seen as an attempt of finding coding methods which still work for stronger inner models, making the definability of certain sets possible. This also applies for the second result of this thesis.

The second theorem deals with the definability of NS_{ω_1} in the presence of a normal, saturated ideal on ω_1 .

Theorem 2. *Assume that $M_1^\#$ exists and that $A \subset \omega_1$ is a stationary, co-stationary set. Then there is a model of ZFC in which the nonstationary ideal restricted to subsets of A is saturated while the full ideal NS_{ω_1} is Δ_1 definable using ω_1 as a parameter.*

Its proof starts again with Shelah’s argument of making NS_{ω_1} saturated, and adds forcings which will write the characteristic functions of any stationary subset of ω_1 into a pattern of canonically definable trees which have a cofinal branch or not. These codings are also used in [1] to show that BPFA and a Σ_4^1 wellorder on the reals are simultaneously possible. We define a suitable class of models which is stationary and the fact that throughout the iteration there is always the stationary complement of A turns out to be crucial when ensuring that this class remains stationary along the way of the iteration. Again we will force to localize the relevant information in such a way that already any suitable model can read it off from a subset of ω_1 only, yielding the desired definability of NS_{ω_1} .

This theorem grew out of an attempt to answer a question in [5], where it is asked whether one could improve Woodin's result of $Con(\text{ZFC} + \text{"there are infinitely many Woodin cardinals"})$ implies $Con(\text{ZFC} + \text{"NS}_{\omega_1}$ is saturated"), and the techniques of our proof were initially intended to solve the problem for having the full NS_{ω_1} saturated. It turned out however that one faces big problems as the stationary class of nice models will lose its stationarity during the iteration as its factors are semiproper only. This raised a serious conflict with desired properties of our core model which served as the ground model during the iteration, and so we decided to tackle the case for the restricted ideal first.

We end this introduction with a short outline of how this thesis is organized. The first chapter very shortly defines the most important tools which are used in the proofs of the two main theorems. RCS -iterations and inner model theory play a fundamental role in the theorems so they are introduced. These belong to the more technical and abstract parts of set theory, but we hope that the reader can fully understand the thesis without being an expert in these subfields as long as he keeps in mind a couple of properties which, in the inner model case, we tried to state as additional axioms. Also a proof of Shelah's result of $Con(\text{ZFC} + \text{"there is a Woodin cardinal"})$ implies $Con(\text{ZFC} + \text{"NS}_{\omega_1}$ is saturated") is included as it is the starting point for our investigations.

In the second chapter we prove Theorem 2 and in the third chapter we prove Theorem 1. They can be read independently of each other.

Contents

1	Preliminaries	5
1.1	\mathcal{S} -semiproperness and RCS-iterations	5
1.2	The canonical inner Model with one Woodin Cardinal	10
1.3	How to make NS_{ω_1} \aleph_2 -saturated	16
2	$\text{NS}_{\omega_1} \upharpoonright A$ is Δ_1-definable	23
2.1	Introduction	23
2.2	The coding forcing	24
2.3	The class \mathcal{S}	26
2.4	The definition of the iteration	28
2.5	The sealing forcings	29
2.6	The coding forcings	29
2.6.1	\mathbb{Q}_α^0	29
2.6.2	\mathbb{Q}_α^1	30
2.6.3	Localization	31
2.7	The definability of NS_{ω_1}	34
2.8	$\text{NS}_{\omega_1} \upharpoonright A$ is \aleph_2 -saturated	35
3	NS_{ω_1} saturated and a projective well-order.	39
3.1	The coding method	39
3.2	Coding the reals	50
3.3	Definition of the iteration	52
3.4	NS_{ω_1} is saturated and a projective well-order. Open Questions.	57

Chapter 1

Preliminaries

1.1 \mathcal{S} -semiproperness and RCS-iterations

This section is a short reminder of the main definitions and properties of notions centered around the fundamental concept of properness. As is well known the definition of a proper notion of forcing was singled out by S. Shelah in the eighties. Proper forcings represent a class of forcings which is closed under iterations with countable support and which contain all the forcings which have the ccc and which are ω -closed. There is a weaker version of properness which Shelah called semiproper which still enjoys a lot of the nice properties of properness which however can not be iterated with countable support in a semiproperness preserving way. Nevertheless there is a generalization of countable support iterations, the so-called revised countable support iteration (RCS-iteration), which yields, applied to semiproper factors, always a semiproper notion of forcing. For our purpose a further generalization of semiproperness is needed, namely the so called \mathcal{S} -semiproper forcings. All these notions will be introduced now, the missing proofs can be found e.g. in M. Viale's notes [20] or in S. Shelah's [19].

Definition 3. *A notion of forcing \mathbb{P} is called proper if and only if there is a regular cardinal λ such that $\lambda > 2^{|\mathbb{P}|}$ and there is a closed unbounded $C \subset [H_\lambda]^\omega$ of elementary submodels $M \prec (H_\lambda, \in, <, \dots)$ (where $<$ defines a well-order. on H_λ and the dots should indicate further possible second order parameters) such that for every condition $p \in \mathbb{P} \cap M$ there exists a $q \leq p$ such that q is (M, \mathbb{P}) -generic; the latter meaning that every maximal antichain $A \subset \mathbb{P}$, $A \in M$ induces a predense subset $A \cap M$ below the condition q .*

It is very well known that there are other ways of defining what it means for a condition $q \in \mathbb{P}$ to be (M, \mathbb{P}) -generic, e.g. it is equivalent to the assertion that for every name for an ordinal $\dot{\alpha} \in M$, $q \Vdash \dot{\alpha} \in M$. Also for a partial order \mathbb{P} to be proper is equivalent to preserve stationary subsets of $[\lambda]^\omega$ for every uncountable cardinal λ .

The next notion is crucial in the proof of the first main result.

Definition 4. A class \mathcal{S} consisting of elements $M \in [H_\theta]^\omega$ for arbitrary cardinals θ is called everywhere stationary if the following two things hold:

1. for all regular cardinals θ , $\mathcal{S} \cap [H_\theta]^\omega$ is a stationary subset of $[H_\theta]^\omega$
2. \mathcal{S} is closed under truncation meaning that for all regular, uncountable cardinals θ and all $M \in \mathcal{S}$ $M \cap H_\theta \in \mathcal{S}$.

Now we can introduce a generalized form of proper forcing:

Definition 5. A notion of forcing \mathbb{P} is \mathcal{S} -proper for an everywhere stationary class \mathcal{S} if there exists a regular, uncountable cardinal θ such that $\theta > 2^{|\mathbb{P}|}$ and there is a club $C \subset [H_\theta]^\omega$ such that for every $M \in C \cap \mathcal{S}$ and every $p \in \mathbb{P} \cap M$ there is a $q \leq p$ which is (M, \mathbb{P}) -generic.

This definition can also be used if \mathcal{S} is not a class but a stationary subset of some $[H(\theta)]^\omega$. It is well known due to Shelah that \mathcal{S} -properness is almost as good as full properness, I.e. it is preserved under countable support iterations, preserves stationary subsets $T \subset [H_\theta]^\omega$ as long as $T \subset \mathcal{S}$ and therefore does not collapse \aleph_1 . In particular this means that during an iteration of \mathcal{S} -proper forcings with countable support, \mathcal{S} remains an everywhere stationary class.

Let us now turn to semiproperness:

Definition 6. Let \mathbb{P} be a partial order, then \mathbb{P} is semiproper if and only if there is a cardinal $\theta > 2^{|\mathbb{P}|}$ and there is a club $C \subset [H_\theta]^\omega$ of elementary submodels $M \prec (H_\theta, \in, <, \dots)$ such that every condition $p \in \mathbb{P} \cap M$ has an (M, \mathbb{P}) -semigeneric condition q below it; and a condition q is (M, \mathbb{P}) -semigeneric if and only if whenever $\dot{\alpha}$ is a name for a countable ordinal in M then $q \Vdash \dot{\alpha} \in M$.

This generalizes in a straightforward way:

Definition 7. Let \mathcal{S} be an everywhere stationary class. A notion of forcing \mathbb{P} is \mathcal{S} -semiproper if for a regular $\theta > 2^{|\mathbb{P}|}$ there is a club $C \subset [H_\theta]^\omega$ of elementary submodels $M \prec (H_\theta, \in, <, \dots)$ such that for every $M \in C \cap \mathcal{S}$ and for every $p \in M$ there is a $q < p$ which is (M, \mathbb{P}) -semigeneric.

It is well known that in fact something stronger is true:

Fact 8. If \mathcal{S} is everywhere stationary then \mathbb{P} is \mathcal{S} semiproper if and only if for every sufficiently large θ and every $M \prec H_\theta$, $M \in \text{Stat}$ such that $p, \mathbb{P} \in M$ there is a $q < p$ such that q is (M, \mathbb{P}) -semigeneric.

Proof. Let $\lambda > 2^{|\mathbb{P}|}$ then there is a club $C \subset H_\lambda$ such that for every $N \in C$ and every $p \in N$ there is a stronger q which is (N, \mathbb{P}) -semigeneric. Let $\theta > \lambda$ and let $M \prec H_\theta$, $M \in \mathcal{S}$ and let $<$ be a fixed well-order. of H_θ then M will have the $<$ -least club C witnessing the above as an element. Hence

$M \cap H_\lambda \in C$ and as $M \in \mathcal{S}$ we get $(M \cap H_\lambda) \in \mathcal{S} \cap C$. Thus we find for every $p \in M \cap H_\lambda$ a $q < p$ which is $(M \cap H_\lambda)$ -semigeneric. As $\lambda > 2^{|\mathbb{P}|}$, q is also (M, \mathbb{P}) -semigeneric. \square

A most important feature of proper notions of forcing is that iterating them with countable support results in a proper forcing again. (Its proof relies heavily on the fact that countable subsets in the extension can be covered by countable subsets of the groundmodel. This has as a consequence that whenever we break an iteration with countable support $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \delta)$ into two halves $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \beta)$ and $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \beta < \alpha < \delta)$, the second half still is a countable support iteration)

Contrary to proper forcings, semiproper forcings will not be preserved when using a countable support iteration. In order to give an example illustrating this fact we have to be able to break an iteration $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \delta)$ into two halves $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \beta)$ for an ordinal $\beta < \delta$, and $\dot{\mathbb{P}}_{[\beta, \delta]}$, where the second half is an iteration over the ground model $V^{\mathbb{P}^\beta}$ of length $\delta - \beta$ with factors which correspond to the $\dot{\mathbb{Q}}_\alpha$. Intuitively it is clear that such a construction should be possible, though on the one hand the factors $\dot{\mathbb{Q}}_\alpha$ are \mathbb{P}_α -names of partial orders, while in the yet not explicitly defined iteration $\dot{\mathbb{P}}_{[\beta, \delta]}$ the factors of the iteration should be \mathbb{P}_β -names of $\dot{\mathbb{P}}_{[\beta, \alpha]}$ -names of partial orders (where $\dot{\mathbb{P}}_{[\beta, \alpha]}$ is a \mathbb{P}_β -name for an iteration of length $\alpha - \beta$). Further the appropriate definition of the second half $\dot{\mathbb{P}}_{[\beta, \delta]}$ should have the consequence that the two step iteration $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \beta) * \dot{\mathbb{P}}_{[\beta, \delta]}$ is forcing equivalent to $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \delta)$. A precise analysis of the situation leads quickly to a rather tedious definition:

Definition 9. Let $(\mathbb{P}_\beta, \dot{\mathbb{Q}}_\beta : \beta < \alpha)$ be a forcing iteration of length α , let $\eta < \alpha$ then we let $\dot{\mathbb{P}}_{[\eta, \alpha]}$ be a \mathbb{P}_η -name defined like this:

$$\forall p \in \mathbb{P}_\eta \forall \dot{s} \in V^{\mathbb{P}^\eta} (p \Vdash_\eta \dot{s} \in \dot{\mathbb{P}}_{[\eta, \alpha]} \text{ if and only if } \forall q <_\eta p \exists r \in \mathbb{P}_\alpha \\ (r \restriction \eta < q \text{ and } r \restriction \eta \Vdash_\eta r \restriction [\eta, \alpha) = \dot{s}).$$

Here the term $r \restriction [\eta, \alpha)$ does not mean the usual check-name for the tail sequence as seen in $V^{\mathbb{P}^\eta}$ (as this could result in more forcing conditions than wanted - $V^{\mathbb{P}^\eta}$ could have more such sequences than V), but rather the \mathbb{P}_η -name for a function with domain $[\eta, \alpha)$ and values at stage ξ are $\dot{\mathbb{P}}_{[\eta, \xi]}$ -names according to the \mathbb{P}_ξ -name $r(\xi)$. Thus the above definition has the more readable characterization that for a \mathbb{P}_η -generic filter G_η over V we have that

$$V[G_\eta] \models \dot{\mathbb{P}}_{[\eta, \alpha]} = \{p \restriction [\eta, \alpha) : p \in \mathbb{P}_\alpha, p \restriction \eta \in G_\eta\}$$

With these clarified notions one can show that

Fact 10. *Every iteration $(\mathbb{P}_\beta, \dot{\mathbb{Q}}_\beta : \beta < \alpha)$ is forcing equivalent to the according two step iteration $(\mathbb{P}_\beta, \dot{\mathbb{Q}}_\beta : \beta < \eta) * \dot{\mathbb{P}}_{[\eta, \alpha]}$. And the forcing $\dot{\mathbb{P}}_{[\eta, \alpha]}$ is forced to be isomorphic to an iteration of length $\alpha - \eta$, i.e.*

$$1 \Vdash_\eta \dot{\mathbb{P}}_{[\eta, \alpha]} \text{ is isomorphic to an iteration of length } \alpha - \eta$$

One must be careful with the second statement of the last fact. Although for an iteration $(\mathbb{P}_\beta, \dot{\mathbb{Q}}_\beta : \beta < \alpha)$ the tail $\dot{\mathbb{P}}_{[\eta, \alpha]}$ can be seen as an iteration in $V^{\mathbb{P}_\eta}$, the type of the iteration can change. As an example take a countable support iteration $(\mathbb{P}_\beta, \dot{\mathbb{Q}}_\beta : \beta < \alpha)$ and an intermediate stage $\eta < \alpha$ such that in $V^{\mathbb{P}_\eta}$ the cofinality of an ordinal $\delta < \alpha$, $\eta < \delta$ gets changed from uncountable to countable cofinality. Then the cut off iteration with ground model $V^{\mathbb{P}_\eta}$ will not be a countable support iteration anymore as at stage δ , which has countable cofinality, nevertheless the direct limit is taken. This situation can not happen when the factors of the iteration are proper and a countable support iteration is taken, as in a generic extension obtained with a proper forcing all new countable sets of ordinals can be covered by old countable sets. The above described phenomenon can be exploited to construct an example of a countable support iteration of forcings which does collapse ω_1 , yet the factors are semiproper. In order to change the cofinality of an uncountable regular cardinal to cofinality ω we use Namba's forcing:

Definition 11. *The partial order $Nm(\omega_2)$, the so called Namba forcing, is defined like this: conditions are perfect trees $T \subset (\omega_2)^{<\omega}$, where perfect means that every node has \aleph_2 -many extensions. The ordering is given by $T_1 \leq T_2$ if and only if $T_1 \subset T_2$.*

One can show that under CH forcing with $Nm(\omega_2)$ preserves \aleph_1 and changes the cofinality of $(\omega_2)^V$ to ω . Moreover if one assumes the existence of a measurable cardinal, then after forcing with an ω -closed, thus CH preserving notion of forcing the Namba forcing is semiproper. Nevertheless countable support iterations of ω_1 -preserving notions of forcing, starting with a Namba forcing can collapse ω_1 :

Assume as a ground model a model where Namba forcing is semiproper and changes the cofinality of ω_2 to ω . Let $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \omega_2)$ be a countable support iteration of length ω_2 such that the first factor \mathbb{P}_0 is the Namba forcing. Further assume that at each stage $\alpha < \omega_2$, $V^{\mathbb{P}_\alpha}$ thinks that $\dot{\mathbb{Q}}_\alpha$ has an antichain of size \aleph_1 , a feature which can be arranged easily. Work in $V^{\mathbb{P}_0}$, fix a cofinal function $f : \omega \rightarrow \omega_2$ and for each $\alpha < \omega_2$ a \mathbb{P}_0 -name of an antichain $A_\alpha \subset \dot{\mathbb{Q}}_\alpha$ of size ω_1 in $V^{\mathbb{P}_\alpha}$. List $A_\alpha = (a_\beta^\alpha : \beta < \omega_1)$ and define a name for a function \dot{g} as follows:

Let \dot{g} be chosen such that for every $n \in \omega$ and every $\beta < \omega_1$ the Boolean value $\llbracket \dot{g}(n) = \beta \rrbracket = 1 \dot{\wedge} a_\beta^{f(n)} \dot{\wedge} 1$, I.e. at the $f(n)$ -th coordinate, the value is $a_\beta^{f(n)}$, while constantly 1 everywhere else.

Now \dot{g} is forced to be a surjective function from ω to ω_1 . Indeed let $p \in \mathbb{P}_{\omega_2}$ be a condition with support $\gamma < \omega_2$ and let $\beta < \omega_1$ be an arbitrary ordinal. Then there is an $n \in \omega$ such that $f(n) > \gamma$ and so p and $a_\beta^{f(n)}$ are compatible. Therefore for every ordinal $\beta < \omega_1$ it is dense to be in the range of \dot{g} and so it is surjective on ω_1 .

Thus one cannot hope in general that countable support iterations of semiproper forcings will preserve ω_1 . One has to iterate semiproper forcings in a more careful way if one wants to preserve ω_1 . Obviously the above example relies on the fact that after the Namba forcing the cofinality of ω_2^V has changed to ω , nevertheless at stage ω_2^V , due to the definition of the countable support iteration, we take the direct limit of the previous factors, though we actually should have taken the inverse limit. If we allow for our conditions in the iteration $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \omega_2)$ \mathbb{P}_β -names of countable sets, instead of the usual countable support, thus using a revised countable support, then the example above will not define a surjection of ω to ω_1 anymore. Indeed this modification will completely rule out the possibility of collapsing ω_1 as revised countable support iterations of semiproper forcings result in a semiproper forcing again.

Definition 12. *Let \mathbb{P}_α be a forcing notion, let $\alpha > 0$. Then \mathbb{P}_α is an RCS-iteration (short for revised countable support) of the factors \mathbb{P}_β , $\beta < \alpha$ if it is a subset of the inverse limit of the forcings $(\mathbb{P}_\beta : \beta < \alpha)$ such that each $p \in \mathbb{P}_\alpha$ satisfies*

- *for each $q < p$ there is an ordinal $\gamma < \alpha$ and a \mathbb{P}_γ -condition r such that $r \leq q \restriction \gamma$ and in the forcing \mathbb{P}_γ it holds that $r \Vdash_\gamma \text{cf}(\alpha) = \omega$ or for each $\beta \geq \gamma$ $p \restriction [\gamma, \beta) \Vdash_{\mathbb{P}_{\gamma, \beta}} p(\beta) = 1$*

Fact 13. *Iterations with RCS-support whose factors are semiproper result in a semiproper forcing notion. Moreover if we split an RCS iteration into two pieces then the tail iteration, as seen from the intermediate model will look like an RCS iteration again. More precisely, if $(\mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha : \alpha < \beta)$ is an RCS iteration then $1 \Vdash_\gamma \dot{\mathbb{P}}_{[\gamma, \beta)}$ is an RCS-iteration, for every $\gamma < \beta$.*

It is well known that properness is equivalent to the preservation of stationary subsets of $[\lambda]^\omega$. As a consequence the iteration of \mathcal{S} -proper forcings causes no new problems as long as we iterate with countable support. The class \mathcal{S} will remain stationary throughout the iteration and we end up with an \mathcal{S} -proper forcing, using the same proof as for countable iterations of proper forcings. For \mathcal{S} -semiproper notions of forcing \mathbb{P} we can still infer that they preserve stationary subsets of ω_1 as long as for every stationary $S \subset \omega_1$, $\{M \cap \omega_1 : M \in \mathcal{S}\} \cap S$ is stationary. Indeed if $S \subset \omega_1$ is stationary, if $p \in \mathbb{P}$, and if \dot{C} denotes a name of a club, then we pick an elementary submodel $M \in \mathcal{S}$ such that $M \cap \omega_1 \in S$ which contains \dot{C} . Let $q < p$ be an (M, \mathbb{P}) -generic condition then $q \Vdash M \cap \omega_1 \in \dot{C}$ and as $M \cap \omega_1 \in S$ we are done.

However when iterating \mathcal{S} -semiproper posets, for \mathcal{S} an everywhere stationary class, we have to take into account that usually the stationarity of \mathcal{S} gets lost during the iteration. Thus a statement like “A two-step iteration of \mathcal{S} -semiproper notions of forcing is \mathcal{S} -semiproper again” does not make sense in general. We can consider the natural enlargement of \mathcal{S} in $V[G]$ though, which adds to each element M of \mathcal{S} the G -interpretations of its names and which is the key notion when dealing with iterations.

Definition 14. *Let $S \subset [H_\theta]^\omega$ be stationary and let \mathbb{P} be an arbitrary notion of forcing with $2^{|\mathbb{P}|} < \theta$. Then set*

$$S[G] := \{M[G] : \mathbb{P} \in M \in S\}$$

This set $S[G]$ will remain a stationary set in $V[G]$:

Lemma 15. *Let $S \subset [H_\theta]^\omega$ be stationary and let \mathbb{P} be an arbitrary notion of forcing with $2^{|\mathbb{P}|} < \theta$ then $S[G]$ is a stationary subset of $V[G]$'s version of $[H_\theta]^\omega$.*

Proof. Fix a name \dot{C} for a club in $[H_\theta]^\omega$. Pick a countable $M \prec H_{\theta+}$ such that M contains \dot{C} and such that $M \cap H_\theta \in S$. Then \dot{C}^G is a club and is an element of $M[G]$, therefore $M[G] \cap H_\theta = (M \cap H_\theta)[G] \in \dot{C}^G$, yet $(M \cap H_\theta)[G] \in S[G]$. \square

Now this enables us to handle two step iterations:

Fact 16. *Let $S \subset H_\theta$ be stationary and assume that \mathbb{P} is a forcing notion with $2^{|\mathbb{P}|} < \theta$ which is S -proper. Let \mathbb{Q} be a notion of forcing in $V[G]$ which is $S[G]$ -proper, then the iteration $\mathbb{P} * \mathbb{Q}$ is S -proper.*

The last Proposition leads to

Fact 17. *Let $(\mathbb{P}_i, \dot{Q}_i : I \in \lambda)$ be an RCS-iteration of forcings for which*

$$1 \Vdash_i \dot{Q}_i \text{ is } \mathcal{S}[\dot{G}_I]\text{-semiproper}$$

holds at each stage $I < \lambda$. Then \mathbb{P}_λ is an \mathcal{S} -semiproper forcing notion.

1.2 The canonical inner Model with one Woodin Cardinal

Constructing universes of set theory with definable wellorders on the reals becomes a reasonable task only when the underlying ground model already satisfies a certain amount of definability which can be exploited. In the presence of sufficiently 'small' large cardinals Gödel's constructible universe L is the ultimate candidate for such a ground model, handing to the mathematician a lot of well studied tools to examine its structure thoroughly.

However as soon as we investigate properties of the set theoretic universe which are only achievable using stronger large cardinal hypotheses L will not suffice anymore as these hypotheses are inconsistent with $V = L$. The statement NS_{ω_1} is \aleph_2 -saturated is exactly one of such properties which need a quite large cardinal, namely a Woodin cardinal, whose existence contradicts $V = L$. We are thus compelled to use a model which is not L , yet having some of its nice properties, plus containing a Woodin cardinal. Of course Inner model theory equips the set theorist with such a model, named M_1 which shall be introduced here shortly.

As inner model theory relies heavily on a lot of nontrivial notions and definitions, whose introduction would soak up a lot of space we will skip them and assume the reader is already familiar with concepts such as

- the notion of an extender, its length and support, ultrapower constructions and their relation to elementary embeddings of the universe,
- Jensen's fine structural hierarchy,
- the notion of acceptability,
- fine extender sequences,
- the definition of active and passive premice,
- the notion of projecta, universality, solidity and soundness,
- iteration games played on sufficiently iterable premice, thus generating an iteration tree and the notion of branches in the iteration tree which drop in model or degree,
- the comparison process of sufficiently iterable premice,
- the definition of a Woodin cardinal,
- plus other notions we probably have forgot to mention.

One can find these concepts introduced e.g. in [11], [13], [9]. It should be noted however that a reader who does not know any of these can skip to the end of the section while still being able to fully understand the thesis, keeping in mind a couple of properties of M_1 which are mentioned as a list of properties stated as Fact 24, and which can be seen as additional axioms we will use during our proofs.

Now to the definition of M_1 : recall first that a premouse \mathcal{M} is called tame whenever its extenders do not overlap a local Woodin cardinal, i.e. whenever E is an extender on the \mathcal{M} sequence and $\lambda = lh(E)$ then

$$\mathcal{J}_\lambda^{\mathcal{M}} \models \forall \xi > crit(E) (\xi \text{ is not Woodin}).$$

We define recursively a sequence of premice \mathcal{N}_ξ starting with

$$\mathcal{N}_0 := (V_\omega, \in, \emptyset, \emptyset).$$

Suppose now that the premouse \mathcal{N}_ξ has already been defined. Then we consider the ω th core of \mathcal{N}_ξ and stop the construction if it does not exist. If it exist we set $\mathcal{M}_\xi := \mathcal{C}_\omega(\mathcal{N}_\xi)$ and split into cases:

1. if $\mathcal{M}_\xi = (J_\gamma^{\vec{E}}, \in, \vec{E}, \emptyset)$ is passive and there is an extender F^* over V , further an extender F over \mathcal{M}_ξ and an ordinal $\nu < \gamma$ such that

$$V_{\nu+\omega} \subset \text{Ult}(V, F^* \text{ and } F \upharpoonright \nu = F^* \cap ([\nu]^{<\omega} \times J_\gamma^{\vec{E}}).$$

Assume further that the structure $(J_\gamma^{\vec{E}}, \in, \vec{E}, F)$ is a tame premouse then pick the least such ν and set

$$\mathcal{N}_{\xi+1} := (J_\gamma^{\vec{E}}, \in, \vec{E}, F)$$

2. if $\mathcal{M}_\xi = (J_\gamma^{\vec{E}}, \in, \vec{E}, H)$ is not passive, (i.e. $H \neq \emptyset$) or there does not exist a background extender F^* , just continue in the J -hierarchy. Set

$$\mathcal{N}_{\xi+1} = (J_{\gamma+1}^{\vec{E} \hat{\ } H}, \in, \vec{E} \hat{\ } H, \emptyset).$$

In the limit steps λ , we let $\omega\eta = \lim \inf_{\xi < \lambda} (\rho_\omega^+)^{\mathcal{M}_\xi}$ and let \mathcal{N}_λ be the unique passive premouse \mathcal{P} of height $\omega\eta$ such that for every $\beta < \eta$, $\mathcal{J}_\beta^{\mathcal{P}}$ is the eventual value of $\mathcal{J}_\beta^{\mathcal{M}_\xi}$ as $\xi \rightarrow \lambda$.

We can use the just described sequence of premice

$$\mathbb{C}^t := (\mathcal{N}_\xi : \mathcal{N}_\xi \text{ exists})$$

to define M_1 , assuming the existence of a Woodin cardinal. As a reminder let us introduce the notion of 1-smallness:

Definition 18. *Let \mathcal{M} be a premouse, then we say that \mathcal{M} is 1-small if the following holds: whenever λ is the critical point of an extender on the \mathcal{M} -sequence, and $\delta < \lambda$ then*

$$\mathcal{J}_\lambda^{\mathcal{M}} \not\neq \text{ there is no Woodin cardinal above } \delta.$$

Suppose first that for every ξ , \mathcal{N}_ξ is 1-small, then it is a result of J.Steel [14] that for every $\xi < \text{Ord}$, \mathcal{N}_ξ is defined and so is \mathcal{N}_∞ . In this case we let M_1 be \mathcal{N}_∞ which is a class sized model which can be shown to contain exactly one Woodin cardinal.

Otherwise if there is a $\xi < \text{Ord}$ such that \mathcal{N}_ξ is not 1-small then fix the least such ξ . \mathcal{N}_ξ is an active premouse. Let $M_1^\#$ be $\mathcal{C}_\omega(\mathcal{N}_\xi)$ and let \mathcal{P} be the result of iterating the last extender of $M_1^\#$ out of the universe. Set $M_1 := \mathcal{J}_\infty^{\mathcal{P}}$, then again M_1 is a class sized model with one Woodin cardinal. To summarize:

Theorem 19. *If there is a Woodin cardinal then in both cases M_1 is a class sized model with exactly one Woodin cardinal and all its initial segments $\mathcal{J}_\beta^{M_1}$ are 1-small and ω -sound.*

We will always assume that $M_1^\#$ exists, i.e. $M_1^\#$ is the least mouse which is not 1-small and M_1 is the result of iterating away the last extender of $M_1^\#$.

It is a wellknown fact that reals which are elements in a sufficiently iterable premouse \mathcal{M} admit an easy (i.e. Δ_2^2) definition with a countable ordinal as parameter, using comparability of mice. The formula for $r \in \mathcal{M}$ just reads like this: r is the α -th real in some sufficiently, i.e. $(\omega, \omega_1 + 1)$ -iterable premouse. Thus the definability of reals in mice is tied to the expressibility of 'sufficient' iterability conditions in the sense of iteration games. For the model M_1 , as was observed by J. Steel in [15] an even easier definition is possible, due to a weakening of the usual iteration game which is still enough to guarantee a certain amount of comparability. We say that a premouse \mathcal{M} is Π_2^1 -iterable if player II has a winning strategy for $\mathcal{I}(\mathcal{M})$, where the latter denotes the new iteration game played on the premouse \mathcal{M} . Roughly speaking J. Steel showed that

- if \mathcal{M} is a premouse which is embeddable into a model of the \mathbb{C}^t -sequence then player II has a winning strategy for the $\mathcal{I}(\mathcal{M})$,
- if on the other hand II has a winning strategy for $\mathcal{I}(\mathcal{M})$ played on the premouse \mathcal{M} then \mathcal{M} can be compared with any premouse \mathcal{N} which embeds into an element of the \mathbb{C}^t -sequence,
- and finally the set of premice $\{\mathcal{M} : \text{II has a winning strategy in } \mathcal{I}(\mathcal{M})\}$ is Π_n definable.

We can use this to observe that a low complexity definition of countable (countable in M_1 that is) initial segments of M_1 is possible in generic extensions $M_1[G]$ which preserve ω_1 . We consider the set of countable premice which are Π_2^1 -iterable, ω -sound, 1-small, and which project to ω .

$$B := \{\mathcal{M} \text{ ctbl premouse} : \mathcal{M} \text{ is } \Pi_2^1\text{-iterable, } \omega\text{-sound, 1-small } \rho_\omega(\mathcal{M}) = \omega\}.$$

If we are in a forcing extension $M_1[G]$ which preserves ω_1 then using Shoenfield absoluteness, the set of countable premice which are initial segments of M_1 and which project to ω form a set of premice which are still Π_2^1 -iterable in $M_1[G]$, ω -sound and 1-small. If we consider in $M_1[G]$ an arbitrary element \mathcal{M} of B and let $\mathcal{N} = \mathcal{J}_\eta^{M_1}$, $\eta < \omega_1$ be an initial segment of M_1 which projects to ω then we can compare these two as \mathcal{M} is Π_2^1 -iterable which suffices for comparison as mentioned above. As both \mathcal{M} and \mathcal{N} are ω -sound and ω -projecting they actually do not move during the iteration and therefore we get that $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$ must hold. If we let the height of $\mathcal{N} = \mathcal{J}_\eta^{M_1}$ vary we see that there is certainly an $\eta < \omega_1$ such that

$\mathcal{M} \trianglelefteq \mathcal{N} = \mathcal{J}_\eta^{M_1}$. Thus the set B defines in $M_1[G]$ a set of initial segments of M_1 , which is cofinal in that for every countable $\mathcal{M} \trianglelefteq M_1$ there is an $\mathcal{N} \in B$ such that $\mathcal{M} \trianglelefteq \mathcal{N}$. As Π_2^1 -iterability is a Π_2^1 -notion, B is itself Π_2^1 -definable. Thus we have shown:

Lemma 20. *Let $M_1[G]$ be an ω_1 -preserving forcing extension of M_1 . Then in $M_1[G]$ there is Π_2^1 -definable set B of premice which are of the form $\mathcal{J}_\eta^{M_1}$ for some $\eta < \omega_1$. B is defined as*

$$B := \{\mathcal{M} \text{ ctbl premouse} : \mathcal{M} \text{ is } \Pi_2^1\text{-iterable, } \omega\text{-sound and not 1-small}\},$$

and the set

$$\{\eta < \omega_1 : \exists \mathcal{N} \in B(\mathcal{N} = \mathcal{J}_\eta^{M_1})\}$$

is cofinal in ω_1 .

Next we want turn to the concept of generic absoluteness. The motivation for this will become clear later once we turn to the proofs of the thesis. For now we just state that we would want our ground model V to be definable in generic extensions $V[G]$, for G a generic filter for a poset \mathbb{P} , i.e. we need a first order formula $\Phi(x)$ such that the class $\{x : V[G] \models \Phi(x)\}$ outputs V again. To achieve this we proceed indirectly using Steel's core model K . As its definition is very involved we will skip it (the interested reader can find it in [18]) and just state that it is so defined that the class K^c is an iterate of K , or dually put K is a Skolem hull of K^c . The proof of its existence initially used a measurable cardinal as well, but eventually R. Jensen and J. Steel found a way around that additional hypotheses.

Theorem 21. *Assume that there is no transitive class model satisfying ZFC and "there is a Woodin cardinal". Then there is a Σ_2 -formula $\Phi(x)$ such that*

$$K := \{x : \Phi(x) \text{ holds}\}$$

is a class model of ZFC, which is iterable. If \mathbb{P} is a notion of forcing of set size and G is a V -generic filter then $K^{V[G]} = K^V$, thus K is absolute for set sized forcing extensions.

Note that as soon as we consider M_1 , cut it at the Woodin cardinal δ we are in the situation of the anti large cardinal assumption and thus can build K in $\mathcal{J}_\delta^{M_1}$. The model we end up with is again $\mathcal{J}_\delta^{M_1}$ which was proved by J.Steel (see [17] for a proof of this). Thus we do have a certain amount of generic absoluteness in M_1 namely there is a formula $\Phi(x)$ which defines K and thus $\mathcal{J}_\delta^{M_1}$ for notions of forcing which have size less than M_1 's Woodin cardinal δ .

We turn now to condensation of sufficiently iterable premice. The central result is the following which was proved by J. Steel and I. Neeman building on the works of Jensen, S.Friedman, Dodd, Mitchell and Schimmerling:

Theorem 22. *Let \mathcal{M} be an ω -sound, $(\omega, \omega_1, \omega_1 + 1)$ -iterable premouse. Let $\pi : \mathcal{H} \rightarrow \mathcal{M}$ be a fully elementary map such that its critical point $\text{cp}(\pi) = \rho_\omega^{\mathcal{H}}$. Then either*

1. \mathcal{H} is a proper initial segment of \mathcal{M}
2. there is an extender E on the \mathcal{M} -sequence such that $\text{lh}(E) = \rho_\omega^{\mathcal{H}}$ and \mathcal{H} is a proper initial segment of $\text{Ult}_0(\mathcal{M}, E)$

A useful observation is the following:

Lemma 23. *Let \mathcal{M} be as in the theorem above. Consider the set S of countable elementary submodels of \mathcal{M} whose transitive collapses are proper initial segments of \mathcal{M} . Then S is stationary in $[\mathcal{M}]^\omega$.*

Proof. Assume for a contradiction that S is not stationary then there is a club C which is $<$ -least in the definable wellorder of the mouse, such that every element H of it does not collapse to an initial segment of \mathcal{M} . Let $\mathcal{N} \triangleright \mathcal{M}$ be a bigger ω -sound, $(\omega, \omega_1, \omega_1 + 1)$ -iterable premouse which sees that C is the $<_{\mathcal{N}}$ -least such club. Let $P \prec_1 \mathcal{N}$ be a countable Σ_1 -substructure which is ω -sound, created as the Σ_1 -hull of $\{\mathcal{M}\}$ in \mathcal{N} . Then $C \in P$ by elementarity and $P \cap [\mathcal{M}]^\omega \in C$ as C is closed and unbounded. But now the collapse of $P \cap [\mathcal{M}]^\omega$ is an initial segment of $\pi(P)$. Yet $\pi(P)$ does collapse nicely as a comparison argument shows. Once we compare $\pi(P)$ with an ω -sound, ω -projecting initial segment of \mathcal{M} nothing will move during the comparison process and so $\pi(P)$ is an initial segment of \mathcal{M} . Thus $\pi(P) \cap [\mathcal{M}]^\omega$ is an initial segment, yet $\pi(P) \cap [\mathcal{M}]^\omega \in C$ which is a contradiction. □

To summarize the above results we formulate an explicit list of axioms which M_1 satisfies and which a reader who does not know about inner model theory should keep in mind when reading this thesis.

Fact 24. *M_1 satisfies the following list of axioms*

1. *There is a Π_2^1 -definable set of reals I whose elements are codes for countable initial segments of M_1 . Moreover these codes are cofinal meaning that for every countable initial segment \mathcal{P} of M_1 there is a code c in I such that \mathcal{N} is an initial segment of c . Further this set still works in all ω_1 -preserving forcing extensions $M_1[G]$ of M_1 .*
2. *In M_1 below its Woodin cardinal Steel's core model K can be constructed and coincides with $\mathcal{J}_\delta^{M_1}$. The definition of K is generically absolute for set forcing extensions of size less than δ .*
3. *If $\mathcal{J}_\eta^{M_1}$ is an initial segment of M_1 , hence an ω -sound, $(\omega, \omega_1, \omega_1 + 1)$ -iterable premouse then there is a stationary subset $S \subset [\mathcal{J}_\eta^{M_1}]^\omega$ such that every $H \in S$ condenses to an initial segment of $\mathcal{J}_\eta^{M_1}$.*

1.3 How to make NS_{ω_1} \aleph_2 -saturated

The investigation of the nonstationary ideal on a regular cardinal has a long history in set theory which is no wonder as stationarity represents one of its most fundamental notions. The question of the length of antichains of stationary subsets modulo nonstationarity generated particular interest due to their central role in generic ultrapower arguments.

Definition 25. *Let κ be a regular cardinal and I an ideal on κ . For a regular cardinal λ we say that I is λ saturated if there are no antichains of length λ in $P(\kappa)\setminus I$, where antichains are meant to be modulo I -small intersections of their elements.*

An equivalent way of saying that I is λ -saturated is therefore the statement that the Boolean algebra $P(\kappa)\setminus I$ has the λ -cc, which highlights the importance of the notion in the context of generic ultrapowers where conditions are elements of I -positive sets ordered by the subset relation.

The question of possible values for antichains modulo I is intimately tied with large cardinals as was already implicitly evident in S. Ulam's work on measurability, showing that for no λ there is a λ^+ -saturated ideal on λ^+ . Forty years later R. Solovay showed that for a regular uncountable cardinal κ , every stationary $S \subset \kappa$ can be partitioned into κ many stationary sets which implies that for every stationary $S \subset \kappa$ the restricted nonstationary ideal $\text{NS}_\kappa \upharpoonright S$ can not be κ -saturated. Thus naturally the question arises whether there are successor cardinals κ such that the nonstationary ideal on κ is κ^+ -saturated. Here a crucial difference between the nonstationary ideal on \aleph_1 and the nonstationary ideal on other regular $\kappa > \aleph_1$ shows up, which has its deeper reasons in the trivial fact that below ω_1 limit ordinals have only one possible cofinality, while at bigger cardinals more possibilities occur. It is a ZFC-theorem of M.Gitik and S.Shelah that NS_κ can not be κ^+ -saturated for any $\kappa > \omega_1$, for $\kappa = \omega_1$ however, NS_{ω_1} can be \aleph_2 -saturated, which was shown first by K. Kunen assuming the existence of a huge cardinal.

The ultimate solution to the problem of the consistency of the statement NS_{ω_1} is \aleph_2 -saturated from optimal large cardinal assumptions was eventually found by S. Shelah who showed around 1985 that already a Woodin cardinal suffices for the consistency of NS_{ω_1} being saturated. As this result and its proof are essential for our work we will give the proof in this section in detail. As a concluding remark we mention that it was only till 2006 that R.Jensen and J.Steel proved that the assumption of a Woodin cardinal is in fact sharp in terms of consistency strength via showing that if the theory $\text{ZFC} + \text{“NS}_{\omega_1}$ on ω_1 is saturated” is consistent then so is $\text{ZFC} + \text{“there is a Woodin cardinal”}$

We start with the preparations for Shelah's result. The material in this section draws heavily from R.Schindler's notes [16] on the problem. Recall first a couple of definitions and facts.

Definition 26. Let A be an arbitrary set then a cardinal κ is A -strong up to the cardinal δ iff $\forall \gamma < \delta \exists j : V \rightarrow M$ which is elementary such that

1. $\text{crit } j = \kappa \wedge \gamma < j(\kappa)$
2. $V_{\kappa+\gamma} \subset M$
3. $A \cap V_{\kappa+\gamma} = j(A) \cap V_{\kappa+\gamma}$

The following fact can be used to define Woodin cardinals:

Fact 27. The following are equivalent

- δ is Woodin
- For any $A \subset V_\delta$,

$$\{\alpha < \delta : \alpha \text{ is } A\text{-strong up to } \delta \}$$

is stationary in δ .

We will need a bit more, namely a Woodin cardinal with a \diamond -sequence living below it:

Definition 28. Let δ be a Woodin cardinal then we say that δ is Woodin with \diamond iff there is a sequence $(a_\kappa : \kappa < \delta)$ such that for each κ , $a_\kappa \subset V_\kappa$ and for every $A \subset V_\delta$ the set

$$\{\kappa < \delta : A \cap V_\kappa = a_\kappa \wedge \kappa \text{ is } A\text{-strong up to } \delta\}$$

is stationary in δ .

In terms of consistency strength this adds nothing to being a Woodin cardinal. If δ is the Woodin cardinal and we force with δ -Cohen forcing then in the resulting generic extension we have that δ is Woodin with \diamond : indeed first note that δ -Cohen forcing is the same as forcing with conditions of the form $(a_\alpha : \alpha < \kappa < \delta \wedge a_\alpha \subset V_\alpha)$, ordered by end extension. Assume now that there is a condition p such that

$$p \Vdash \tau \subset V_\delta \wedge \sigma \subset \delta \text{ is club in } \delta.$$

We have to show that there is a stronger condition $q < p$, $q = (a_\alpha : \alpha < \lambda)$ and a cardinal $\kappa < \delta$ for which

$$q \Vdash \kappa \in \sigma \text{ is } \tau\text{-strong up to } \delta \wedge \tau \cap \kappa = a_\kappa.$$

We construct by recursion a descending sequence of conditions $(p_\kappa : \kappa < \delta)$ such that the length of each p_κ is μ_κ and such that the following points are obeyed:

1. $\{\mu_\kappa : \kappa < \delta\}$ is a club in δ .
2. For every κ there is some $C_\kappa \subset \mu_\kappa$ which is unbounded in μ_κ such that $p_\kappa \Vdash \sigma \cap \mu_\kappa = C_\kappa$ and consequently $p_\kappa \Vdash \mu_\kappa \in \sigma$.
3. For every κ there is some $A_\kappa \subset V_\kappa$ such that $p_\kappa \Vdash \tau \cap V_{\mu_\kappa} = A_\kappa$
4. for every κ , $a_{\mu_\kappa} = A_\kappa$
5. If $p_{\kappa+1} \not\Vdash \kappa$ is τ -strong up to δ , then there is an $\alpha < \mu_{\kappa+1}$ such that

$$p_{\kappa+1} \Vdash \kappa \text{ is not } \tau\text{-strong up to } \alpha$$

(Note here that $\kappa + 1$ makes such a choice always possible. It is impossible for κ limit)

That such a sequence exists is easily seen. Now set $A := \bigcup_{\kappa < \delta} A_\kappa$, and as δ is Woodin and the set $\{\mu_\kappa : \kappa < \delta\}$ is a club in δ we find a point $\kappa = \mu_\kappa$ such that κ is A -strong up to δ . Now pick the condition $q := (a_\lambda : \lambda < \kappa + 1)$ which by 2,3 and 4 satisfies

$$q \Vdash \kappa \in \sigma \wedge \tau \cap \kappa = a_\kappa.$$

To finish we need to show that also $q \Vdash \kappa$ is τ -strong up to δ . Assume for a contradiction the opposite then by property 5 there exists an $\alpha < \mu_{\kappa+1}$ such that

$$p_{\kappa+1} \Vdash \kappa \text{ is not } \alpha\text{-strong up to } \delta.$$

But the elementary embedding $j : V \rightarrow M$ which witnesses that κ is A -strong up to δ can be lifted to $j' : V[G] \rightarrow M'$, and by the δ -closure of the forcing it will witness that κ is A -strong up to δ , a contradiction.

The usage of the \diamond -sequence at the Woodin cardinal is crucial for the proof of the existence of a model where NS_{ω_1} is \aleph_2 -saturated. We will use it as a guideline for the iteration. Whenever we hit a stage α such that the \diamond -sequence $(a_\beta : \beta < \delta)$ at stage α is the name of a maximal antichain of stationary subsets of ω_1 of length ω_2 we want to change its length to \aleph_1 , and further ensure that this maximal antichain remains maximal in all stationary subsets of ω_1 -preserving outer models. The next forcing notion does exactly what we demand:

Definition 29. Assume that \vec{S} is an antichain of stationary subsets of ω_1 . Then the so called sealing forcing $\mathbb{S}(\vec{S})$ consists of conditions of the form (p, c) where $p : \alpha + 1 \rightarrow \vec{S}$ is a function and $c : \alpha + 1 \rightarrow \omega_1$ is a function with closed image and such that

$$\forall \xi \leq \alpha (c(\xi) \in \bigcup_{i \in \xi} p(i))$$

holds. We let $(q, d) < (p, c)$ if q and d end-extend p and c respectively.

It is well known that the sealing forcing $\mathbb{S}(\vec{S})$ is ω -distributive and preserves all stationary subsets of \vec{S} , thus $\mathbb{S}(\vec{S})$ is stationary subsets of ω_1 preserving if \vec{S} is maximal.

We can also consider a stationary, co-stationary set $A \subset \omega_1$ and the nonstationary ideal restricted to subsets of A ,

$$\text{NS}_{\omega_1} \upharpoonright A := \{B \subset A : B \text{ is stationary}\}.$$

It is natural to ask the same question we asked for the full ideal NS_{ω_1} , for the restricted version $\text{NS}_{\omega_1} \upharpoonright A$, namely whether it can be \aleph_2 -saturated. Surprisingly its positive answer has a simpler structure than for full NS_{ω_1} . The reason behind this is that the set A enables us to see that the according A -sealing forcings are particularly nice, namely ω -distributive.

Definition 30. *Assume that $A \subset \omega_1$ is a stationary, co-stationary subset of ω_1 and that \vec{S}_A is an antichain of stationary subsets of A . Then we can also seal it off using the straightforward generalization, denoted by $\mathbb{S}(\vec{S})_A$ of the already introduced sealing forcing $\mathbb{S}(\vec{S})$. Conditions of $\mathbb{S}(\vec{S})_A$ are pairs (p, c) where $p : \alpha + 1 \rightarrow \vec{S}_A$ is a function and $c : \alpha + 1 \rightarrow \omega_1$ is a function with range a closed subset of ω_1 such that*

$$\forall \xi \leq \alpha (c(\xi) \in A \rightarrow c(\xi) \in \bigcup_{I \in \xi} p(I))$$

The idea now to force NS_{ω_1} to be \aleph_2 -saturated is to seal off one by one all the long antichains of stationary subsets with an iteration. The first obstacle one immediately encounters is how we catch our tail during this process, for which the usage of the Woodin cardinal will be crucial. Another difficulty is the following: One has to ensure that the iteration stays stationary set preserving to avoid utter chaos. The sealing forcing $\mathbb{S}(\vec{S})$ is stationary set preserving as long as \vec{S} is maximal, but there does not exist a theory for iterations of stationary set preserving forcings in a stationary set preserving way. Thus we are compelled to work with semiproper notions of forcing and *RCS*-iterations instead. We demand to only seal off a maximal antichain \vec{S} when the forcing $\mathbb{S}(\vec{S})$ is also semiproper. But this leaves us with the possibility of not sealing off every long antichain during the iteration. That these difficulties still do not prevent the proof is outlined here in detail:

Theorem 31. *Assume that δ is a Woodin cardinal with \diamond . Then there exists a semiproper forcing \mathbb{P} of size δ such that in $V[G]$, NS_{ω_1} is \aleph_2 -saturated and $\delta = \aleph_2$.*

Proof. We use a \diamond -sequence on δ to determine at each stage $\alpha < \delta$ which forcing to use. Assume that during our iteration we have arrived at stage α . Then let \mathbb{Q}_α be

1. the sealing off forcing of a maximal antichain σ^{G_α} if the diamond sequence at stage α is the \mathbb{P}_α -name σ for a maximal antichain of stationary subsets of ω_1 and the sealing forcing is semiproper.
2. the collapse of 2^{\aleph_2} to \aleph_1 else.

At limit stages we use the *RCS*-limit. This already suffices. Assume for a contradiction that NS_{ω_1} is not \aleph_2 -saturated in $V[G]$, I.e. there is a maximal antichain $\vec{S} = (S_i : I < \omega_2)$ in $P(\omega_1)/\text{NS}_{\omega_1}$. Let τ be a \mathbb{P} -name for the sequence. As $V[G] \models \aleph_2 = \delta$ for our Woodin cardinal δ , we claim that it is possible to find an inaccessible κ below δ such that the following three properties hold:

1. κ is $\mathbb{P} \oplus \tau$ -strong up to δ in V
2. $\kappa = \omega_2^{V[G \upharpoonright \kappa]}$
3. $\vec{S} \upharpoonright \kappa = (S_i : I < \kappa) = (\tau \cap V_\kappa)^{G \upharpoonright \kappa}$ is the maximal antichain in $V[G \upharpoonright \kappa]$ which is picked by the \diamond -sequence at stage κ .

This is clear as we can assume that our \diamond -sequence lives on the stationary subset of inaccessible cardinals below δ , and for all inaccessible κ property 2 automatically holds. Moreover the sets

$$C_1 := \{\kappa < \delta : \vec{S} \upharpoonright \kappa = (S_i : I < \kappa) = (\tau \cap V_\kappa)^{V[G \upharpoonright \kappa]}\}$$

and

$$C_2 := \{\kappa < \delta : \forall \alpha \forall S \in P(\omega_1) \cap V^{\mathbb{P}^\alpha} \text{ stationary } \exists \bar{S} \in \vec{S} \upharpoonright \kappa (S \cap \bar{S} \notin \text{NS})\}$$

are both clubs, therefore hitting the stationary set T consisting of the points $\kappa < \delta$ where $\tau \cap V_\kappa = a_\kappa$ (remember: a_κ is the κ -th element of the \diamond -sequence $(a_\alpha : \alpha < \delta)$) and κ is τ -strong up to δ . Thus if κ is in the nonempty intersection $C_1 \cap C_2 \cap T$ then 1 and 2 are satisfied, and the recursive definition of our forcing \mathbb{P} yields that at stage κ , as $a_\kappa = \tau \cap V_\kappa$, the sealing forcing $\mathbb{S}((\tau \cap V_\kappa)^{G \upharpoonright \kappa})$ is at least considered, and in order to show property 3, it suffices to show that $(\tau \cap V_\kappa)^{G \upharpoonright \kappa} = \vec{S} \upharpoonright \kappa$ is maximal in $V[G \upharpoonright \kappa]$. But this is clear as by the definition of RCS iteration and as $|\mathbb{P}_\alpha| < \kappa$ we take at inaccessible κ 's the direct limit of the \mathbb{P}_α 's, thus each stationary $S \subset \omega_1$ in $V^{\mathbb{P}^\kappa}$ is already included in a $V^{\mathbb{P}^\alpha}$ for $\alpha < \kappa$. So we have ensured the existence of a κ with all the 3, above stated properties.

Now the forcing $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ can not be semiproper at stage κ , as otherwise we would have to force with it, therefore killing the antichain \vec{S} . So there exists a condition $(p, c) \in \mathbb{S}(\vec{S} \upharpoonright \kappa)$ such that the set

$$\bar{T} := \{X \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} : |X| = \aleph_0 \wedge (p, c) \in X \wedge \nexists Y \supset X (Y \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \wedge |Y| = \aleph_0 \wedge (X \cap \omega_1 = Y \cap \omega_1) \wedge \exists (q, d) \leq (p, c) ((q, d) \text{ is } Y\text{-semigeneric}))\}.$$

is stationary in $V[G \upharpoonright \kappa]$, and by construction of our iteration, the κ -th forcing in \mathbb{P} is $Col(\omega_1, 2^{\aleph_2})$, so in $V[G \upharpoonright \kappa + 1]$ there is a surjection $f : \omega_1 \rightarrow (H_{\kappa+})^{V[G \upharpoonright \kappa]}$. As $Col(\omega_1, 2^{\aleph_2})$ is proper the set \bar{T} remains stationary in $V[G \upharpoonright \kappa + 1]$ which implies that

$$T := \{\alpha < \omega_1 : f''\alpha \in \bar{T} \wedge \alpha = f''\alpha \cap \omega_1\}$$

is stationary in $V[G \upharpoonright \kappa + 1]$. As the tail $\mathbb{P}_{[\kappa+2, \delta)}$ remains semiproper, seen as an iteration with $V[G \upharpoonright \kappa + 1]$ as ground model, we can infer that T remains stationary in $V[G]$ and hence there exists an $i_0 < \delta$ such that

$$(**) \quad T \cap S_{i_0} \text{ is stationary in } V[G].$$

Let us shortly reflect the situation we are in. The idea is to find a model $X \in \bar{T}$ such that we *can* find a $(X, \mathbb{S}(\vec{S} \upharpoonright \kappa))$ -semigeneric condition $(q, d) < (p, c)$, thus arriving at a contradiction. In order to do so we have to ensure that $\alpha = X \cap \omega_1$ is in some $S_i \in \vec{S} \upharpoonright \kappa$. As \vec{S} was assumed to be maximal there is indeed an index $i_0 < \delta$ which is as desired, this index however might be bigger than κ . This is where the large cardinal assumption comes into play. We can find an elementary embedding $j : V \rightarrow M$ such that $j(\kappa) > i_0$, thus it seems that $j(\vec{S} \upharpoonright \kappa)$ is now long enough to have S_{i_0} as an element. But this is not correct as \vec{S} was not assumed to be definable and therefore $j\vec{S} \upharpoonright \kappa \neq \vec{S} \upharpoonright j(\kappa)$. We have to use more than just the elementary embedding, namely that Woodiness fixes even a predicate with j . Indeed if we let $\lambda > i_0$ such that $(\tau \cap V_\lambda)^{G \upharpoonright \lambda} = \vec{S} \upharpoonright \lambda$, and let $j : V \rightarrow M$ be such that $j(\tau) \cap V_\lambda = \tau \cap V_\lambda$ then $j(\tau \cap V_\kappa) = j(\tau) \cap V_{j(\kappa)}$ and $V_{j(\kappa)} = V_\lambda \cup V_{j(\kappa)-V_\lambda}$, thus $j(\vec{S} \upharpoonright \kappa) = j(\tau \cap V_\kappa)^{G \upharpoonright \kappa}$ contains $(\tau \cap V_\lambda)$ and thus S_{i_0} .

First let $\lambda < \delta$, $\lambda > \max(i_0, \kappa + 1)$ be such that $(\tau \cap V_\lambda)^{G \upharpoonright \lambda} = \vec{S} \upharpoonright \lambda$, so we have $(\tau \cap V_\lambda)^{G \upharpoonright \lambda}(i_0) = S_{i_0}$. As κ was chosen to be $\mathbb{P} \oplus \tau$ -strong up to δ we let $j : V \rightarrow M$ be an elementary embedding with critical point κ , such that M is transitive, $M^\kappa \subset M$, $V_{\lambda+\omega} \subset M$, $j(\mathbb{P}) \cap V_\lambda = \mathbb{P} \cap V_\lambda$, and $j(\tau) \cap V_\lambda = \tau \cap V_\lambda$.

H should denote the generic filter for the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ over $M[G \upharpoonright \lambda]$. Then we lift j to an elementary embedding

$$j^* : V[G \upharpoonright \kappa] \rightarrow M[G \upharpoonright \lambda, H].$$

Notice that $(V_{\lambda+\omega})^{V[G \upharpoonright \lambda]} = (V_{\lambda+\omega})^{M[G \upharpoonright \lambda]}$.

Now we let $(X_i : I < \omega_1) \in V[G \upharpoonright \kappa + 1]$ be an increasing continuous chain of countable elementary substructures of $(H_{j(\kappa+)}^{M[G \upharpoonright \kappa+1]})$ with $\{\tau \cap V_\lambda, i_0\} \subset X_0$ satisfying for all $I < \omega_1$ the following three properties:

- (a) $I \in X_{I+1}$
- (b) $f''(X_i \cap \omega_1) \subset X_i$
- (c) $j^{*''}(X_i \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]}) \subset X_i$

Let $\bar{G} := G \upharpoonright [\kappa + 2, \lambda]$, then we have that

$$\{X_i[\bar{G}] \cap \omega_1 : I < \omega_1\} \in V[G \upharpoonright \lambda]$$

is a club in ω_1 so intersecting it with the stationary set defined in (**) we find some $I < \omega_1$ such that $X_i[\bar{G}] \cap \omega_1 = X_i \cap \omega_1 \in T \cap S_{i_0}$.

Write $X := X_i, \alpha := X \cap \omega_1$. As at stage κ we had to force with the ω -closed $Col(2^{\aleph_2}, \aleph_1)$ we know that $X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]} \in V[G \upharpoonright \kappa]$. Remember that $f \in V[G \upharpoonright \kappa + 1]$ was chosen as a surjection of ω_1 onto $(H_{\kappa+})^{V[G \upharpoonright \kappa]}$, so as $\alpha \in T$ by definition of T $f''\alpha \in \bar{T}$ and $\alpha = f''\alpha \cap \omega_1$, and hence by (b)

$$f''\alpha \subset X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]} \in V[G \upharpoonright \kappa].$$

As $\alpha = f''\alpha \cap \omega_1$, $f''\alpha \in \bar{T}$ and $f''\alpha \subset X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]}$ we get that $X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]} \in \bar{T}$ and therefore

$$(***) \quad j^*(X \cap H_{\kappa+})^{V[G \upharpoonright \kappa]} \in j^*(\bar{T}).$$

Note that our second generic H , denoting the generic filter for the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ over $M[G \upharpoonright \lambda]$ has not been specified yet. As the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ over $M[G \upharpoonright \lambda]$ is semi-proper we have that there is a condition q in the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ which is $(X[G], \mathbb{P}_{[\lambda+1, j(\kappa)]})$ -semigeneric. If we pick H such that $q \in H$ then by semigenericity of q we obtain $X[\bar{G}, H] \cap \omega_1 = X[\bar{G}] \cap \omega_1 = X \cap \omega_1 = \alpha \in S_{i_0} = (\tau \cap V_\lambda)^{G \upharpoonright \lambda}(i_0) \in X[\bar{G}, H]$. But also due to (c) we have that

$$j^*(X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]}) = j''(X \cap (H_{\kappa+})^{V[G \upharpoonright \kappa]} \subset X[\bar{G}, H].$$

This gives us the desired contradiction as we can find an $(X[\bar{G}, H], j(\mathbb{S}(\vec{S} \upharpoonright \kappa)))$ -semigeneric condition below $j(p, c) = (p, c)$. Indeed we can just list the countably many names for countable ordinals in $X[\bar{G}, H]$ along with conditions of $j(\mathbb{S}(\vec{S} \upharpoonright \kappa))$ deciding them below (p, c) and let $(p', c') \in j(\mathbb{S}(\vec{S} \upharpoonright \kappa))$ be just the condition with $\text{dom}(c') = \text{dom}(d') = \alpha + 1$, $c'(\alpha) = \alpha$ and $p'(I) = S_{i_0}$ for some $I < \alpha$. So $X[\bar{G}, H]$ together with $(p', c') < (p, c)$ witness that $j^*(X \cap H_{\kappa+})^{V[G \upharpoonright \kappa]} \notin j^*(\bar{T})$, contradicting (***) □

Chapter 2

$\text{NS}_{\omega_1} \upharpoonright A$ is Δ_1 -definable

2.1 Introduction

In this section we want to give a proof of the following theorem

Theorem 32. *Assume that $M_1^\#$ exists, and let $A \in M_1$ be a stationary, co-stationary subset of ω_1 . Then there is a generic extension $M_1[G]$ via a set sized forcing such that in $M_1[G]$ NS_{ω_1} is Δ_1 -definable in $H(\omega_3)$ with parameter ω_1 and such that $\text{NS}_{\omega_1} \upharpoonright A$ is \aleph_2 -saturated.*

Before starting the actual proof we outline its route roughly: The main idea is to use Shelah's proof of the saturation of NS_{ω_1} from a Woodin cardinal as a starting point and try to add certain forcings which code additional information of the universe. This additional information can be used to obtain a nicer definition of stationarity on ω_1 . The extra coding forcings should be chosen in such a way that they will not interfere with the usual sealing forcings which push the saturation of NS_{ω_1} down to \aleph_2 , and should be robust enough in that a once coded information should be preserved in all future further generic extensions of the universe.

To be a little more precise the proof is via a δ -long (δ the Woodin cardinal) countable support iteration of semiproper and \mathcal{S} -proper forcings. We will use a \diamond -sequence $(a_\alpha)_{\alpha < \delta}$ to guide the iteration. At each stage α of the iteration we look at the α -th entry of the sequence. We will distinguish two different cases, a coding stage and a sealing stage and start to describe the first of the two stages:

If a_α is the \mathbb{P}_α -name of a stationary subset $S \subset A$ then we code the characteristic function of S into a pattern of nicely definable trees which should have a cofinal branch or be bounded, meaning in that particular context that there are no cofinal branches through the tree in no ω_1 -preserving outer model. This is followed by a bunch of forcings which add a more local version of this information to the universe. We shall see that the iteration of these forcings is \mathcal{S} -proper.

If a_α is a \mathbb{P}_α -name of a long (i.e. of length \aleph_2) maximal antichain in the structure $P(A)/\text{NS}_{\omega_1}$ then we seal off this antichain after collapsing its length to \aleph_1 but only if this forcing is semiproper. This seemingly redundant move will enable us to argue for stationary set preservation for the resulting iteration, as, contrary to the semiproper case, there is no iteration theory of stationary set preserving notions of forcing. This forcing is seen to be A^c -proper and semiproper. In the remaining case we just do nothing and force with the trivial forcing.

To summarize we arrive at a countable support iteration of \mathcal{S} -proper and A^c -proper forcings, therefore the iteration is $\mathcal{S}(A^c)$ -proper where $\mathcal{S}(A^c)$ should denote the class $\{M \in \mathcal{S} : M \cap \omega_1 \in A^c\}$. We will see soon that the class \mathcal{S} is projective stationary, i.e. $\mathcal{S}(A^c)$ of models remains everywhere stationary, hence the iteration with countable support yields a $\mathcal{S}(A^c)$ -proper extension of the ground model M_1 . Consequently all new countable sets of ordinals can be covered by countable sets of ordinals living in the ground model. We use this observation to argue that stationary subsets of ω_1 are preserved in the just described iteration:

We can look at the iteration in a different way. As we only seal off when the sealing forcing is semiproper, and as the forcing from coding stages is \mathcal{S} -proper (as to be seen soon), hence \mathcal{S} -semiproper we drop the countable support iteration for a second and use an *RCS*-iteration instead, i.e. we use all the factors of the iteration but iterate them using the revised countable support instead of the plain countable support. We arrive at a \mathcal{S} -semiproper generic extension of the ground model M_1 which is stationary set preserving as \mathcal{S} is projective stationary. Yet, as $\mathcal{S}(A^c) = \{M \in \mathcal{S} : M \cap \omega_1 \in A^c\}$ remains everywhere stationary, countable subsets of ordinals in the generic extension can be covered by countable sets in the ground model. Consequently the *RCS*-iteration is in fact just a countable support iteration. Indeed at every stage $\alpha < \delta$ we see that $V^{\mathbb{P}_\alpha}$ is a $\mathcal{S}(A^c)$ -proper forcing extension of V , thus whenever there is an ordinal $\beta > \alpha$ such that $\text{cof}(\beta)$ is countable in $V^{\mathbb{P}_\alpha}$, it has already been of countable cofinality in V . Thus the *RCS*-iteration is just a countable support iteration, and we can iterate with countable support to end up with a \mathcal{S} -semiproper extension of the ground model. As a consequence the final model, when using countable support iteration for the iterands, preserves stationary subsets of ω_1 .

2.2 The coding forcing

As already mentioned we need a coding forcing which harmonizes with the stationary sealing forcings we need to keep the saturation of NS_{ω_1} low at \aleph_2 . Our ground model for the iteration will be M_1 , the canonical inner model with one Woodin cardinal. We cut M_1 at the Woodin cardinal δ to build Steel's K there. We have already seen that $K^{\mathcal{J}_\delta^{M_1}} = \mathcal{J}_\delta^{M_1}$. The move

towards K has the advantage that K has a nice first order definition, as opposed to M_1 , and generic absoluteness below δ is a well known fact for K . We use K -trees $T_\alpha(\xi) := ((\alpha^{+\xi})^{<(\alpha^{+\xi})})^K$, where β is a cardinal from K , and either force a cofinal branch through it or we ensure that there can not be any cofinal branches in ω_1 -preserving outer models. This way we can code arbitrary 0,1-patterns into sequences of cofinal or bounded trees. The big advantage of this method is that once we decided to write a certain pattern using these trees this information will prevail in all ω_1 -preserving outer models. Thus we don't have to reconsider earlier information during the iteration. These sequences of cofinal or bounded trees will code characteristic functions of stationary subsets of ω_1 . Of course the process of specializing a tree off or shooting a cofinal branch should preserve stationary subsets of ω_1 to be useful for our purpose. Moreover we should be able to iterate these forcings in a nice way. That this is indeed the case is the content of the next proposition.

Proposition 33 (Coding via specializing trees). *Assume GCH, $\beta > \omega_1$ regular, and let \mathcal{S} be a stationary class which is projective stationary, i.e. for every stationary $S \subset \omega_1$, $\mathcal{S}(S) = \{M \in \mathcal{S} : M \cap \omega_1 \in S\}$ is everywhere stationary. Suppose that \mathbb{Q} is an \mathcal{S} -proper notion of forcing of size less than β and G is V -generic. Then:*

1. $T(\beta) := ((\beta^+)^{<(\beta)})^V$ viewed as a forcing is \mathcal{S} -proper over $V[G]$.
2. There is a proper forcing $\mathbb{R} \in V[G]$ of size β^{++} that destroys the properness of $T(\beta)$. More specifically if H is \mathbb{R} -generic over $V[G]$, then in any ω_1 preserving outer model of $V[G][H]$ there is no branch through $T(\beta)$ which is $T(\beta)$ -generic over V .

Proof. We shall show first that $T(\beta) := ((\beta^+)^{<(\beta)})^V$ is \mathcal{S} -proper over $V[G]$ for G a generic filter for the forcing \mathbb{Q} . This is trivial as the \mathcal{S} -proper forcing \mathbb{Q} will not change the stationarity of \mathcal{S} in $V^{\mathbb{Q}}$. Thus if we pick any countable $M \in \mathcal{S} \subset V$ and fix a condition $p \in T(\beta) \cap M$, we can find a stronger $q < p$ which is $(M, T(\beta))$ -generic, as $T(\beta)$ is defined in V and sufficiently closed there. So $T(\beta)^V$ is \mathcal{S} -proper in $V[G]$.

To prove the second statement, first add β^{++} Cohen reals with a finite support product over $V[G]$, then Lévy collapse β^{++} to ω_1 and let $V[G][H_1][H_2]$ denote the resulting model, which is a proper forcing extension of $V[G]$. By an observation of J.Silver we know that each β -branch through $T(\beta)$ which lies in $V[G][H_1][H_2]$, in fact lies already in $V[G][H_1]$. Indeed if \dot{b} is a name in $V[G][H_1]$ for a new $T(\beta)$ -branch, then we can build a binary ω -tree of conditions in the Lévy collapse and by its ω_1 closure each branch has a lower bound, resulting in $2^{\aleph_0} = \beta^{++}$ -many different interpretations of \dot{b} . Thus $(T(\beta))^V$ has in $V[G][H_1]$ $2^{\aleph_0} = \beta^{++}$ many nodes on a fixed level which is impossible as $V \models GCH$.

Thus $T(\beta)$ has at most ω_1 -many branches in $V[G][H_1][H_2]$, and none of those branches is cofinal in β^+ , therefore none of the branches is $T(\beta)$ -generic over V . Also every node is included in a β -branch. This enables us to use Baumgartner's method of "specializing a tree off a small set of branches".

Fact 34. *If T is tree of height ω_1 , such that each node is contained in a cofinal branch, and with at most \aleph_1 many cofinal branches, then there is a ccc forcing \mathbb{P} such that if G is \mathbb{P} -generic then whenever W is an outer model of $V[G]$ with the same ω_1 then each cofinal branch through W belongs already to V .*

Now we can use the forcing \mathbb{P} from the Fact above to build the model $V[G][H_1][H_2][H_3]$ and obtain that whenever W is an ω_1 preserving outer model of $V[G][H_1][H_2][H_3]$ then each branch through the tree $T(\beta)$ in W is in fact in $V[G][H_1][H_2]$, and therefore as already noted in $V[G][H_1]$. As no branch through $T(\beta)$ in $V[G][H_1]$ is cofinal in β^+ (by the ccc), and $T(\beta)$ generic branches are necessarily cofinal in β^+ , we are done. \square

2.3 The class \mathcal{S}

Note that in the following we constantly use the fact that M_1 below the Woodin cardinal coincides with Steel's core model K as build in $\mathcal{J}_\delta^{M_1}$. The move toward K has the advantage that, contrary to M_1 K admits a first order definition. This does not cause any problems as long as we are below δ (which we are all the time). As M_1 is the ground model of our iteration, we do not have access to full condensation. Knowing that a sufficiently elementary submodel M of some $\mathcal{J}_\eta^{M_1}$ collapses to an M_1 initial segment is nevertheless of highest importance in our proof. The reason for this is lies in the fact that we use K definable trees, living on K cardinals. It is therefore desirable that these trees still live on K initial segments even after we transitively collapse, in order to not completely lose control of the things we are talking about. These considerations will become clearer once the everywhere stationary class \mathcal{S} , introduced for these very reasons is seen in action during the proof.

We introduce without much further ado:

Definition 35. *Let \mathcal{S} be a class of countable sets. We say that \mathcal{S} is everywhere stationary if*

1. *for every regular cardinal θ , $\mathcal{S} \cap H(\theta)$ is a stationary subset of $[H(\theta)]^\omega$, and*
2. *\mathcal{S} is closed under truncation, I.e $X \cap H_\theta \in \mathcal{S}$ whenever $X \in \mathcal{S}$ and θ a regular cardinal.*

Definition 36. Let M be a (sufficiently) elementary submodel of some K initial segment \mathcal{J}_η^K . We say that M collapses nicely if the following demands are met:

1. the transitive collapse \bar{M} is an initial segment of K , I.e. $\bar{M} \triangleleft K$, and
2. whenever $\omega_1^{\bar{M}}$ is a cardinal in an K -initial segment \mathcal{J}_η^K then already all cardinals of \bar{M} remain cardinals in \mathcal{J}_η^K , and
3. \bar{M} is K -correct, meaning that if $\Phi(x)$ denotes the first order formula defining K and use it to define K inside the transitive model \bar{M} then $K^{\bar{M}} = \bar{M}$.

This list seems quite daring on first sight, nevertheless there are plenty of nicely collapsing submodels:

Lemma 37. Let \mathcal{S} denote the class of nicely collapsing submodels of M_1 . Then \mathcal{S} is everywhere stationary in M_1 . Moreover \mathcal{S} is projective stationary, meaning that for every stationary $X \subset \omega_1$, the set $\mathcal{S}(X) := \{M \in \mathcal{S} : M \cap \omega_1 \in X\}$ remains an everywhere stationary class.

Proof. The first item is just the Lemma 23 on Condensation we have proved already. For the second part we have to show that club often $\bar{M} = \mathcal{J}_\eta^{M_1}$ is K -correct, but this is easy as we could have just picked an M which is elementary in the universe with respect to the K defining formula Φ .

\mathcal{S} is trivially closed under truncation and all the considerations so far hold for $H(\theta) \cap M$ as well, thus $H(\theta) \cap M \in \mathcal{C} \cap \mathcal{S}$ and \mathcal{S} is everywhere stationary.

Finally to show that \mathcal{S} is projective stationary we assume the opposite, thus there is a stationary $X \subset \omega_1$ and a fixed regular cardinal θ for which there is a club $C \subset [H_\theta]^\omega$ such that every $M \in C$ does not collapse to an initial segment of M_1 or satisfies $M \cap \omega_1 \notin X$. Let C be the least such club in the wellorder. We pick a regular $\lambda > \theta$ and the Σ_n -Skolem hull denoted by N of $\{X, \theta\}$ in H_λ . We can assume that $N \cap \omega_1 \in X$. It is clear that $N \cap H_\theta \in C$, however we can arrange that N collapses to an initial segment of M_1 and so does $N \cap H_\theta$ which is a contradiction. To see that we can always assume that N collapses to an initial segment of M_1 we first note that N projects to ω as it is not fully elementary in H_λ and can be chosen to be ω -sound. If we pick a countable initial segment of M_1 , say $\mathcal{J}_\eta^{M_1}$ which projects to ω and is ω -sound as well we can start to compare these two models. But now both models will not move during the comparison and so $\bar{N} \triangleleft \mathcal{J}_\eta^{M_1}$ as desired. □

The role of \mathcal{S} and its somewhat peculiar definition will become clearer during the advance of the proof. The reason we have to constantly fall back

on \mathcal{S} is that in the end we want suitable countable models to be able to see the patterns of cofinal or bounded trees we inscribed during the iteration. As these trees are defined in M_1 on M_1 -cardinals, the countable models should be correct about their M_1 and their sequence of M_1 -cardinals. As condensation in general fails in M_1 we are compelled to work with \mathcal{S} .

2.4 The definition of the iteration

We now describe the iteration in detail which is used to prove the main theorem. It utilizes a δ -long countable support iteration of \mathcal{S} -semiproper forcings, the factors of it will be explained in the following subsections. Our ground model is the inner model M_1 with one Woodin cardinal δ , which is also Woodin with \diamond as was shown. Actually we are working all the time below the Woodin cardinal δ , and we emphasize again that $\mathcal{J}_\delta^{M_1} = K$. We will therefore use M_1 and K synonymously from now on and hope that it will not confuse the reader. Remember that in M_1 GCH does hold and there exists a well-order, denoted by $<$ of M_1 . Fix a stationary, co-stationary subset $A \subset \omega_1$. The goal is to arrange the sealing stages, where we seal a maximal antichain of stationary subsets of ω_1 off and the coding stages, where we code the characteristic function χ_S of a stationary $S \subset \omega_1$ into an according pattern of cofinal and bounded (i.e. trees which have no branch of the height of the tree) trees in such a way that they will not interfere with each other.

We use the \diamond -sequence $(a_\alpha)_{\alpha < \delta}$ to determine with which forcing we should force at stage α . Thus assume that $\alpha < \delta$ and we have already constructed \mathbb{P}_β for $\beta \leq \alpha$. We define the forcing $\dot{\mathbb{Q}}_\alpha$ in $V^{\mathbb{P}_\alpha}$ as follows:

1. if a_α is a \mathbb{P}_α -name of a stationary subset S of ω_1 then we code the characteristic function of S into a pattern of trees, followed by forcings which will localize the inscribed information. These forcings will be specified below in a detailed description.
2. if a_α is a \mathbb{P}_α -name of a maximal antichain of stationary subsets of our fixed stationary, co-stationary $A \subset \omega_1$, then use the sealing forcing to seal it off, provided the sealing forcing is \mathcal{S} -semiproper. If it is not force with the usual Levy collapse $Col(\aleph_1, 2^{\aleph_2})$.
3. else collapse 2^{\aleph_2} to \aleph_1 or force to create a default pattern of cofinal trees

The third point of the definition deserves an explanation: our goal is that we only create ω_1 -length patterns of cofinal and bounded trees which correspond to characteristic functions of stationary subsets of ω_1 . We allow a default pattern however to fill in gaps in the sequence of patterns. The reason that we do so is to avoid the case of accidentally forced patterns during our

iteration. Indeed if there is a gap in between two characteristic functions of stationary sets, it could happen that the noise of the forcings we use create an unwanted pattern of trees in the gap, which we cannot kill off once it is produced. This potentially ruins our argument. To avoid this degenerated case we set up the iteration in such a way that a default pattern is forced in all gaps between characteristic functions of stationary sets. This is cumbersome to explain in the recursive way we chose above, therefore we used the blurry words in the third case and hope that this remark makes the point clear.

2.5 The sealing forcings

Whenever a_α at stage α is the \mathbb{P}_α -name of a long maximal antichain in $P(A)/\text{NS}_{\omega_1}$ then we have to seal it off using the following notion of forcing:

Definition 38. *Assume that \vec{S}_A is a maximal antichain of stationary subsets of A , where $A \subset \omega_1$ is stationary, co-stationary. Conditions of the sealing forcing $\mathbb{S}(\vec{S})_A$ are pairs (p, c) where $p : \alpha + 1 \rightarrow \vec{S}_A$ is a function and $c : \alpha + 1 \rightarrow \omega_1$ is a function with range a closed subset of ω_1 such that*

$$\forall \xi \leq \alpha (c(\xi) \in A \rightarrow c(\xi) \in \bigcup_{i \in \xi} p(i))$$

It is well known that $\mathbb{S}(\vec{S})_A$ is ω -distributive, preserves stationary subsets of ω_1 but is not necessarily semiproper. The definition of $\mathbb{S}(\vec{S})_A$ still makes sense if the antichain \vec{S}_A is not maximal. In that case forcing with $\mathbb{S}(\vec{S})_A$ turns the antichain into a maximal one. If A is not stationary then the definition of $\mathbb{S}(\vec{S})_A$ is again meaningful and the forcing $\mathbb{S}(\vec{S})_A$ has a dense subset which is countably closed.

2.6 The coding forcings

We turn now to the forcings mentioned in case 1. They are defined as a three step iteration of forcings $\mathbb{Q}_\alpha^0 * \mathbb{Q}_\alpha^1 * \mathbb{Q}_\alpha^2$ which will be defined now:

2.6.1 \mathbb{Q}_α^0

For brevity we use in the following often the notion $(K)_\eta$ for an ordinal η which just means \mathcal{J}_η^K . Assume we arrived in our iteration at stage α , α a cardinal of the ground model K which is M_1 below the Woodin cardinal δ , and the \diamond -sequence $(a_\beta)_{\beta < \delta}$ at stage α is the \mathbb{P}_α -name of a stationary subset $S \subset \omega_1$. Then we want to write the characteristic function χ_S of S into a pattern of canonical trees for which we either shoot a cofinal branch through, or make sure that there will never be a cofinal branch in each ω_1 -preserving outer model.

More specifically we let \mathbb{Q}_α^0 be an ω_1 -length iteration of forcings $(\mathbb{P}_\xi : \xi < \omega_1)$ defined as follows: For $\xi < \omega_1$ we consider the forcing $T_\alpha(\xi)$ consisting of the tree of K -sequences of elements of $(\alpha^{+\xi})^K$ of length less than $(\alpha^{+\xi})^K$, i.e. we let $T_\alpha(\xi) = ((\alpha^{+\xi})^{<(\alpha^{+\xi})})^K$. This forcing remains \mathcal{S} -proper over $K[G_\alpha]$, as was shown in Lemma 33. However, again by Lemma 33 there exists a forcing $\mathbb{R}_\alpha(\xi)$ of size $\alpha^{+(\xi+2)}$ such that after forcing with $\mathbb{R}_\alpha(\xi)$, if H denotes the generic for $\mathbb{R}_\alpha(\xi)$, each outer model of $K[G_\alpha][H]$ which preserves ω_1 can not contain any branch through $T_\alpha(\xi)$ which is cofinal in $\alpha^{+\xi}$. Now we take the stationary set $S \subset \omega_1$ and code the pattern of S into an ω_1 -block of trees $T_\alpha(\xi) := ((\alpha^{+\xi})^{<(\alpha^{+\xi})})^K$, using only every third cardinal successor of α in K for a nontrivial forcing in order to guarantee \mathcal{S} -properness.

1. if $\xi \in S$ then we let $\mathbb{P}_{\xi \cdot 3}$ be the forcing which shoots a cofinal branch through the tree $T_\alpha(\xi^{+++})$ using the tree forcing with conditions that are nodes in the tree $T_\alpha(\xi^{+++})$.
2. if $\xi \notin S$ then pick $\mathbb{P}_{\xi \cdot 3}$ to specialize the tree $T_\alpha(\xi^{+++})$, using the already defined \mathcal{S} -proper specialization forcing.
3. if the index ξ of \mathbb{P}_ξ is not of the form $\eta \cdot 3$ for an $\eta < \omega_1$ we let \mathbb{P}_ξ be the trivial forcing.

We define \mathbb{Q}_α^0 to be the countable support iteration of the just defined \mathbb{P}_ξ , $\xi < \omega_1$.

Note that as K satisfies the GCH and as we only used every third K cardinal we ensure that the condition on the size of the forcings in Lemma 33 is met, thus each iterand \mathbb{P}_ξ is an \mathcal{S} -proper forcing and so is \mathbb{Q}_α^0 . Note further that if G^0 is \mathbb{Q}_α^0 -generic then in $M_1[G_\alpha][G^0]$ the model sees the stationary set S via the pattern of the trees $(T_\alpha(\alpha))^K$, having a cofinal branch or not.

To avoid unwanted patterns we can without any problems demand that all ω_1 -blocks are used, meaning that each block has a pattern (possibly a dummy pattern) written on it.

2.6.2 \mathbb{Q}_α^1

To define the next forcing \mathbb{Q}_α^1 we first fix a sufficiently big initial segment \mathcal{J}_η^K of K such that the generic filter $G_\alpha * G^0$ is also generic over \mathcal{J}_η^K . We then collapse the size of the structure $\mathcal{J}_\eta^K[G_\alpha][G^0]$ to ω_1 via the usual Levy collapse, $\mathbb{Q}_\alpha^1 := \text{Coll}(\lambda, \omega_1)$, where $\lambda = |\mathcal{J}_\eta^K|$. If G^1 is \mathbb{Q}_α^1 -generic then in $K[G_\alpha][G^0][G^1]$ there exists a subset $X_\alpha \subset \omega_1$ such that whenever M is a transitive model of ZFC, which is K -correct and $X_\alpha \in M$ then M will say that:

there is a K -cardinal α such that the previously coded stationary set S can be read off from a pattern of cofinal or bounded branches through canonical K -trees starting at α .

2.6.3 Localization

Our goal is to add a set $Y_\alpha \subset \omega_1$ which reflects the desired property down to all suitable models of size \aleph_0 . To be more specific we want a $Y_\alpha \subset \omega_1$ for which the following holds:

$\heartsuit \forall \beta < \omega_1 \forall M$ countable, transitive model of \mathbf{ZF}^- , $(Y_\alpha \cap \omega_1) \in M$, $(K)^M \subset \mathcal{J}_\eta^K$ for some $\eta < \omega_1$, $(\omega_1)^M = (\omega_1)^{(K)^M}$ then M satisfies that it can decode out of $Y_\alpha \cap \omega_1$ an ordinal $\bar{\alpha}$ such that for every $\xi < \omega_1$ ($\xi \in S$ if and only if $T_{\bar{\alpha}}(\xi^{+++})$ has a cofinal branch in M).

To force property \heartsuit we just use approximations of size \aleph_0 . Thus our forcing \mathbb{Q}_α^2 consists of the set of ω_1 -Cohen conditions $p : |p| \rightarrow 2$ in $K[G_\alpha][G^0][G^1]$ with the properties that

- The domain of p is a limit ordinal $< \omega_1$.
- The even part of p codes $X_\alpha \cap \omega_1$.
- For any limit ordinal $\xi < \omega_1$, $\xi \leq |p|$ and any transitive \mathbf{ZF}^- -model M of size ω_1 , with $\omega_1^M = (\omega_1)^{K^M}$ and which contains $p \upharpoonright \xi$, moreover satisfies that $(K)^M \subset (K)_\eta$ for some $\eta < \omega_1$ and $(\omega_1)^M = (\omega_1)^{(K)^M}$ we have that there is $\bar{\alpha}$ such that $\forall \xi < (\omega_1)^M$ $\xi \in S$ if and only if $(T_{\bar{\alpha}}(\xi^{+++}))^{(K)^M}$ has a branch in M which is cofinal in $(\bar{\alpha})^{+(\xi)}$.

Claim 39. *The forcing \mathbb{Q}_α^2 is \mathcal{S} -proper.*

Proof. Note that we can assume, via arguing by induction on the length of the iteration that \mathcal{S} is everywhere stationary in $K^{\mathbb{P}_\alpha * \mathbb{Q}_\alpha^0 * \mathbb{Q}_\alpha^1}$ as $\mathcal{S}(A^c)$ is. Note also that the forcing \mathbb{Q}_α^2 has the extendibility property, meaning that given some condition $p \in \mathbb{Q}_\alpha^2$ and some countable limit ordinal $\gamma > |p|$, we can always find a condition q extending p , such that $|q| = \gamma$. This is easily seen as given a condition $p \in \mathbb{Q}_\alpha^2$, one can extend its length to γ in such a way that the even entries still code X_α up to γ , and on the odd entries one codes an ordinal $\eta > \gamma$ into an ω -block of the odd entries up to γ . Let q be such a sequence. Then no model N which has q as an element can have γ as its ω_1 as it can use q to see the countability of $\eta > \gamma$. Thus the third property is automatically satisfied and the sequence q of length γ is a condition in \mathbb{Q}_α^2 . We want to show the \mathcal{S} -properness now, thus we shall exhibit a club $C \subset [H(\theta)^{K[G_\alpha][G^0][G^1]}]^\omega$ such that whenever we pick an arbitrary countable elementary submodel $M \in C \cap \mathcal{S}$ and some condition $p \in M \cap \mathbb{Q}_\alpha^2$ we can show that there is a stronger q which is (M, \mathbb{Q}_α^2) -generic. Choose C to be a club of elementary submodels of some $(K)_\sigma[G_\alpha][G^0][G^1]$, such that the set X_α is in every element of C , and let the ordinal σ be large enough to enable $(K)_\sigma[G_\alpha][G^0][G^1]$ to define its K properly, and therefore is able to realize that

1. $(K)_{\alpha^1}$ is an initial segment of the K -version which is computed in the model $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$. Moreover as $(K)_{\alpha^1}$ is coded into the set X_{α} , the model $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ will see that the set coded into X_{α} is an initial segment of its own version of K . We write $dec(X_{\alpha})$ to denote the mouse which is coded into X_{α} . Thus $dec(X_{\alpha}) \triangleleft (K)^{(K)_{\sigma}[G_{\alpha}][G^0][G^1]}$ does hold.
2. As $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ and $(K)_{\alpha^1}$ do have the same ω_1 , the sequence of K -trees, as computed in $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ and $(K)_{\alpha^1}$ respectively, which starts at the ordinal α (which is coded into X_{α}), will coincide. To be more precise, the sequence of K -cardinals starting at α , as computed in $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ and $(K)_{\alpha^1}$ respectively coincide. The trees $T_{\alpha}(\xi^{+++})$ as computed in $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ and $(K)_{\alpha^1}$ respectively coincide, and they do or do not have cofinal branches simultaneously.

Note that such a σ always exists due to the Reflection Principle, and the definability of K in small forcing extensions.

Now let M be from $C \cap \mathcal{S}$. We list all the dense subsets D_n which are elements of M and construct a descending sequence of \mathbb{Q}_{α}^2 -conditions $p = q_0 > q_1 > \dots$. We can demand that for every $k \in \omega$, q_k lies in the corresponding dense set D_k . Further we demand that $\sup_{k \in \omega} |dom(q_k)| = \xi = M \cap \omega_1$. If we can ensure that the limit of the q_k 's, denoted with q_{ω} is a condition we would be finished, as q_{ω} is $(M, \mathbb{Q}_{\alpha}^2)$ -generic. As the first two properties of conditions in \mathbb{Q}_{α}^2 hold automatically for q_{ω} , we have to ensure the third property. Note first that if $\xi < |q_{\omega}|$ holds then property 3 will hold automatically as well. Thus we shall show that the lower bound q_{ω} satisfies

- (*) For any countable, transitive ZF^- model N such that $q_{\omega} \in N$ and μ an ordinal such that $(K)^N \subset (K)_{\mu}$, if $\xi = \omega_1^N = (\omega_1^K)^N$ and $(\omega_1)^N = (\omega_1)^{(K)_{\mu}}$ then q_{ω} codes a $(K)^N$ -cardinal $\bar{\alpha}$ and certain generic filters such that if one builds the sequence of $(K)^N$ -cardinals $(\bar{\alpha}^{+\tau})^{(K)^N}$, and the according trees $(T_{\bar{\alpha}}(\xi^{+++}))^{(K)^N}$ then it holds that $\xi \in \dot{S}[G_{\alpha}][G^0][G^1]$ iff there is a cofinal branch through $(T_{\bar{\alpha}}(\xi^{+++}))^{(K)^N}$ in N .

Now if we take our countable $M \prec (K)_{\sigma}[G_{\alpha}][G^0][G^1]$, $M \in \mathcal{S}$ such that $X_{\alpha} \in M$, by elementarity we have that the two properties of $(K)_{\sigma}[G_{\alpha}][G^0][G^1]$ which we listed above still hold for M . Thus

1. M thinks that the real initial segment $(K)_{\alpha^1}$ is coded into X_{α} . Further M is also able to realize that this real initial segment is an initial segment of its own K .
2. $(\omega_1)^M = ((\omega_1)^K)^M$ and moreover M computes the same pattern of trees having cofinal branches or not as its own version of K . To be more precise the sequence of K -cardinals starting at α does not depend on whether we compute it in M or in $(K)^M$. And likewise for the pattern of trees having cofinal branches or not.

Thus its transitive collapse \bar{M} will decode the same information out of $\pi(X_\alpha) = X_\alpha \cap (\omega_1)^{\bar{M}}$, where π denotes the collapsing function. I.e. \bar{M} thinks that $X_\alpha \cap (\omega_1)^{\bar{M}}$ codes an initial segment of its own version of K together with an ordinal $\bar{\alpha}$ and some generically added branches through trees, which have all the information to see the intended pattern of cofinal or bounded trees $T_{\bar{\alpha}}(\xi^{+++})$ as computed in $(K)^{\bar{M}}$. But as M was assumed to be in \mathcal{S} it will collapse nicely, thus $\bar{M} = (K)_\xi$ and $X_\alpha \cap (\omega_1)^{\bar{M}}$ codes an initial segment $(K)_\eta$ of $(K)_\xi$.

Whenever N is a countable, transitive ZF^- -model, which satisfies that N is elementary with respect to the K defining first order formula $\Phi(x)$, $K^N \subset K_\mu$ for a $\mu < \omega_1$, $q_\omega \in N$, and $\xi = (\omega_1)^N = ((\omega_1)^{(K)})^N$, then it contains, as the even entries of q_ω code X_α , the set $X_\alpha \cap \xi$. So if N decodes the information packed into $X_\alpha \cap \xi$, it will obtain the ZF^- -model from above, namely $(K)_\eta \subset (K)^{\bar{M}}$, the $(K)_\eta$ -cardinal $\bar{\alpha}$, and certain generically added cofinal branches through trees, such that $(K)_\eta$ together with the generically added branches read off the intended pattern of bounded and cofinal trees. As N was assumed to be smart, i.e. able to see each K initial segment in it as such, N will be able to realize that $(K)_\eta$ is in fact also an initial segment of its own K , thus $(K)_\eta \subset (K)^N$.

The goal is to show that the model $(K)^N$ which the countable model N creates will define exactly the same sequence of cardinals starting at $\bar{\alpha}$ and exactly the same pattern of cofinal or bounded trees, as the model $(K)_\eta$. For the pattern which is seen by $(K)_\eta$ is the right one and we would be finished. In order to show this, we crucially use the definition of \mathcal{S} . First note that from our assumptions on N we get that there is a $\mu < \omega_1$ such that $(K)^N \subset (K)_\mu$ and $(\omega_1)^N = (\omega_1)^{(K)\mu}$. Thus, lining up already shown things, we have that

$$(K)_\eta \subset (K)^N \subset (K)_\mu$$

and their versions of ω_1 all coincide. Note now that as $M \in \mathcal{S}$ thus the transitive collapse \bar{M} is of the form $(K)_\zeta$ for some $\zeta < \omega_1$. By the definition of \mathcal{S} we have that $(K)^{(K)\zeta} = (K)_\zeta$, and as seen above we also have

$$(K)_\eta \subset (K)_\zeta.$$

Comparing ζ and μ we split into cases.

First assume that $\zeta \leq \mu$. Then remember that $(K)_\zeta$ has the property that whenever a larger (K) -initial segment has the same ω_1 then already all $(K)_\zeta$ -cardinals remain cardinals in the larger K -initial segment. As $(K)_\mu$ is of such a form and shares the same ω_1 , we infer that all cardinals of $(K)_\zeta$ are still cardinals in $(K)_\mu$. But by elementarity we have that all cardinals of $(K)_\eta$ are still cardinals in $(K)_\zeta$, thus the sequence of cardinals after $\bar{\alpha}$ is the same as evaluated in $(K)_\eta$ and $(K)_\mu$. And as $(K)_\eta \subset (K)^N \subset (K)_\mu$, the sequence of cardinals after $\bar{\alpha}$ in $(K)^N$ is the same as the sequence of cardinals after $\bar{\alpha}$ in $(K)_\eta$. Thus the sequence of trees in both models live

on the same ordinals. As $(\omega_1)^{(K)_\eta} = (\omega_1)^{(K)^N}$ the pattern of cofinal or bounded trees must coincide as well as otherwise the $(\omega_1)^{(K)^N}$ would have been collapsed.

Secondly assume that $\mu > \zeta$. Then $(K)_\eta \subset (K)^N \subset (K)_\mu \subset (K)_\zeta$ and as already noted above, the sequence of cardinals after $\bar{\alpha}$ does not depend on the model $(K)_\eta$ or $(K)_\zeta$. But this also implies that the sequence of cardinals does not depend on whether we compute them in $(K)^N$ or $(K)_\eta$. Thus the trees live on the same ordinals, independently from where we construct them, and again, even the pattern of bounded or cofinal trees agrees as $(\omega_1)^N = (\omega_1)^{(K)_\mu}$. This finishes the proof. \square

This finishes the definition of the three step iteration we use whenever the α -th entry of the \diamond -sequence a_α is the name of a stationary subset of ω_1 . What is left is to show that in the resulting model, the nonstationary ideal NS_{ω_1} is indeed Δ_1 -definable over $H(\omega_3)$ and $\text{NS}_{\omega_1} \upharpoonright A$ is \aleph_2 -saturated.

2.7 The definability of NS_{ω_1}

Goal of this section is the proof that in our final model $M_1[G_\delta]$ NS_{ω_1} is $\underline{\Delta}_1$ -definable over $H(\omega_3)$. The parameter will be ω_1 . Remember that the only patterns of length ω_1 which occur on trees $T(\xi)^K$ in $M_1[G_\delta]$ are the ones which code stationary sets.

Lemma 40. *If G denotes the generic for the forcing notion defined at the beginning of the last section. Then in $M_1[G]$, the nonstationary ideal is Δ_1 -definable over $H(\omega_3)$ using the parameter $H(\omega_1)$.*

Proof. Let $S \subset \omega_1$ be an arbitrary stationary subset of $M_1[G]$. By the inaccessibility of the Woodin cardinal δ , which represents the length of the iteration, we know that the \mathbb{P}_δ -name \dot{S} of S is in fact a \mathbb{P}_β -name for some $\beta < \delta$. Thus there is a stage $\alpha < \delta$ such that at stage α the name \dot{S} is considered by the \diamond -sequence. The rules of our iteration then force the characteristic function of S into the sequence of K -trees starting at the K -cardinal α , followed by a localization of this information.

We therefore know that there is a set $Y_\alpha \subset \omega_1$ such that

- \heartsuit For every countable, transitive model $M(\in L_{\omega_1}[Y_\alpha][K_{\omega_1}])$ of ZF^- which has $Y_\alpha \cap \omega_1^M$ as an element and which satisfies that $K^M \subset \mathcal{J}_\eta^K$ and $\omega_1^M = \omega_1^{K^M}$, there exists an ordinal $\beta \in M$ such that for every $\xi < \omega_1^M$: $\xi \in S$ if and only if $M \models T_\beta(\xi^{+++})$ has a $\beta^{+\xi}$ -cofinal branch.

Note that property \heartsuit can be written as Σ_1 over $H(\omega_3)$ with parameter K_{ω_1} , i.e. ω_1 . As stationarity is automatically Π_1 over $H(\omega_2)$ we see that in

$M_1[G_\delta]$ stationary sets have a $\underline{\Delta}_1$ definition over $H(\omega_3)$. What is left is to show that the definition \heartsuit characterizes stationarity in $M_1[G_\delta]$.

For that assume that S is an arbitrary subset of ω_1 and assume that there is a set $Y_\alpha \subset \omega_1$ such that \heartsuit holds. We want to conclude that S is indeed stationary. Note first that \heartsuit also holds for transitive models M which are uncountable. Indeed if M would be an uncountable, transitive model such that the antecedens of \heartsuit does hold but not its conclusion, then we can consider a countable elementary $N \prec M$ which is an element of \mathcal{S} and collapse it to obtain a countable transitive \bar{N} . By the definition of \mathcal{S} , \bar{N} collapses to a M_1 -initial segment $\bar{N} = \mathcal{J}_\eta^{M_1}$ for some countable η . Thus \bar{N} is a countable, transitive model M for which \heartsuit fails by elementarity, which is a contradiction. to our assumption that \heartsuit is true.

Hence as \heartsuit holds for models of arbitrary size we can consider large initial segments $\mathcal{J}_\eta^{M_1}$ of M_1 , for which the antecedens of \heartsuit holds trivially. Thus $\mathcal{J}_\eta^{M_1}$ witnesses that the pattern of the characteristic function of S is written into trees which are build using K of $\mathcal{J}_\eta^{M_1}$, I.e. the real K . But remember that we have set up the iteration \mathbb{P}_δ in such a way that the only patterns of cofinal or bounded trees $T_\beta(\xi)^K$ which can arise are the ones which code stationary subsets of ω_1 . Thus the pattern which codes S must stem from a stationary set which is what we wanted. □

2.8 $NS_{\omega_1} \upharpoonright A$ is \aleph_2 -saturated

What is left is to show that we indeed have the restricted nonstationary ideal saturated after forcing with the iteration.

Lemma 41. *In the final model $M_1[G_\delta]$ of the iteration the restricted nonstationary ideal $NS_{\omega_1} \upharpoonright A$ on ω_1 is \aleph_2 -saturated.*

Proof. Note first that for every $\alpha < \delta$ the forcing \mathbb{P}_α is \mathcal{S} -semiproper. Indeed each factor of the iteration is \mathcal{S} -proper, or \mathcal{S} -semiproper, thus if we iterate with countable support we obtain an \mathcal{S} -semiproper notion of forcing as we have argued at the beginning of the proof in section 2.1. This suffices to see that the iteration is stationary sets preserving, which is of highest importance for the following argument.

Assume now for a contradiction that in $M_1[G_\delta]$ there is a $\delta = \omega_2^{M_1[G_\delta]}$ -long antichain of stationary subsets of A . Denote this antichain with S and let τ denote its \mathbb{P}_δ -name. We claim that it is possible to find an inaccessible κ below δ such that the following three properties hold:

1. κ is $\mathbb{P} \oplus \tau$ -strong up to δ in M_1
2. $\kappa = \omega_2^{M_1[G \upharpoonright \kappa]}$

3. $\vec{S} \upharpoonright \kappa = (S_i : I < \kappa) = (\tau \cap (M_1)_\kappa)^{G \upharpoonright \kappa}$ is the maximal antichain in $M_1[G \upharpoonright \kappa]$ which is picked by the \diamond -sequence at stage κ .

This is clear as we can assume that our \diamond -sequence lives on the stationary subset of inaccessible cardinals below δ , and for all inaccessible κ property 2 automatically holds. Moreover the sets

$$C_1 := \{\kappa < \delta : \vec{S} \upharpoonright \kappa = (S_i : I < \kappa) = (\tau \cap (M_1)_\kappa)^{M_1[G \upharpoonright \kappa]}\}$$

and

$$C_2 := \{\kappa < \delta : \forall \alpha \forall S \in P(\omega_1) \cap M_1^{\mathbb{P}^\alpha} \text{ stationary } \exists \bar{S} \in \vec{S} \upharpoonright \kappa (S \cap \bar{S} \notin NS)\}$$

are both clubs, therefore hitting the stationary set T consisting of the points $\kappa < \delta$ where $\tau \cap (M_1)_\kappa = a_\kappa$ (remember: a_κ is the κ -th element of the \diamond -sequence $(a_\alpha : \alpha < \delta)$) and κ is τ -strong up to δ . Thus if κ is in the nonempty intersection $C_1 \cap C_2 \cap T$ then 1 and 2 are satisfied, and the recursive definition of our forcing \mathbb{P} yields that at stage κ , as $a_\kappa = \tau \cap (M_1)_\kappa$, the sealing forcing $\mathbb{S}((\tau \cap (M_1)_\kappa)^{G \upharpoonright \kappa})$ is at least considered, and in order to show property 3, it suffices to show that $(\tau \cap (M_1)_\kappa)^{G \upharpoonright \kappa} = \vec{S} \upharpoonright \kappa$ is maximal in $M_1[G \upharpoonright \kappa]$. But this is clear as by the RCS iteration properties we take at inaccessible κ 's the direct limit of the \mathbb{P}_α 's, thus each stationary $S \subset \omega_1$ in $M_1^{\mathbb{P}^\kappa}$ is already included in a $M_1^{\mathbb{P}^\alpha}$ for $\alpha < \kappa$. So we have ensured the existence of a κ with all the 3, above stated properties.

Now the forcing $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ can not be \mathcal{S} -semiproper at stage κ , as otherwise we would have to force with it, therefore killing the antichain \vec{S} . So there exists a condition $(p, c) \in \mathbb{S}(\vec{S} \upharpoonright \kappa)$ such that the set

$$\begin{aligned} \bar{T} := \{X \prec (H_{\kappa^+})^{M_1[G \upharpoonright \kappa]} : |X| = \aleph_0 \wedge (p, c) \in X \wedge \nexists Y \supset X (Y \prec (H_{\kappa^+})^{M_1[G \upharpoonright \kappa]} \\ \wedge |Y| = \aleph_0 \wedge (X \cap \omega_1 = Y \cap \omega_1) \wedge \exists (q, d) \leq (p, c) ((q, d) \text{ is } Y\text{-semigeneric})\} \\ \cap \mathcal{S}. \end{aligned}$$

is a stationary subset of \mathcal{S} in $M_1[G \upharpoonright \kappa]$, and by construction of our iteration, the κ -th forcing in \mathbb{P} is $Col(\omega_1, 2^{\aleph_2})$, so in $M_1[G \upharpoonright \kappa + 1]$ there is a surjection $f : \omega_1 \rightarrow (H_{\kappa^+})^{M_1[G \upharpoonright \kappa]}$. As $Col(\omega_1, 2^{\aleph_2})$ is proper the set \bar{T} remains stationary in $M_1[G \upharpoonright \kappa + 1]$ which implies that

$$T := \{\alpha < \omega_1 : f''\alpha \in \bar{T} \wedge \alpha = f''\alpha \cap \omega_1\}$$

is stationary in $M_1[G \upharpoonright \kappa + 1]$. As the tail $\mathbb{P}_{[\kappa+2, \delta]}$ remains \mathcal{S} -semiproper, seen as an iteration with $M_1[G \upharpoonright \kappa + 1]$ as ground model, we can infer that T remains stationary in $M_1[G]$ and therefore either $\neg A \cap T$ is stationary or there exists an $i_0 < \delta$ such that

$$(**) \quad T \cap S_{i_0} \text{ is stationary in } M_1[G].$$

Assume first that $\neg A \cap T$ is stationary. We want to derive a contradiction. Fix an $\alpha \in \neg A \cap T$, and let $Z := f''\alpha \in \bar{T}$. We can list inside $Z \prec (H_{\kappa^+})^{M_1[G \upharpoonright \kappa]}$ all the $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ -names for countable ordinals and a descending sequence of $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ -conditions $(p, c) > (p_0, c_0) > (p_1, c_1), \dots$ deciding more and more of them. We can assume that $\sup(\text{dom}(p_i)) = \alpha$ and thus the lower bound of the sequence $(p_i, c_i)_{i \in \omega}$ denoted by (p', c') with $\text{dom}(p') = \alpha + 1$ and $p'(\alpha) \in \vec{S} \upharpoonright \kappa$ is in fact already a condition in $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ as $\alpha \notin A$. Thus we have found a $(Z, \mathbb{S}(\vec{S} \upharpoonright \kappa))$ -semigeneric condition below (p, c) contradicting $Z \in \bar{T}$.

Thus (**) must hold. Let us shortly reflect the situation we are in. The idea is to mimic the short argument from above in our new case, i.e. we want to find a model $X \in \bar{T}$ such that we *can* find a $(X, \mathbb{S}(\vec{S} \upharpoonright \kappa))$ -semigeneric condition $(q, d) < (p, c)$, thus arriving at a contradiction. For that it would be sufficient to find an X such that

1. $X \cap \omega_1 \in T \cap S_{i_0}$ for some $i_0 < \kappa$
2. $f''(X \cap \omega_1) \subset X$
3. $S_{i_0} \in X$

The first thing to note here is that already the first item can be impossible to fulfill, for we can not restrict the size of i_0 and it might as well happen that $i_0 > \kappa$, and thus there is no chance of finding an $\mathbb{S}(\vec{S} \upharpoonright \kappa)$ -condition which is $(X, \mathbb{S}(\vec{S} \upharpoonright \kappa))$ -semigeneric in the way described above. Here we use the large cardinal property of δ being Woodin for rescue. We can find an elementary embedding $j : M_1 \rightarrow N$ such that $j(\kappa) > i_0$, and which fixes the predicate \vec{S} up to a cardinal $\lambda > i_0$. The task has now changed to find a $(j(X), j(\mathbb{S}(\vec{S} \upharpoonright \kappa)))$ -semigeneric condition below (c, p) in order to get a contradiction.

First let $\lambda < \delta$, $\lambda > \max(i_0, \kappa + 1)$ be such that $(\tau \cap (M_1)_\lambda)^{G \upharpoonright \lambda} = \vec{S} \upharpoonright \lambda$, so we have $(\tau \cap (M_1)_\lambda)^{G \upharpoonright \lambda}(i_0) = S_{i_0}$. As κ was chosen to be $\mathbb{P} \oplus \tau$ -strong up to δ we let $j : M_1 \rightarrow N$ be an elementary embedding with critical point κ , such that N is transitive, $N^\kappa \subset N$, $(M_1)_{\lambda+\omega} \subset N$, $j(\mathbb{P}) \cap (M_1)_\lambda = \mathbb{P} \cap (M_1)_\lambda$, and $j(\tau) \cap (M_1)_\lambda = \tau \cap (M_1)_\lambda$.

H should denote the generic filter for the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{N[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ over $N[G \upharpoonright \lambda]$. Then we lift j to an elementary embedding

$$j^* : M_1[G \upharpoonright \kappa] \rightarrow N[G \upharpoonright \lambda, H].$$

Notice that $((M_1)_{\lambda+\omega})^{M_1[G \upharpoonright \lambda]} = ((M_1)_{\lambda+\omega})^{N[G \upharpoonright \lambda]}$.

Now we let $(X_k : k < \omega_1) \in M_1[G \upharpoonright \kappa + 1]$ be an increasing continuous chain of countable elementary substructures of $(H_{j(\kappa)^+})^{N[G \upharpoonright \kappa + 1]}$ with $\{\tau \cap (M_1)_\lambda, i_0\} \subset X_0$ satisfying for all $k < \omega_1$ the following three properties:

- (a) $k \in X_{k+1}$

(b) $f''(X_k \cap \omega_1) \subset X_k$

(c) $j^{**}(X_k \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]}) \subset X_k$

Let $\bar{G} := G \upharpoonright [\kappa + 2, \lambda]$, and consider the set

$$\{X_k[\bar{G}] \cap \omega_1 : k < \omega_1\}$$

which is a club in $M_1[G \upharpoonright \lambda]$. We can demand that for all $k < \omega_1$, $X[\bar{G}] \cap \omega_1 = X \cap \omega_1$. And by the stationarity of $T \cap S_{i_0}$ we obtain an index $k \in \omega_1$ such that

$$X_k[\bar{G}] \cap \omega_1 = X_k \cap \omega_1 \in T \cap S_{i_0}.$$

Write $X := X_k$, $\alpha := X \cap \omega_1$. As at stage κ we had to force with the ω -closed $Col(2^{\aleph_2}, \aleph_1)$ we know that $X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \in M_1[G \upharpoonright \kappa]$. Remember that $f \in M_1[G \upharpoonright \kappa + 1]$ was chosen as a surjection of ω_1 onto $(H_{\kappa+})^{M_1[G \upharpoonright \kappa]}$, so as $\alpha \in T$ by definition of T $f''\alpha \in \bar{T}$ and $\alpha = f''\alpha \cap \omega_1$, and hence by (b)

$$f''\alpha \subset X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \in M_1[G \upharpoonright \kappa].$$

As $\alpha = f''\alpha \cap \omega_1$, $f''\alpha \in \bar{T}$ and $f''\alpha \subset X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]}$ we get that $X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \in \bar{T} \subset \mathcal{S}$ and therefore

$$(***) \quad j^*(X \cap H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \in j^*(\bar{T}) \subset j^*(\mathcal{S}).$$

Note that our second generic H , denoting the generic filter for the segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{N[G \upharpoonright \lambda]}$ of $j(\mathbb{P})$ over $N[G \upharpoonright \lambda]$ has not been specified yet. The segment $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{N[G \upharpoonright \lambda]}$ of $j(\mathbb{P} \upharpoonright \kappa)[G \upharpoonright \lambda]$ over $N[G \upharpoonright \lambda]$ is $j^*(\mathcal{S})[G \upharpoonright \lambda]$ -semiproper. By (***), $j^*(X \cap H_{\kappa+}^{M_1[G \upharpoonright \kappa]})(\bar{G}) \in j^*(\bar{T})(\bar{G})$. So we can say that as \bar{T} is closed under supersets with the same intersection with ω_1 so is $j(\bar{T})$. Hence $X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \in \bar{T}$ implies that $j^*(X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]}) \in j^*(\bar{T})$ which in turn implies that $X \in j^*(\bar{T})$ by property (c), and finally we see that $X[\bar{G}] \in j^*(\bar{T})[G \upharpoonright [\kappa + 1, \lambda]] \subset j(\mathcal{S})[G \upharpoonright \lambda]$. By the $j(\mathcal{S})[G \upharpoonright \lambda]$ -semiproperness of $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{N[G \upharpoonright \lambda]}$ we can now conclude that any condition $r \in \mathbb{P}_{[\lambda+1, j(\kappa)]} \cap X[\bar{G}]$ has a stronger condition $q \in \mathbb{P}_{[\lambda+1, j(\kappa)]}$ which is $(X[\bar{G}], \mathbb{P}_{[\lambda+1, j(\kappa)]})$ -semigeneric. If we pick H such that $q \in H$ then by semigenericity of q we obtain $X[\bar{G}, H] \cap \omega_1 = X[\bar{G}] \cap \omega_1 = X \cap \omega_1 = \alpha \in S_{i_0} = (\tau \cap (M_1)_\lambda)^{G \upharpoonright \lambda}(i_0) \in X[\bar{G}, H]$. But also due to (c) we have that

$$j^*(X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]}) = j^{**}(X \cap (H_{\kappa+})^{M_1[G \upharpoonright \kappa]}) \subset X[\bar{G}, H].$$

This gives us the desired contradiction as we can find an $(X[\bar{G}, H], j(\mathbb{S}(\vec{S} \upharpoonright \kappa)))$ -semigeneric condition below $j(p, c) = (p, c)$. Indeed we can just list the countably many names for countable ordinals in $X[\bar{G}, H]$ along with conditions of $j(\mathbb{S}(\vec{S} \upharpoonright \kappa))$ deciding them below (p, c) and let $(p', c') \in j(\mathbb{S}(\vec{S} \upharpoonright \kappa))$ be just the condition with $\text{dom}(c') = \text{dom}(d') = \alpha + 1$, $c'(\alpha) = \alpha$ and $p'(I) = S_{i_0}$ for some $I < \alpha$. So $X[\bar{G}, H]$ together with $(p', c') < (p, c)$ witness that $j^*(X \cap H_{\kappa+})^{M_1[G \upharpoonright \kappa]} \notin j^*(\bar{T})$, contradicting (***) \square

Chapter 3

NS_{ω_1} saturated and a projective well-order.

Goal of this section is a the proof of the following result:

Theorem 42. *Assume that $M_1^\#$ exists, then there exists a generic extension $M_1[G]$ of M_1 such that in $M_1[G]$ NS_{ω_1} is \aleph_2 -saturated and there is a lightface Σ_4^1 -definable wellorder on the reals.*

Its proof is organized as follows. First we introduce a method which codes reals into triples of ordinals (β, γ, δ) . This method was invented by A. Caicedo and B. Velickovic [2] building on the work of J. Moore on his set mapping reflection principle (see [12]), which he used to show that BPFA implies that the continuum is \aleph_2 . Their coding is introduced thoroughly in the first section of this chapter. The downside of this coding is that, as opposed to the coding we used in the last chapter, we do not have any control of when exactly a certain code gets written down in the universe. However this difficulty does not prevent a nice definition of a wellorder on the reals which is localizable, meaning that already small fragments of the universe, provided they contain a certain amount of information, are able to witness the coding. The relevant definitions for that are done in the second section and the theorem is finally proved in the last section, where we combine the coding forcings with forcings which seal off the long antichains in $P(\omega_1) \setminus \text{NS}_{\omega_1}$.

3.1 The coding method

The definition of the coding is a rather convoluted one. We have to introduce a couple of definitions and start with

Definition 43. *A \vec{C} -sequence, or a ladder system, is a sequence $(C_\alpha : \alpha \in \omega_1, \alpha \text{ a limit ordinal})$, such that for every α , $C_\alpha \subset \alpha$ is cofinal and the ordertype of C_α is ω .*

For three subsets $x, y, z \subset \omega$ we can consider the oscillation function. First turn use the set x into an equivalence relation \sim_x , defined on the set $\omega - x$ as follows: for natural numbers in the complement of x satisfying $n \leq m$ let $n \sim_x m$ if and only if $[n, m] \cap x = \emptyset$. This enables us to define:

Definition 44. For a triple of subset of natural numbers (x, y, z) list the intervals $(I_n : n \in k \leq \omega)$ of equivalence classes of \sim_x which have nonempty intersection with both y and z . Then the oscillation map $o(x, y, z,) : k \rightarrow 2$ is defined to be the function satisfying

$$o(x, y, z)(n) = \begin{cases} 0 & \text{if } \min(I_n \cap y) \leq \min(I_n \cap z) \\ 1 & \text{else} \end{cases}$$

Next we want to define how suitable countable subsets of ordinals can be used to code reals. For that suppose that $\omega_1 < \beta < \gamma < \delta$ are fixed limit ordinals, and that $N \subset M$ are countable subsets of δ . Assume further that $\{\omega_1, \beta, \gamma\} \subset N$ and that for every $\eta \in \{\omega_1, \beta, \gamma\}$, $M \cap \eta$ is a limit ordinal and $N \cap \eta < M \cap \eta$. We can use (N, M) to code a finite binary string. Namely let \bar{M} denote the transitive collapse of M , let $\pi : M \rightarrow \bar{M}$ be the collapsing map and let $\alpha_M := \pi(\omega_1)$, $\beta_M := \pi(\beta)$, $\gamma_M := \pi(\gamma)$, $\delta_M := \bar{M}$. These are all countable limit ordinals. Further set $\alpha_N := \sup(\pi''(\omega_1 \cap N))$ and let the height $n(N, M)$ of α_N in α_M be the natural number defined by

$$n(N, M) := \text{card}(\alpha_N \cap C_{\alpha_M})$$

where C_{α_M} is an element of our previously fixed ladder system. As $n(N, M)$ will appear quite often in the following we write shortly n for $n(N, M)$. Note that as the ordertype of each C_α is ω , and as $N \cap \omega_1$ is bounded below $M \cap \omega_1$, $n(N, M)$ is indeed a natural number. Now we can assign to the pair (N, M) a triple (x, y, z) of finite subsets of natural numbers as follows:

$$x := \{\text{card}(\pi(\xi) \cap C_{\beta_M}) : \xi \in \beta \cap N\}.$$

Note that x again is finite as $\beta \cap N$ is bounded in the cofinal in β_M -set C_{β_M} , which has ordertype ω . Similarly we define

$$y := \{\text{card}(\pi(\xi) \cap C_{\gamma_M}) : \xi \in \gamma \cap N\}$$

and

$$z := \{\text{card}(\pi(\xi) \cap C_{\delta_M}) : \xi \in \delta \cap N\}.$$

Again it is easily seen that these sets are finite subsets of the natural numbers. We can look at the oscillation $o(x \setminus n, y \setminus n, z \setminus n)$ (remember we let $n := n(N, M)$) and if the oscillation function at these points has a domain bigger or equal to n then we write

$$s_{\beta, \gamma, \delta}(N, M) := \begin{cases} o(x \setminus n, y \setminus n, z \setminus n) \upharpoonright n & \text{if defined} \\ * & \text{else} \end{cases}$$

Similarly we let $s_{\beta,\gamma,\delta}(N, M) \upharpoonright l = *$ when $l > n$. The seemingly arbitrary move in considering the oscillation of the triple $(x \setminus n, y \setminus n, z \setminus n)$ instead of just the oscillation of (x, y, z) will become clear after the next definition. Finally we are able to define what it means for a triple of ordinals (β, γ, δ) to code a real r .

Definition 45. For a triple of limit ordinals (β, γ, δ) , we say that it codes a real $r \in 2^\omega$ if there is a continuous increasing sequence $(N_\xi : \xi < \omega_1)$ of countable sets of ordinals whose union is δ and which satisfies that whenever $\xi < \omega_1$ is a limit ordinal then there is a $\nu < \xi$ such that

$$r = \bigcup_{\nu < \eta < \xi} s_{\beta,\gamma,\delta}(N_\eta, N_\xi)$$

As a short remark, note that if we would define the $s_{\beta,\gamma,\delta}$ function with the full oscillation of (x, y, z) instead, then our above definition would become useless. Indeed if N_ξ is a continuous increasing sequence whose union is δ then for some fixed limit $\xi < \omega_1$, and if we construct the sets x, y, z for the pair (N_ν, N_ξ) , $\nu < \xi$, the set x would for increasing ν eventually just become an enumeration of $\text{card}(\eta \cap C_{\beta_M})$ for $\eta \in \beta \cap N_\nu$. Likewise the sets y and z would just become eventually an enumeration of the natural numbers, and thus completely useless for any coding purpose. The idea to overcome this, is to constantly throw away already coded information, namely the n , and code the real r over and over again into the oscillation pattern.

Our next task is to combine the forcings which make NS_{ω_1} \aleph_2 -saturated, and the forcings which will generically add witnesses for ordinal triples coding reals r in such a way that in the end a projective well-order of the reals is possible. For that we should take a closer look at the forcings which add coding witnesses. These forcings have in common that their generics exist, assuming the so called Mapping Reflection Principle (MRP), which was introduced by Justin Moore. The MRP is a forcing axiom whose consistency strength lies in between the proper forcing axiom PFA and the bounded proper forcing axiom BPFA. Moore famously used the MRP to show that already the bounded proper forcing axiom suffices to decide the value of the continuum, namely $\text{ZFC} + \text{BPFA} \vdash 2^{\aleph_0} = \aleph_2$. We follow Caicedo Velickovic in proving that MRP implies that for every real r there is a triple of ordinals (β, γ, δ) such that $\omega_1 < \beta < \gamma < \delta < \omega_2$ which code the real r . In fact the witnesses for the coding can be added by small proper forcings, which will allow us to combine these with the forcings which seal off long antichains in $P(\omega_1) \setminus \text{NS}_{\omega_1}$. For the definition of the MRP we need the following local version of stationarity:

Definition 46. Let θ be a regular cardinal, X be an uncountable set, let $M \prec H_\theta$ be a countable elementary submodel which contains $[X]^\omega$. Then $S \subset [X]^\omega$ is M -stationary if for every club subset C of $[X]^\omega$, $C \in M$ it holds

that

$$C \cap S \cap M \neq \emptyset.$$

Definition 47. Let X be an uncountable set, $N \in [X]^\omega$ and $x \subset N$ finite. Then the Ellentuck topology on the set $[X]^\omega$ is generated by base sets of the form

$$[x, N] := \{Y \in [X]^\omega : x \subset Y \subset N\}.$$

From now on whenever we say open we mean open with respect to the Ellentuck topology.

Definition 48. Let X be an uncountable set, let θ be a large enough regular cardinal so that $[X]^\omega \in H_\theta$. Then a function Σ is said to be open stationary if and only if its domain is a club $C \subset [H_\theta]^\omega$ and for every countable $M \in C$, $\Sigma(M) \subset [X]^\omega$ is open and M -stationary.

Equipped with these notions we can introduce the mapping reflection principle:

Definition 49. Let Σ be an open stationary function with domain some club $C \subset [H_\theta]^\omega$ and range $P([X]^\omega)$. Then there is a continuous sequence of models $(N_\xi : \xi < \omega_1)$ in $\text{dom}(\Sigma)$ such that for every limit ordinal ξ there is a $\nu < \xi$ such that for every η with $\nu < \eta < \xi$, $N_\eta \cap X \in \Sigma(N_\xi)$.

Key here is that these sequences of models which witness the truth of the MRP can always be forced with a proper forcing:

Proposition 50. PFA proves MRP.

Proof. The goal is to show that the natural forcing which adds a continuous sequence of models witnessing the MRP for a stationary set mapping Σ is always proper. So given such a function Σ with domain $C \subset [H_\theta]^\omega$ and range $P([X]^\omega)$ we let \mathbb{P}_Σ be the partial order whose elements are functions $p : \alpha + 1 \rightarrow \text{dom}(\Sigma)$, α countable, which are continuous and \in -increasing, and which additionally satisfy the MRP-condition on its limit points, namely that for every $0 < \nu < \alpha$ there is a $\nu_0 < \nu$ such that $p(\xi) \cap X \in \Sigma(p(\nu))$ for every ξ in the interval (ν_0, ν) . The order is by extension. The first thing to note is that sets of the form $D_\alpha := \{p \in \mathbb{P}_\Sigma : \alpha \in \text{dom}(p)\}$ are always dense. This is true as the trivially dense sets $D_x := \{p \in \mathbb{P}_\Sigma : \exists \beta \in \text{dom}(p)(x \in p(\beta))\}$ ensure that whenever we force with \mathbb{P}_Σ there will be a surjection from $\{\alpha : \exists p \in G(\alpha \in \text{dom}(p))\}$ onto the uncountable X . Thus once we show that the forcing \mathbb{P}_Σ is proper, and therefore ω_1 -preserving the ω_1 -many dense sets D_α and PFA will give the desired reflecting sequence. Note that we will not use the density of the D_α 's to show that \mathbb{P}_Σ is proper, so we avoid a circular reasoning.

To see that \mathbb{P}_Σ is proper we fix a large enough cardinal λ and a countable, elementary submodel $M \prec H_\lambda$ which contains Σ , \mathbb{P}_Σ , a condition $p \in \mathbb{P}_\Sigma$,

and the structure $H_{|\mathbb{P}_\Sigma|+}$. We list all the dense subsets (D_0, D_1, \dots) of \mathbb{P}_Σ which we can find in M and build by recursion a descending sequence of conditions $(p_i : I \in \omega)$ in M , starting at $p_0 := p$ hitting the corresponding D_{I-1} . Assume that we have already built conditions up to $I \in \omega$. We let N'_i be a countable elementary submodel of $H_{|\mathbb{P}_\Sigma|+}$ containing H_θ , D_i , \mathbb{P}_Σ and p_i , and build the club of countable structures $C_i := \{N'_i \cap X : N'_i \text{ as just described}\}$. Note that this club will be in M , and further that for every club on $[H_\theta]^\omega$ which is in M , the set $M \cap H_\theta$ will be in the club. Thus the set $M \cap H_\theta$ will be in the domain of Σ and by the definition of Σ , the set $\Sigma(M \cap H_\theta)$ is $M \cap H_\theta$ -stationary and open. So there is an $N_i \in C_i \cap \Sigma(M \cap H_\theta) \cap M$, and by the definition of the Ellentuck topology, there is a finite subset of N_i called x_i such that $[x_i, N_i] \subset \Sigma(M \cap H_\theta)$. We first extend the condition p_i to $q_i := p_i \cup \{\zeta_i + 1, \text{hull}^{H_\theta} p_i(\zeta_i) \cup x_i\}$, for ζ_i the maximum of the domain of p_i . This condition q_i will also be in N'_i as all its defining parameters are, thus as N'_i also contains D_i we can extend the condition q_i to a $p_{I+1} \in N'_i \cap D_i$. Note that as we are working in N'_i , no matter how we extend q_i , the range of the extended condition intersected with X will always be contained in $N_i = N'_i \cap X$, and as $\Sigma(M \cap H_\theta \supset [x_i, N_i])$, it will also be contained in $\Sigma(M \cap H_\theta)$. Then if we set $p_\omega := \bigcup_{I \in \omega} p_I \cup (\omega, (M \cap H_\theta))$ then this will be a condition in \mathbb{P}_Σ , which is by construction below p and (M, \mathbb{P}_Σ) -generic, thus the forcing is proper. \square

We can use the MRP to show that in its presence the function $s_{\beta\gamma\delta}$ eventually becomes stable in the following way:

Proposition 51. *Assume that the MRP does hold and let $\omega_1 < \beta < \gamma < \delta < \omega_2$ be limit ordinals with uncountable cofinality. Then there is a continuous sequence of countable sets of ordinals $(N_\xi : \xi < \omega_1)$ such that $\bigcup_{\xi < \omega_1} N_\xi = \delta$ and such that for every limit ordinal $\xi < \omega_1$ and every $n \in \omega$ there is a $s_\xi^n \in 2^n \cup \{*\}$ and there is a $\nu < \xi$ such that*

$$s_{\beta\gamma\delta}(N_\eta, N_\xi) \upharpoonright n = s_\xi^n$$

for every η in the interval (ν, ξ) . Moreover already BPFA suffices for the conclusion.

The proof is a consequence of the following

Lemma 52. *Again assume the MRP and suppose that for every $\alpha < \omega_1$ we have a partition of α into two sets K_α^i , $I \in 2$ such that both are clopen sets in the usual order topology on ω_1 . Then there is a club $C \subset \omega_1$ such that for every limit point ξ of C there is an $I \in 2$ such that $C \setminus K_\xi^i$ is bounded below ξ . Already BPFA does suffice for the conclusion.*

Proof. Consider ω_1 as a subset of $[\omega_1]^\omega$, then it is easy to see that the property "not an ordinal in $[\omega_1]^\omega$ " is an open set in the Ellentuck topology, hence ω_1 is closed in $[\omega_1]^\omega$. The restriction of the Ellentuck topology to ω_1 yields the usual order topology on ω_1 .

If $M \prec H_\theta$ is countable and $\alpha = M \cap \omega_1$ then one of the K_α^i 's is M -stationary. Indeed assume not then there are two clubs C_0 and C_1 in M which have empty intersection with K_α^0 and K_α^1 respectively. But then $C_0 \cap C_1 = \emptyset$ in M and by elementarity in H_θ which is a contradiction. Now set

$$\Sigma(M) := [\omega_1]^\omega \setminus K_\alpha^i$$

where K_α^{1-I} is the stationary part of the partition. As this is an open stationary map we can apply the MRP to obtain a sequence $(N_\xi : \xi < \omega_1)$ such that for all limits ξ there is a $\nu < \xi$ such that $N_\eta \cap \omega_1 \in \Sigma(N_\xi)$ for every $\eta \in (\nu, \xi)$. Then $C := \{N_\xi \cap \omega_1 : \xi < \omega_1\}$ is a club as required.

To see that already BPFA suffices for the conclusion note that the statement of the Lemma is of the form Σ_1 with parameters $K_\alpha^i \subset \omega_1$. We have already seen that each instance of the MRP can be forced by a proper forcing. As BPFA is equivalent to the assertion that $H_{\aleph_2} \prec V^{\mathbb{P}}$ for every proper \mathbb{P} we see that already BPFA is sufficient for the statement. \square

proof of the Proposition. Consider a triple $(\omega_1 < \beta < \gamma < \delta < \omega_2)$ of cofinality ω_1 and let $(N_\xi : \xi < \omega_1)$ be a continuous sequence of countable sets of ordinals whose union is δ . We can assume without loss of generality that $\{\omega_1, \beta, \gamma\} \subset N_0$ and that $N_\eta \cap \omega_1 < N_\xi \cap \omega_1$ holds for $\eta < \xi$ and similar inequalities hold for β and γ . For a fixed natural number $n \in \omega$ and an ordinal $\xi < \omega_1$ we can apply the previous Lemma to obtain a partition of ξ into $2^n + 1$ many sets $\{K_\xi^I : I \in 2^{n+1}\}$ via setting

$$\eta \in K_\xi^I \text{ if and only if } s_{\beta\gamma\delta}(N_\eta, N_\xi) \upharpoonright n = s_i$$

for s_i the i -th element of $2^n \cup \{*\}$ in some fixed well-order. This partition consists even of clopen sets as already finitely many elements $\{n_0, n_1, \dots, n_k\}$ of N_ξ suffice that $s_{\beta\gamma\delta}(\{n_0, \dots, n_k\}, N_\xi) = s_{\beta\gamma\delta}(N_\eta, N_\xi)$ and thus each K_ξ^I is clopen. We apply the previous Lemma to obtain for every $n \in \omega$ a club $C_n \subset \omega_1$ such that for every limit point $\xi \in C$ eventually all points of C below ξ are in some fixed K_ξ^I . To K_ξ^I corresponds an $s_\xi^n \in 2^n \cup \{*\}$ and we have that for every limit $\xi \in C_n$ there is an $\eta < \xi$ such that $s_{\beta\gamma\delta}(N_\eta, N_\xi) \upharpoonright n = s_\xi^n$ for every $\eta \in (\nu, \xi)$. If we set $C := \bigcap_{n \in \omega} C_n$ then the sequence $(N_\xi : \xi \in C)$ is as desired. \square

We will use the MRP to show that for every real r there is a proper notion of forcing which will introduce a sequence of countable models which witnesses that the real r is coded by a triple (β, γ, δ) of limit ordinals below \aleph_2 . The procedure is the same as before, namely we show first that MRP

implies that every real r is coded by a triple of ordinals, and then show that in fact BPPA suffices for the conclusion. We start with fixing an $F : [\omega_4]^{<\omega} \rightarrow \omega_4$. We will define a sequence of two player games \mathcal{G}_ν^F for $\nu < \omega_1$.

I	β_0	γ_0	δ_0	β_1	$\gamma_1 \dots$
II	κ_0, ϵ_0	λ_0, ϑ_0	μ_0, ϖ_0	$\kappa_1, \epsilon_1 \dots$	

The rules of the game \mathcal{G}_ν^F are as follows: player I starts by playing an ordinal $\beta_0 < \omega_2$, then player II responds with a pair of ordinals $\kappa_0 \leq \epsilon_0 < \omega_2$, $\beta_0 \leq \kappa_0$. In the next round player I plays an ordinal $\gamma_0 < \omega_3$ and II responds with playing ordinals $\lambda_0 \leq \vartheta_0 < \omega_3$ such that $\gamma_0 \leq \lambda_0$. This is followed by player I picking a $\delta_0 < \omega_4$ and II playing $\mu_0 \leq \varpi_0 < \omega_4$ with $\delta_0 \leq \mu_0$. Then I goes back to play below ω_2 : he picks a $\beta_1 < \omega_2$ greater than ϵ_0 and II responds with a pair κ_1, ϵ_1 such that $\beta_1 \leq \kappa_1$, and the game continues with I playing below ω_3 , II responding, I playing below ω_4 , II responding and so on. The first player who violates the rules loses. Otherwise the game will determine an infinite sequence of ordinals and we set

$$X := cl_F(\{\kappa_n, \lambda_n, \mu_n : n \in \omega\} \cup \nu)$$

where cl_F should denote the closure under the function F and $\nu < \omega_1$ is the index of the ν -th game \mathcal{G}_ν^F . We say that player II wins if the following conditions hold:

- $X \cap \omega_1 = \nu$
- $X \cap [\beta_n, \beta_{n+1}) \subset [\beta_n, \epsilon_n)$ for all $n \in \omega$.
- $X \cap [\gamma_n, \gamma_{n+1}) \subset [\gamma_n, \vartheta_n)$ for all $n \in \omega$.
- $X \cap [\delta_n, \delta_{n+1}) \subset [\delta_n, \varpi_n)$ for all $n \in \omega$.

If player I wins then he has won already after finitely many stages so by the Gale-Stewart theorem \mathcal{G}_ν^F is determined for every $\nu < \omega_1$. Player I wins \mathcal{G}_ν^F only for few $\nu < \omega_1$ as is shown now. Let

$$A_F := \{\nu < \omega_1 : \text{player I has a winning strategy in } \mathcal{G}_\nu^F\}$$

then

Lemma 53. *A_F is nonstationary.*

Proof. The proof is by contradiction, thus assume that A_F is stationary for some function $F : [\omega_4]^\omega \rightarrow \omega_4$. First notice that there is a winning strategy for I which works for all $\nu \in A_F$ simultaneously. Indeed we can define a strategy σ for I which maps a given state s of the game to the supremum of all the ordinals of the form $\sigma_\nu(s)$, $\nu \in A_F$, where σ_ν is a winning strategy for I. This strategy σ is again a winning strategy for I as a supposed play

of the game where I follows σ but nevertheless yields a victory of II will immediately give II a way of winning against any σ_ν , contradicting the fact that σ_ν is a winning strategy for I.

A similar consideration shows that we can further assume that whenever player I has to pick an ordinal $\gamma_{n+1} < \omega_3$, he can do so without looking at what player II decided to play for κ_n, ϵ_n . Indeed note that after all there are only \aleph_2 -many choices for II to make and I can simply play the supremum of all the answers, which again is a winning strategy for I. Likewise for playing $\delta_{n+1} < \omega_4$, I does not have to care about the choices of II for λ_n and ϑ_n . We will assume that the winning strategy σ for player II has the above described properties

Now we describe a way for II to win against the strategy σ , thus giving us the desired contradiction. We first fix a large enough regular θ and an increasing, continuous elementary chain of \aleph_1 -sized models of length $\omega_1 \cdot \omega$

$$P_0 \prec P_1 \prec \dots P_\xi \prec \dots \prec H_\theta, \quad \xi < \omega_1 \cdot \omega$$

such that $F, A_F, \sigma \in P_0$. Let $N_n := P_{\omega_1 \cdot n}$ and let N be the union of the N_n 's. Let

$$\zeta_n := \sup(N_n \cap \omega_2)$$

$$\eta_n := \sup(N_n \cap \omega_3)$$

$$\theta_n := \sup(N_n \cap \omega_4)$$

then all of these ordinals have cofinality ω_1 and the club subsets of N_n which climb up to them are elements of N_{n+1} as the sequence is an elementary chain. We fix a countable $M \prec N$ containing all the just mentioned clubs and such that $\{\zeta_n : n \in \omega\} \subset M$, $\{\eta_n : n \in \omega\} \subset M$, $\{\theta_n : n \in \omega\} \subset M$ and $F, A_F, \sigma \in M$ holds. Further by the assumed stationarity of A_F we can demand that $\alpha := M \cap \omega_1 \in A_F$. We will describe now a game where I follows σ which will nevertheless result in a victory for II. Assume that we are in the n -th round and the position p_{n-1} of the game looks like this $p_{n-1} = (\beta_0, (\kappa_0, \epsilon_0), \gamma_0, \dots, \delta_{n-1}, (\mu_{n-1}, \varpi_{n-1}))$. Assume further that $p_{n-1} \in N_n$. Then I plays according to σ , thus $\beta_n = \sigma(p_{n-1})$ and as both σ and p_{n-1} are in N_n , β_n will also be in N_n . Then II will follow by choosing $\kappa_n \in M \cap N_{n+1}$ such that $\zeta_n \leq \kappa_n < \omega_2$. Let $\epsilon_n := \sup(M \cap N_{n+1} \cap \omega_2)$. Recall that $\zeta_{n+1} = \sup(N_{n+1} \cap \omega_2)$ has cofinality ω_1 and as M is countable, $\epsilon_n < \zeta_{n+1}$. As N_{n+1} contains a club set which converges to ζ_{n+1} and this club is in M we conclude that $\epsilon_n \in N_{n+1}$. Player II plays (κ_n, ϵ_n) . Player I then answers with $\sigma(p_{n-1}(\beta_n, (\kappa_n, \epsilon_n))) = \gamma_n$. As σ was assumed to be not dependent on the previous choice of player I we know that $\gamma_n \in N_n$. II responds with picking a $\lambda_n \in M \cap N_{n+1}$ with $\eta_n \leq \lambda_n < \omega_3$ and $\vartheta_n := \sup(M \cap N_{n+1} \cap \omega_3)$. The same reasoning as above yields that $\vartheta_n < \eta_{n+1}$ and $\vartheta_n \in N_n$. Player I the answers with δ_n with the help of the strategy σ .

Again, as $\sigma \in N_n$, and as σ does not depend on player I's choice of λ_n and ϑ_n , $\delta_n \in N_n$. Player II then follows with playing $\mu_n \in M \cap N_{n+1}$ such that $\theta_n \leq \mu_n < \omega_4$. If we let $\varpi_n := \sup(M \cap N_{n+1} \cap \omega_4)$ then again $\varpi_n < \theta_{n+1}$ and $\varpi_n \in N_{n+1}$. II then picks (μ_n, ϖ_n) . This defines the next stage p_n of the game and by summing up already shown things we see that $p_n \in N_{n+1}$. If we let

$$X := cl_F(\{\kappa_i, \lambda_i, \mu_i : I \in \omega\} \cup \alpha)$$

then $X \subset M$ as all the relevant information is present in M . This implies that $X \cap \omega_1 = \alpha$. By construction we also have that $\beta_{n+1} \in N_{n+1}$ and $\epsilon_n = \sup(M \cap N_{n+1} \cap \omega_2)$, thus $X \cap [\beta_n, \beta_{n+1}) \subset X \cap [\beta_n, \epsilon_n)$ for every n . Likewise we have that for every n , $X \cap [\gamma_n, \gamma_{n+1}) = X \cap [\gamma_n, \vartheta_n)$ and $X \cap [\delta_n, \delta_{n+1}) = X \cap [\delta_n, \varpi_n)$. Thus II has won the game \mathcal{G}_α^F , however $\alpha \in A_F$ which is a contradiction. \square

If we assume that MRP holds then arbitrary reals r can be coded into triples of ordinals, as is shown now:

Lemma 54. *If MRP holds then every real r is coded into a triple (β, γ, δ) such that the each element of the triple has cofinality ω_1 and $\omega_1 < \beta < \gamma < \delta < \omega_2$.*

Proof. Let r be an arbitrary real. We let θ be a regular cardinal which is large enough, and let the open stationary set mapping Σ^r to be defined on the club of countable elementary submodels $M \prec H_\theta$ with values set as follows:

$$\Sigma^r(M) := \{N \in [M \cap \omega_4]^\omega : s_{\omega_2, \omega_3, \omega_4}(N, M \cap \omega_4) \text{ is an initial segment of } r\}$$

We have to check that Σ^r is indeed open and stationary. Openness is clear as already finite information $T \subset N$ suffices to determine the value of $s_{\omega_2, \omega_3, \omega_4}(N, M \cap \omega_4)$. Thus if $N \in \Sigma^r(M)$ then already an open set of the form $[m, N]$ will be a subset of $\Sigma^r(M)$. To show that Σ^r is stationary needs more work and the proof of this will be postponed. Instead we finish the proof of the lemma under the assumption that Σ^r is open stationary.

Using MRP we obtain a reflecting sequence $(M_\xi : \xi < \omega_1)$ for Σ^r . Let $M := \bigcup_{\xi < \omega_1} M_\xi$ and let \bar{M} be its transitive collapse with π the collapsing map. Let $\beta = \pi(\omega_2)$, $\gamma = \pi(\omega_3)$, $\delta = \pi(\omega_4)$. All these ordinals are in the interval (ω_1, ω_2) and their cofinalities are ω_1 . Let $N_\xi := \pi(M_\xi \cap \omega_4)$ and we get that

$$s_{\beta, \gamma, \delta}(N_\eta, N_\xi) = s_{\omega_2, \omega_3, \omega_4}(M_\eta \cap \omega_4, M_\xi \cap \omega_4)$$

for every $\eta < \xi < \omega_1$.

Moreover for every ξ limit we have continuity in that $N_\xi = \bigcup_{\eta < \xi} N_\eta$. Which implies that the $n(N_\eta, N_\xi)$ converges to ω for $\eta \rightarrow \xi$, where as a short

remainder $n(N_\eta, N_\xi)$ was defined to be the size of the set $C_{\pi(\omega_1)} \cap \pi''(\omega_1 \cap N_\eta)$ for π the collapsing function of N_ξ . As $s_{\beta, \gamma, \delta}(N_\eta, N_\xi) = s_{\omega_2, \omega_3, \omega_4}(M_\eta \cap \omega_4, M_\xi \cap \omega_4)$, and as $(M_\xi : \xi < \omega_1)$ is a reflecting sequence for Σ^r we know that there is a $\nu < \xi$ such that

$$r = \bigcup_{\nu < \eta < \xi} s_{\beta, \gamma, \delta}(N_\eta, N_\xi).$$

This ends our proof as the triple (β, γ, δ) code r as desired. \square

Lemma 55. *Already BPFA suffices for the conclusion of the last Lemma*

Proof. We will use Bagaria's characterization of BPFA again, namely that BPFA is equivalent to the assertion that $H_{\aleph_2} \prec_1 V^{\mathbb{P}}$ for every proper \mathbb{P} . Note that in the proof of the previous Lemma a proper forcing will add a reflecting sequence for the open stationary function Σ^r and the existence of a triple of ordinals coding the reals r is a Σ_1 statement with parameters the ladder system C and the real r . Thus already BPFA suffices to guarantee such a triple. \square

What is still left to show is that the function Σ^r is M -stationary for every M in its domain.

Lemma 56. *Let Σ^r be the function from the proof of Lemma 77,*

$$\Sigma^r(M) := \{N \in [M \cap \omega_4]^\omega : s_{\omega_2, \omega_3, \omega_4}(N, M \cap \omega_4) \text{ is an initial segment of } r\}.$$

Then Σ^r is M -stationary for every $M \prec H_\theta$, where θ is a sufficiently large regular cardinal.

Proof. The set X_{Σ^r} which comes along with Σ^r is ω_4 . As the closed filter on $[\omega_4]^\omega$ is generated by sets which are closed under functions $F : [\omega_4]^{<\omega} \rightarrow \omega_4$ it suffices to show that whenever $M \prec H_\theta$ and $F : [\omega_4]^{<\omega} \rightarrow \omega_4$ is a function in M , then there is an $X \in M$ which is closed under F and for which $X \in \Sigma^r(M)$ does hold. We will do this using the previously defined games \mathcal{G}_ν^F and the already proved fact that there is a club $C \subset \omega_1$ of ordinals ν for which player II has a winning strategy.

Thus let M be an arbitrary elementary submodel of H_θ . We will argue almost entirely inside M . Let \bar{M} as always denote the transitive collapse of M , let π be the collapsing function and set $\alpha_M := \pi(\omega_1)$, $\beta_M := \pi(\omega_2)$, $\gamma_M := \pi(\omega_3)$ and $\delta_M := \pi(\omega_4)$. Let C be our fixed ladder system, and for an ordinal $\rho < \omega_4$ let $ht_{\omega_i}(\rho)$, the height of ρ in ω_i be defined as $ht_{\omega_i}(\rho) := \text{card}(\pi(\rho) \cap C_{\pi(\omega_i)})$. Our goal is to show that we can find a $\nu \in C \cap M$ and play finitely many rounds of the game \mathcal{G}_ν^F inside M such

that if T is the finite set of relevant ordinals played by II and $X := cl_F(T \cup \nu)$ then $s_{\omega_2 \omega_3 \omega_4}(X, M \cap \omega_4)$ is an initial segment of r and thus in $\Sigma^r(M)$. The two players I and II will collaborate to achieve this.

We consider the set of winning strategies σ_ν for II and what they do at stage one. Let us assume that I plays 0 in all of his three first moves. II will respond with pairs (κ'_0, ϵ'_0) , $(\lambda'_0, \vartheta'_0)$, and (μ'_0, ϖ'_0) . We can consider the suprema of the sets $\{\epsilon'_0 : \nu \in C\}$, $\{\vartheta'_0 : \nu \in C\}$ and $\{\varpi'_0 : \nu \in C\}$ and call them ϵ , ϑ and ϖ respectively. These suprema are elements of M but we leave M for a moment to compare the height of ϵ below ω_2 , the height of ϑ below ω_3 and the height of ϖ below ω_4 . We pick the maximum of these three natural numbers, call it k and fix a $\nu \in C \cap M$ be such that its height below ω_1 , say n is bigger than k . We ensured this way that for the set X we will construct inductively $s_{\omega_2 \omega_3 \omega_4}(X, M \cap \omega_4)$ will be defined on the first n digits.

The first round of the game is played as follows: player I simply plays 0 three times and player II responds using his winning strategy σ_ν . His answers, say (κ_0, ϵ_0) , (λ_0, ϑ_0) and (μ_0, ϖ_0) are such that the height of ϵ_0 below ω_2 , the height of ϑ_0 below ω_3 and the height of ϖ_0 below ω_4 are less than n and thus if we let X_0 be the $cl_F(\{\kappa_0, \lambda_0, \mu_0\} \cup \nu)$ then $s_{\omega_2 \omega_3 \omega_4}(X_0, M \cap \omega_4) = o(x \setminus n, y \setminus n, z \setminus n) \upharpoonright n$ (where the sets x, y and z are defined as usual) will be $*$. Thus we start our construction without accidentally having coded some information already in the first step.

Assume now inductively that rounds $0, 1, \dots, I-1$ have already been played and that $X_{I-1} := cl_F(\{\kappa_0, \lambda_0, \mu_0, \dots, \kappa_{I-1}, \lambda_{I-1}, \mu_{I-1}\} \cup \nu)$ is such that $s_{\omega_2 \omega_3 \omega_4}(X_{I-1}, M \cap \omega_4)$ codes the previously fixed real r on the first $I-2$ many digits. Our goal is to code r 's $I-1$ -th digit. For that purpose I plays an ordinal β_i in M such that $ht_{\omega_2}(\beta_i) > \max\{ht_{\omega_3}(\vartheta_{I-1}), ht_{\omega_4}(\varpi_{I-1}), n\}$. Player II responds according to his winning strategy σ_ν with the pair (κ_I, ϵ_I) . We have by the rules of the game that $\beta_i \leq \kappa_i \leq \epsilon_i$. We now split into cases according the $(I-1)$ -th entry of r .

Case 1.

$r(I-1) = 0$. Then player I picks a γ_i such that $ht_{\omega_3}(\gamma_i) > ht_{\omega_2}(\epsilon_i)$ and player II answers with (λ_i, ϑ_i) according to σ_ν . Then I plays δ_i such that $ht_{\omega_4}(\delta_i) > ht_{\omega_3}(\vartheta_i)$ and II plays again what σ_ν tells him to, say (μ_i, ϖ_i) . By the rules of the game \mathcal{G}_ν^F we have that $\gamma_i \leq \lambda_i \leq \vartheta_i$ and $\delta_i \leq \mu_i \leq \varpi_i$. Hence the height of any point in the interval (β_i, ϵ_i) is less than the height of any point in the interval (γ_i, ϑ_i) which are less than the height of any point of the interval (δ_i, ϖ_i) . Thus when going to the images under the M collapse and constructing the according sets x, y and z we see that for $X_i := X_{I-1} \cup \{\kappa_i, \lambda_i, \mu_i\}$, $s_{\omega_2 \omega_3 \omega_4}(cl_F(X_i \cup \nu))(I-1) = 0$ as desired. As ν is a winning strategy for II the already coded information remains untouched

when passing from X_{I-1} to X_i .

Case 2.

If $r(I - 1) = 1$ then we argue very similar to case 1, we just have to ensure that we switch the heights of the ordinals in the intervals. To be more precise we make player I and II play in such a way that the height below ω_2 of any point in the interval (β_i, ϵ_i) is smaller than the height below ω_4 of any point in the interval (δ_i, ϖ_i) which in turn is smaller than the height below ω_3 of any point in the interval (γ_i, ϑ_i) . It is easy to see that such a play exists.

If I and II play the game as described above they will code the real r up to n in $n+1$ many rounds. Let $X := X_{n+1} = cl_F(\{\kappa_i, \lambda_i, \mu_i : I \leq n+1\} \cup \nu)$ then as ν is a winning strategy in \mathcal{G}_ν^F it follows that $s_{\omega_2\omega_3\omega_4}(X, M \cap \omega_4) \upharpoonright n = r \upharpoonright n$ and as we played entirely within the elementary submodel $M \prec H_\theta$ and since X is closed under F we have an F -closed witness for $\Sigma^r(M) \cap M \neq \emptyset$. Thus Σ^r is M stationary. □

Note that the coding method just described does go well with the sealing forcings we use to seal off long antichains in $P(\omega_1) \setminus \text{NS}_{\omega_1}$ as we only use them whenever the forcing is semiproper. We finally have the techniques to prove that, assuming the existence of M_1^\sharp , there is a model of ZFC where NS_{ω_1} is \aleph_2 saturated and which has a Σ_4^1 -well-order. on the reals.

3.2 Coding the reals

We briefly summarize the main results of the last section:

- (†) Given ordinals $\omega_1 < \beta < \gamma < \delta < \omega_2$ of cofinality ω_1 , there exists a proper notion of forcing $\mathbb{P}_{\beta\gamma\delta}$ such that after forcing with it the following holds: There is an increasing continuous sequence $(N_\xi : \xi < \omega_1)$ such that $N_\xi \in [\delta]^\omega$ whose union is δ such that for every limit $\xi < \omega_1$ and every $n \in \omega$ there is $\nu < \xi$ and $s_\xi^n \in 2^n$ such that

$$s_{\beta\gamma\delta}(N_\eta, N_\xi) \upharpoonright n = s_\xi^n$$

holds for every η in the interval (ν, ξ) . We say then that the triple (β, γ, δ) is stabilized.

- (‡) Further if we fix a real r there is a proper notion of forcing \mathbb{P}_r such that the forcing will produce for a triple of ordinals $(\beta_r, \gamma_r, \delta_r)$ of size and cofinality \aleph_1 a continuous, increasing sequence $(P_\xi : \xi < \omega_1)$,

$P_\xi \in [\delta_r]^\omega$ such that $\bigcup P_\xi = \delta_r$ and such that for every limit $\xi < \omega_1$ there is a $\nu < \xi$ such that

$$\bigcup_{\nu < \eta < \xi} s_{\beta_r, \gamma_r, \delta_r}(P_\eta, P_\xi) = r.$$

We say then that the real r is determined by the triple $(\beta_r, \gamma_r, \delta_r)$.

We can use these two forcing notions to set up a well-order. on the reals, which is suitable for our purposes. First we define a partial function f assigning triples of ordinals to reals:

Definition 57. For a real r we let $f(r)$ be the antilexicographically least triple (β, γ, δ) for which there is a sequence of models $(N_\xi : \xi < \omega_1)$, $N_\xi \in [\delta]^\omega$, $\bigcup(N_\xi) = \delta$ such that the set

$$\{\xi < \omega_1 : \exists \nu < \xi (\bigcup_{\nu < \eta < \xi} s_{\beta, \gamma, \delta}(N_\eta, N_\xi) = r)\}$$

is club containing (if there is such a triple).

Note that this function f is only a partial function. However we can force any given real to be in the domain of f using the proper forcing described in (‡). We use the function f to define a partial well-founded order, which will become a well-order. during our iteration :

Definition 58. Let r, s be two reals for which $f(r)$ and $f(s)$ is defined. Then we write

$$r < s \text{ iff } f(r) <_{\text{antilex}} f(s).$$

It is clear that $<$ will become a well-order. as soon as every real $r \in \omega^\omega$ is in the domain of f . The crucial property of $<$ is the following:

Lemma 59. Suppose that M is a transitive model such that every triple of ordinals $\omega_1 < (\beta, \gamma, \delta) < \omega_2$ is stabilized. Suppose further that $M \models x < y$, for two reals $x, y \in M$ then $x < y$ will hold in every stationary subset-preserving forcing extension $M[G]$ of M .

Proof. Assume for a contradiction that there is a stationary set preserving notion of forcing \mathbb{P} such that $M[G] \models y < x$, for a generic G for \mathbb{P} . Thus G must have added a continuous, increasing sequence of countable sets of ordinals $(N_\xi : \xi < \omega_1)$ such that for a triple (β, γ, δ) the set $\{\xi < \omega_1 : \exists \nu < \xi (\bigcup_{\nu < \eta < \xi} s_{\beta, \gamma, \delta}(N_\eta, N_\xi) = y)\}$ is club containing, and (β, γ, δ) is antilexicographically less than the least triple $(\beta_x, \gamma_x, \delta_x)$ for x . But we have assumed that every triple of ordinals $< \omega_2$ in M is already stabilized, I.e. there exists in M an increasing continuous sequence of countable sets of ordinals $(P_\xi : \xi < \omega_1)$ such that $\bigcup_{\xi < \omega_1} P_\xi = \delta$. By a hand by hand

argument carried out in $M[G]$ we obtain a club $C = \{\xi < \omega_1 : P_\xi = N_\xi\}$ and for every limit point ξ of C we have that $\bigcup_{\nu < \eta < \xi} s_{\beta\gamma\delta}(N_\eta, N_\xi)$ is the same as $\bigcup_{\nu < \eta < \xi} s_{\beta\gamma\delta}(P_\eta, P_\xi)$. Thus

$$M[G] \models \{\xi < \omega_1 : \bigcup s_{\beta\gamma\delta}(P_\eta, P_\xi) = y\} \text{ is club containing.}$$

But as y and $(P_\xi : \xi < \omega_1)$ are objects in M and the function $s_{\beta\gamma\delta}$ is absolute we can consider the set also in M . As $M[G]$ is a stationary set-preserving generic extension of M we have that already M sees that the set $\{\xi < \omega_1 : \exists \nu < \eta < \xi (s_{\beta\gamma\delta}(P_\eta, P_\xi) = y)\}$ is club containing in M as otherwise the complement would be stationary and therefore stationary in the extension. This is a contradiction to $M \models x < y$. \square

Thus the order $<$, once witnessed in a transitive model such that every triple of ordinals is stabilized, will not change in all outer models which preserve stationary subsets of ω_1 . This has as a consequence that we can build up the well-order. $<$ gradually during a forcing iteration and once we have a model of the just described type which sees that $x < y$, then $x < y$ in our final model of the iteration, as long as the iteration preserves stationary subsets of ω_1 . This enables us to localize the well-order. $<$, thus arriving at a projective well-order. of small complexity.

3.3 Definition of the iteration

Next we describe how to code reals nicely while making NS_{ω_1} \aleph_2 -saturated. In order to get NS_{ω_1} \aleph_2 -saturated we need an *RCS*-iteration of length κ , where κ is the Woodin cardinal. Again we let a \diamond -sequence decide what to do in our forcing iteration. We fix such a \diamond -sequence $(a_\alpha : \alpha < \kappa)$ in the ground model $V = M_1$, and let \vec{C} be the ladder system we obtain using the Σ_3^1 -well-order. of the reals of M_1 , which will still be a ladder system in small forcing extensions of M_1 which preserve ω_1 . We describe first informally how the iteration looks like. As always we have stages which are used to code information yielding the definable well-order. and stages where we seal off long antichains in $P(\omega_1)/\text{NS}_{\omega_1}$. We ensure that we code all the reals we generate during the iteration into triples of ordinals (β, γ, δ) using the proper forcing of (\ddagger) . At the same time we ensure that all the triples of ordinals below ω_2 stabilize using the forcing described in (\dagger) . Additionally whenever our \diamond -sequence hits the name of a long antichain in $P(\omega_1)/\text{NS}_{\omega_1}$ we seal it off. As we have stationarily many inaccessible cardinals below the Woodin κ we will hit stationarily often inaccessible stages α such that the model $(M_1)_\alpha[G_\alpha]$ satisfies the following:

- (1) $(M_1)_\alpha[G_\alpha] \models \text{ZF}^-$
- (2) $(M_1)_\alpha[G_\alpha] \models \forall \beta\gamma\delta < \omega_2 ((\beta, \gamma, \delta) \text{ is stabilized})$

(3) $(M_1)_\alpha[G_\alpha] \models \forall r \in \omega^\omega \exists (\beta_r, \gamma_r, \delta_r)$ (r is determined by $(\beta_r, \gamma_r, \delta_r)$).

Whenever we hit such a stage everything $(M_1)_\alpha[G_\alpha]$ sees about $<$ will be preserved in all future extensions in our iteration by Lemma 61. Thus we will additionally localize the information which is seen by $(M_1)_\alpha[G_\alpha]$ concerning $<$, i.e. for every pair of reals $x < y$ in $(M_1)_\alpha[G_\alpha]$ we add a subset $Y_{x,y}$ of ω_1 such that every countable transitive model N which contains $Y_{x,y} \cap \omega_1^N$ will also see that $x < y$. This uses a proper forcing again. As all the iterands are proper or semiproper, using an *RCS*-iteration will yield a semiproper notion of forcing. In the end we will argue that indeed NS_{ω_1} is saturated and there is a Σ_4^1 -definable well-order on the reals.

We start now with a more detailed description of how the iteration should look like. We will construct the iteration recursively, so assume that $\alpha < \kappa$ and we have already constructed \mathbb{P}_β for $\beta \leq \alpha$. We define the forcing $\dot{\mathbb{Q}}_\alpha$ in $V^{\mathbb{P}_\alpha}$ as follows:

(i) Assume that a_α is the \mathbb{P}_α -name of a real r_α . Then we let $\dot{\mathbb{Q}}_\alpha$ be the \mathbb{P}_α -name of the forcing which codes r_α into a triple of ordinals $(\beta_{r_\alpha}, \gamma_{r_\alpha}, \delta_{r_\alpha})$, such that $\beta_{r_\alpha}, \gamma_{r_\alpha}, \delta_{r_\alpha} < \omega_2$ and using the already fixed \vec{C} -sequence. This forcing is followed by considering all the triples of ordinals $(\beta', \gamma', \delta')$ which are antilexicographically below $(\beta_{r_\alpha}, \gamma_{r_\alpha}, \delta_{r_\alpha})$ and which have not been stabilized yet. We use an *RCS*-iteration of forcings which stabilize each such triple $(\beta', \gamma', \delta')$. As a summary $\dot{\mathbb{Q}}_\alpha$ is an ω_1 -long *RCS* iteration of proper forcings resulting in a proper forcing, and we obtain a model where the real r_α is coded into the triple $(\beta_{r_\alpha}, \gamma_{r_\alpha}, \delta_{r_\alpha})$ with the help of the ladder system \vec{C} , and each other triple of ordinals below it will be stabilized.

Otherwise force with $\text{Col}(2^{\aleph_2}, \aleph_1)$, the usual Lévy collapse which collapses 2^{\aleph_2} down to \aleph_1 .

(ii) Assume that α is an inaccessible, further that a_α is the \mathbb{P}_α -name of a maximal antichain S_α in $P(\omega_1)/\text{NS}_{\omega_1}$, and assume that the sealing forcing $\mathbb{S}(S_\alpha)$ is semiproper. Then force with it, i.e. let $\dot{\mathbb{Q}}_\alpha$ be $\mathbb{S}(S_\alpha)$.

Otherwise force with $\text{Col}(2^{\aleph_2}, \aleph_1)$.

(iii) If α is an inaccessible and if $(M_1)_\alpha[G_\alpha]$ is a model such that property (1) (2) and (3) from above holds, then we first collapse its size down to \aleph_1 . We consider a pair of reals $x, y \in (M_1)_\alpha[G_\alpha]$ such that $(M_1)_\alpha[G_\alpha] \models x < y$ and construct a set $Y_{x,y} \subset \omega_1$ which should code in a nice way the information that $(M_1)_\alpha[G_\alpha] \models x < y$ (this will be specified below). Now code the set $Y_{x,y}$ into a real $s_{x,y}$ using almost disjoint coding forcing relative to an almost disjoint family of reals in M_1 . We pick the pair $(x < y)$ in such a way that every such pair we create during our iteration will be considered at some inaccessible stage α . This can be easily achieved using some bookkeeping function.

The points (i) and (ii) are clear, thus we shall discuss (iii) in detail: Note first that it is straightforward to code the previously fixed ladder system \vec{C} into a subset X of ω_1 such that for every limit ordinal $\xi < \omega_1$, if we decode $X \cap \xi$ we end up with $\vec{C} \upharpoonright \xi$. Assume now that we are in the situation described in (iii), thus α is an inaccessible and $(M_1)_\alpha[G_\alpha]$ is a model of ZF^- , (1),(2) and (3). We collapse its size to \aleph_1 using Lévy-collapse and let H be the generic filter. Let x, y be two reals in $(M_1)_\alpha[G_\alpha]$ such that $(M_1)_\alpha[G_\alpha] \models x < y$. We describe now the set $X_{x,y} \subset \omega_1$ which codes the information:

Claim 60. *In $M_1[G_\alpha][H]$ there is a set $X_{x,y} \subset \omega_1$ such that $X_{x,y}$ can be recursively partitioned into 6 subsets such that the following holds:*

1. $\text{dec}_1(X_{x,y}) = \vec{C}$
2. $\text{dec}_2(X_{x,y}) = x$
3. $\text{dec}_3(X_{x,y}) = y$
4. $\text{dec}_4(X_{x,y})$ is the set consisting of the triple of ordinals $(\beta_x, \gamma_x, \delta_x)$ and the continuous sequence of models $(N_\xi^x : \xi < \omega_1)$ such that $(\beta_x, \gamma_x, \delta_x)$ is the antilexicographically least triple which codes x with the help of \vec{C} and witnessing sequence $(N_\xi^x : \xi < \omega_1)$
5. $\text{dec}_5(X_{x,y})$ is the set similar defined as in the previous point but with y instead of x .
6. $\text{dec}_6(X_{x,y})$ is the set of all the continuous increasing sequences $(N_\xi^{(\beta', \gamma', \delta')} : \xi < \omega_1)$ for $(\beta', \gamma', \delta') < (\beta_x, \gamma_x, \delta_x)$ of sets of ordinals which witness that the triple $(\beta', \gamma', \delta')$ stabilizes at a real which is neither x nor y .

The construction of such a set $X_{x,y}$ is straightforward. As a consequence, for every transitive model M of ZFC which contains $X_{x,y}$, M will see that $x < y$ as it contains all the relevant information. The goal now is to rewrite $X_{x,y}$ into a set $Y_{x,y} \subset \omega_1$ such that every countable, transitive M which contains $Y_{x,y} \cap \omega_1^M$ already sees that $x < y$ using the local ladder $\vec{C} \upharpoonright \omega_1^M$. We can force the existence of such a set $Y_{x,y}$ with a proper notion of forcing.

Lemma 61. *There is a proper notion of forcing \mathbb{R} which introduces a set $Y_{x,y} \subset \omega_1$ such that if H' is an \mathbb{R} generic filter $M_1[G_\alpha][H][H']$ satisfies that there is subset $Y_{x,y}$ of ω_1 such that if $\xi = \omega_1^N$ for a countable transitive model N then there are recursively definable decoding functions dec_i such that the following holds:*

1. $\text{dec}_1(Y_{x,y}) = \vec{C}$ and for every limit ordinal $\xi < \omega_1$, with $\xi \omega = \xi$, $\text{dec}_1(Y_{x,y} \upharpoonright \xi) = \vec{C} \upharpoonright \xi$.
2. $\text{dec}_2(Y_{x,y}) = x$

3. $dec_3(Y_{x,y}) = y$
4. $dec_4(Y_{x,y} \upharpoonright \xi)$ is the set consisting of a triple of ordinals $(\beta_x^\xi, \gamma_x^\xi, \delta_x^\xi)$ and a continuous sequence of sets of ordinals $(N_I^{x,\xi} : I \in \xi)$ such that x is coded by the triple $(\beta_x^\xi, \gamma_x^\xi, \delta_x^\xi)$ using the local ladder system $\vec{C} \upharpoonright \xi$ and witnessed by $(N_I^{x,\xi} : I \in \xi)$.
5. $dec_5(Y_{x,y})$ is defined similar as 4 but with x replaced by y .
6. $dec_6(X_{x,y})$ is the set of all the continuous increasing sequences $(N_I^{\xi,(\beta',\gamma',\delta')}) : I < \xi$ for $(\beta', \gamma', \delta') < (\beta_x^\xi, \gamma_x^\xi, \delta_x^\xi)$ of sets of ordinals which witness that the triple $(\beta', \gamma', \delta')$ stabilizes at a real which is neither x nor y using the local ladder $\vec{C} \upharpoonright \xi$.

Proof. Working in $M_1[G_\alpha][H]$ we have that $(M_1)_\alpha[G_\alpha]$ is a model of ZF^- and (1) and (2) of size \aleph_1 . By the previous Claim we know that there is a set $X_{x,y}$ such that all the points in the claim are true. Fix a model of the form $(M_1)_\eta[G_\alpha][H]$ for $\eta > \alpha$ which contains $X_{x,y}$ and consider the club C of countable, elementary submodels containing $X_{x,y}$. We can easily arrange that $X_{x,y}$ satisfies already (1) of the Claim. For M an arbitrary element of C the following holds:

$$(M_1)_\eta[G_\alpha][H] \models X_{x,y} \text{ satisfies (1) – (6) of the previous Claim}$$

thus for each $M \in C$:

$$M \models X_{x,y} \text{ satisfies (1) – (6)}$$

thus

$$\bar{M} \models \bar{X}_{x,y} \text{ satisfies (1) – (6)}$$

But $\bar{X}_{x,y} = X_{x,y} \upharpoonright \omega_1^M$ and as the decoding functions dec_i are absolute for transitive models we get that $X_{x,y} \upharpoonright \xi$ satisfies (1)-(6) for ξ 's which are the ω_1 of some $M \in C$. In order to get the full statement of the Claim we add additional information to $X_{x,y}$ which yields $Y_{x,y}$ such that any countable transitive model N which contains $Y_{x,y} \cap \omega_1^N$ must have its ω_1 to be an ω_1^M for some $M \in C$. To achieve this we use forcing.

Let \mathbb{R} be the following partial order: conditions $p \in \mathbb{R}$ are ω_1 -Cohen conditions, i.e. functions from limit ordinals $\xi < \omega_1$ to 2 which satisfy:

1. the even ordinals of $\{\eta < \xi : p(\eta) = 1\}$, where $\xi = dom(p)$ code the set $X_{x,y} \cap \xi$.
2. for every limit ordinal ξ , if M is countable and transitive and $\xi = \omega_1^M$ and $(p \upharpoonright \xi) \in M$ then $p(\xi)$ seen as a subset of ξ satisfies points (1) – (6) for every limit $\eta < \xi$.

Note that whenever we do have a condition $p \in \mathbb{R}$, and $\xi < \omega_1$ is a limit ordinal we can extend p to $q < p$ such that $\xi \in \text{dom}(q)$. This is clear as we can write into the first odd ω -block of q following $\text{dom}(p)$ a surjection of ξ to ω . Then no countable transitive model M which contains q can have its ω_1 at ξ , thus the second property for being a condition in \mathbb{R} is satisfied automatically. Thus the set $D_\eta := \{p \in \mathbb{R} : \eta \in \text{dom}(p)\}$ is dense for every $\eta < \omega_1$ and the generic will produce a subset of ω_1 , $Y_{x,y}$ with the desired properties (1) – (6) for countable, transitive models in $M_1[G_\alpha][H]$. This already suffices as we will see below that the forcing \mathbb{R} is also ω -distributive.

What is left is to show that the forcing \mathbb{R} is proper: for that we pick the $(M_1)_\eta[G_\alpha][H]$ from above and let the club C be the set of all countable elementary submodels of it which contain the set $X_{x,y}$ from above. If $M \in C$, and $p \in \mathbb{R} \cap M$ then we shall construct a $q < p$ which is (M, \mathbb{R}) -generic. We list all the dense sets D_n in M and recursively construct a descending sequence of conditions starting at $p = p_0 > p_1 > \dots$ such that $p_n \in D_n$ and such that $\text{sup}(\text{dom}(p_n)) = \omega_1 \cap M$. If we can show that the limit p_ω is a condition in \mathbb{R} , we are done. Thus we have to show that whenever $p_\omega \cap \xi$ is contained in a countable, transitive model N such that $\xi = \omega_1^N$ then it will satisfy conditions (1) – (6). This is clear by definition of \mathbb{R} for every $\xi < \text{dom}(p_\omega)$. If $\xi = \text{dom}(p_\omega)$ then as $\xi = \omega_1^M$, $M \in C$ and p_ω codes $X_{x,y} \cap \xi$ we know by the above that $X_{x,y} \cap \xi$ satisfies (1) – (6), and so does $Y_{x,y} \cap \xi$ if we change the decoding functions in the obvious way. Thus p_ω is a condition in \mathbb{R} as desired and the forcing \mathbb{R} is proper.

Note that the same argument shows that \mathbb{R} is also ω -distributive, and so the added set $Y_{x,y}$ has the desired properties not only for countable sets in $M_1[G_\alpha][H]$ but also for sets in $M_1[G_\alpha][H][H']$ as desired. □

Now that we have constructed the localized set $Y_{x,y}$ for two reals $x < y$ in $(M_1)_\alpha[G_\alpha]$ we code $Y_{x,y} \subset \omega_1$ into a real $r_{x,y}$ using almost disjoint coding. We fix an M_1 -family $(r_\alpha : \alpha < \omega_1)$ of almost disjoint reals and add a real $r_{x,y}$ such that the following holds:

$$\forall \xi < \omega_1 (\xi \in Y_{x,y} \text{ iff } r_{x,y} \cap r_\xi \text{ is almost disjoint.})$$

This forcing is ccc, therefore proper and for the real $r_{x,y}$ the following holds:

- ♡ Every countable, transitive model M which contains $r_{x,y}$ and $(M_1)_{\omega_1^M}$ sees that $x < y$ using the ladder system $\vec{C} \upharpoonright \omega_1^M$.

This ends the discussion of the forcing we use in case (iii).

3.4 NS_{ω_1} is saturated and a projective well-order. Open Questions.

In this section we show that the model we obtain following the definition given in the previous section will indeed satisfy that NS_{ω_1} is saturated and there is a Σ_4^1 definable well-order. on the reals. We start with the well-order. first.

Lemma 62. *Let G be a generic filter for the forcing defined in the last section. Then*

$$M_1[G] \models \text{there is a } \Sigma_4^1 \text{ definable well-order. on the reals.}$$

Proof. We can use the cofinal set of M_1 -initial segments in $M_1[G]$ which is Π_2^1 definable to construct a Σ_3^1 -definable ladder system \vec{C} in $M_1[G]$. Simply let $(\alpha, C_\alpha) \in \vec{C}$ if and only if there is a countable M_1 initial segment M which contains (α, C_α) and which sees that C_α is the $<_M$ -least set in M (where $<_M$ denotes the usual definable well-order. on the mouse M) which is cofinal and has ordertype ω . This definition is Σ_3^1 and we let $\psi(x)$ be the defining formula, I.e. for every $C \subset \omega_1$ we have that $\psi(C)$ if and only if $C \in \vec{C}$. We fix the ladder system \vec{C} defined via ψ and code, using the machinery described and explained above, everything relative to it. We claim that the following defines the well-order. $<$ in $M_1[G]$:

- (*) $x < y$ if and only if $\exists r \forall M (M \text{ countable and transitive and } (M_1)_{\omega_1^M} \in M \text{ then } \psi(\text{dec}(r, (M_1)_{\omega_1^M})) \text{ and } M \models x < y \text{ using the local ladder system } \vec{C} \cap \omega_1^M)$.

Here we write $(\text{dec}(r, (M_1)_{\omega_1^M}))$ for the set we obtain when we decode the real using the almost disjoint family of $(M_1)_{\omega_1^M}$ -reals and obtain a subset of ω_1^M . Thus the statement $\psi(\text{dec}(r, (M_1)_{\omega_1^M}))$ tells us that the ladder system coded into the real is the previously fixed ladder system \vec{C} relativized to ω_1^M . The direction from left to right is clear as whenever $x < y$ in $M_1[G]$ then the names of x and y will appear at some earlier stage $\beta < \kappa$ in our iteration. Thus $x < y$ will be witnessed already at some inaccessible stage $\alpha < \kappa$ as we used a bookkeeping function which looks at each name of a pair of reals unboundedly often on the inaccessibles below κ . Note that in this case the definition of our iteration guarantees us that we have added a real $r_{x,y}$ such that \heartsuit holds which is exactly what we want.

For the direction from right to left note first that the right hand side of (*) holds also for large enough uncountable models M containing r and $(M_1)_{\omega_1}$ and which satisfy that each triple of ordinals below ω_2^M is stabilized. Indeed if we assume that M is uncountable, $r \in M$ and $(M_1)_{\omega_1} \in M$ and every triple of ordinals is stabilized and M does decide that $y < x$, then a

countable elementary submodel $N \prec M$ containing r will see as well that $y < x$ using the ladder system which is coded into r . This ladder system must be our fixed \vec{C} . But now the transitive collapse $\pi(N) = \bar{N}$ contains r and we can assume that also $(M_1)_{\omega_1^{\bar{N}}}$ is in \bar{N} . It will still think that $y < x$ using the ladder system $\pi(\vec{C}) = \vec{C} \cap \omega_1^{\bar{N}}$, but \bar{N} is just as in the assumption for the right hand side for $(*)$. This gives us the contradiction. as \bar{N} cannot see both $x < y$ and $y < x$. Thus if we let M be big enough such that every triple of ordinals below ω_2 is stabilized then it will see that $x < y$. But by Lemma 61 $x < y$ must hold in $M_1[G]$.

Note further that the definition of the well-order. is of the form $\exists r \forall M (\Pi_2^1 \rightarrow \Pi_3^1)$ and therefore Σ_4^1 . □

What is left is to show that in $M_1[G]$ the nonstationary ideal NS_{ω_1} is indeed \aleph_2 -saturated. But this does not cause any problems as the coding forcings were all seen to be proper, the sealing forcings were only used when semiproper and we used *RCS*-iteration for the limit steps. Therefore the iteration yields a semiproper, thus stationary set preserving extension of M_1 and we can just repeat Shelah's proof that NS_{ω_1} is \aleph_2 -saturated in the final model. Hence the following is true, which ends the proof of the theorem:

Lemma 63. *If G denotes the generic filter for the iteration then in $M_1[G]$ the nonstationary ideal NS_{ω_1} is \aleph_2 -saturated.*

We end this thesis with a couple of remarks. At first glimpse it might seem that the techniques introduced in the second chapter, i.e. isolating a suitable class of models \mathcal{S} and use K -definable objects to code patterns would as well yield a solution to the problem of projectively definable wellorders on the reals in the presence of NS_{ω_1} saturated. Indeed this was our strategy for quite some time, it turned out however that this approach is fruitless. The reason for this is that sealing the antichains in $P(\omega_1) \setminus NS_{\omega_1}$ is a semiproper forcing only, hence our stationary class \mathcal{S} will lose its stationarity along the iteration. This fact compels us to work with the augmented version of \mathcal{S} , namely $\mathcal{S}[G]$, for G the generic filter. But models in $\mathcal{S}[G]$ are of the form $M[G]$, i.e. M with all the names in it evaluated with the help of G . These models $M[G]$ are not necessarily generic extensions of M , thus defining K in them causes utter chaos, although K can be defined in $M \in \mathcal{S}$ nicely. This problems led me to consider other methods of coding which finally resulted in the proof described in this chapter. Note however that the methods developed here do not seem to get an easy proof of the following natural extension of the problem in chapter 2:

Question 1. *How to get a model of ZFC in which NS_{ω_1} is \aleph_2 -saturated and the full nonstationary ideal NS_{ω_1} is δ_1 -definable with parameter ω_1 ?*

Another natural question is whether the δ_4^1 wellorder is optimal while having NS_{ω_1} saturated. By the results of Hjorth and Woodin, under the assumption that there is a measurable cardinal, the Σ_4^1 wellorder is optimal but are there ways to do better in the absence of a measurable?

Question 2. *Is there a model of ZFC in which NS_{ω_1} is saturated and there is a Δ_3^1 -definable wellorder on the reals?*

and finally the very interesting problem

Question 3. *Is it possible to have a model of ZFC in which NS_{ω_1} is saturated and CH holds?*

Bibliography

- [1] A. Caicedo and S.D. Friedman *BPFA and Projective Wellorderings of the Reals* Journal of Symbolic Logic, pp. 1126–1136, 2011.
- [2] A. Caicedo and B. Velickovic *The bounded proper forcing axiom and well-orderings of the reals* Mathematical Research Letters, 13 (2-3) (2006). pp. 393-408
- [3] V. Fischer and S.D. Friedman *Cardinal characteristics and projective wellorders* Annals of Pure and Applied Logic 161, pp. 916-922, 2010.
- [4] S.D. Friedman *Fine Structure and Class Forcing* De Gruyter, 2000.
- [5] S.D. Friedman and Liuzhen Wu *Large cardinals and the Δ_1 -definability of the nonstationary ideal* To appear.
- [6] L. Harrington *Long Projective Wellorderings* Ann. Math. Logic 12, pp. 1-24, 1977.
- [7] G. Hjorth *The size of the ordinal u_2* J. London Math. Soc.(2), 52 (3), pp. 417-433, 1995.
- [8] A. Kanamori *The Higher Infinite. Second Edition* Springer, 2003.
- [9] B. Löwe and J. Steel *An Introduction to Core Model theory* Sets and Proofs, Cambridge University Press, pp. 103-159.
- [10] D.A. Martin and J.R. Steel *Iteration trees* J. Amer. Math. Soc. 7, no.1, pp.1-73, 1994.
- [11] W. Mitchell and J. Steel *Fine Structure and Iteration Trees* Lecture Notes in Logic, Springer, 1994
- [12] J. Moore *Set Mapping Reflection* Journal of Mathematical Logic, 5(1),pp. 87-98, 2005.
- [13] J.Steel *An Outline of Inner Model Theory* Handbook of Set Theory, Springer, pp. 1595-1685, 2011.

- [14] *Inner models with many Woodin cardinals* Annals of Pure and applied Logic 65, pp.185-209, 1993.
- [15] J. Steel *Projectively well-ordered Inner Models* Annals of pure and applied Logic, pp.77-104, 1995.
- [16] R. Schindler *On NS_{ω_1} being saturated* Online Notes, http://www.math.uni-muenster.de/u/rds/sat_ideal_better_version.pdf 20.6.2016.
- [17] R. Schindler *Core models in the presence of Woodin cardinals*, J. Symb. Logic 71, pp. 1145-1154, 2006.
- [18] E. Schimmerling *A Core Model Toolbox and Guide* Handbook of Set Theory, Springer, pp. 1685-1753.
- [19] S. Shelah *Proper and Improper Forcing* Springer, 1998.
- [20] M. Viale, G. Audrito and S. Steila *A Boolean algebra approach to semiproper iterations* Online Notes, http://www.logicatorino.altervista.org/matteo_viale/semiproperforcing.pdf 20.6.2016.
- [21] H.W. Woodin *The Axiom of Determinacy, Forcing Axioms, and the Nonstationary Ideal* De Gruyter, 2001.