

Heterotrophy among Cyanobacteria

Published as part of the ACS Omega virtual special issue "Phytochemistry".

Ronald Stebegg,* Georg Schmetterer, and Annette Rompel*



Cite This: *ACS Omega* 2023, 8, 33098–33114



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Cyanobacteria have been studied in recent decades to investigate the principle mechanisms of plant-type oxygenic photosynthesis, as they are the inventors of this process, and their cultivation and research is much easier compared to land plants. Nevertheless, many cyanobacterial strains possess the capacity for at least some forms of heterotrophic growth. This review demonstrates that cyanobacteria are much more than simple photoautotrophs, and their flexibility toward different environmental conditions has been underestimated in the past. It summarizes the strains capable of heterotrophy known by date structured by their phylogeny and lists the possible substrates for heterotrophy for each of them in a table in the Supporting Information. The conditions are discussed in detail that cause heterotrophic growth for each strain in order to allow for reproduction of the results. The review explains the importance of this knowledge for the use of new methods of cyanobacterial cultivation, which may be advantageous under certain conditions. It seeks to stimulate other researchers to identify new strains capable of heterotrophy that have not been known so far.



1. INTRODUCTION

Cyanobacteria have evolved to a photolithoautotrophic growth mode^{1,2} because they use light as an energy source (photo-trophic), the inorganic water molecule as an electron source (lithotrophic) for NADPH production, and carbon dioxide, that has only one carbon atom, as the sole carbon source (autotrophic) (the different types of trophies and their possible combinations are explained in Table 1). This implies that autotrophic organisms must metabolically form all covalent bonds between carbon atoms. Unlike other phototrophic prokaryotes such as purple bacteria and green sulfur bacteria, which perform an anoxygenic mode of photosynthesis and have only one photosystem, cyanobacteria are defined as prokaryotes capable of oxygenic photosynthesis and possess two photosystems (I, for a review about photosystem I, please see ref 3, and II, for a review about photosystem II, please see ref 4). Oxygenic photosynthesis requires two distinct photosystems to ensure sufficient reduction potential for water splitting (removal of electrons from the very electronegative oxygen and the release of protons and molecular dioxygen) on one hand and the production of enough ATP needed for carbon fixation in the Calvin cycle on the other. In eukaryotes such as algae and land plants, only oxygenic photosynthesis takes place, which consists of two photosystems. This fact is consistent with the endosymbiotic theory⁵ that the chloroplasts of algae and plants descended from cyanobacteria that invaded eukaryotic cells, the progenitor of modern algae and plants.

Due to their production of free molecular oxygen, cyanobacteria were among the first evolutionary organisms to protect themselves against radical oxygen species (ROS). As a result, cyanobacteria evolved aerobic respiration in addition to their oxygenic photosynthesis. The oxygen and organic molecules that have been formed in the light can be utilized for respiration in the dark. In contrast to eukaryotic algae and embryophytes, where these two processes are spatially separated in chloroplasts and mitochondria, they take place in the same compartment in cyanobacteria.⁶

While all cyanobacteria, eukaryotic algae, and land plants can survive in periods of darkness by aerobic respiration, few strains of cyanobacteria^{7–9} and eukaryotic algae^{10–12} but no plants are able to grow in prolonged darkness, where carbon dioxide can no longer be fixed since NADPH and ATP production only occurs during the light reaction. In the dark, organic molecules are used as a source of carbon, energy, and electrons, and this growth mode is called chemoorganoheterotrophic growth (chemotrophic because energy is obtained through chemical reactions, organotrophic because the electrons are derived from organic molecules, and heterotrophic because organic substrates

Received: April 2, 2023

Accepted: July 10, 2023

Published: September 6, 2023




Table 1. Various Combinations of Trophies of Naturally Occurring Organisms

Growth mode	Energy source	Electron source	Carbon source
photolithoautotrophic	light	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S, H ₂ O)	molecules consisting of only one carbon atom (e.g., CO ₂ , HCOOH, CH ₃ OH, CH ₄)
photolithoheterotrophic	light	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S, H ₂ O)	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)
chemolithoautotrophic	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S)	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S)	molecules consisting of only one carbon atom (e.g., CO ₂ , HCOOH, CH ₃ OH, CH ₄)
chemolithoheterotrophic	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S)	inorganic compounds (e.g., H ₂ , Fe ²⁺ , NH ₃ , H ₂ S)	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)
photoorganoheterotrophic	light	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)
chemoorganoheterotrophic	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)	organic molecules consisting of at least two carbon atoms (e.g., glucose, fructose, sucrose, glycerol, ...)
photomixotrophic	light	inorganic and organic molecules	molecules consisting of one carbon atom and molecules consisting of at least two carbon atoms

normally used for growth consist of at least two carbon atoms, as illustrated in Table 1). In the following sections, the terms photolithoautotrophic and chemorganoheterotrophic are abbreviated to photoautotrophic and chemoheterotrophic. Normally, cyanobacteria capable of chemoheterotrophic growth can use only one or two distinct organic molecules as a carbon source, and the substrates vary between cyanobacterial strains. The most prevalent substrates are sugars such as glucose, fructose, and sucrose and the alcohol glycerol.^{7–9,13–23} This is consistent with the fact that the genomes of many strains of cyanobacteria contain genes that are either responsible for the uptake of sugars or for their metabolism.²⁴ Import through the outer membrane is more likely to occur through nonspecific porins,²⁵ while for transport through the cytoplasmic membrane, homologues of the glucose transporter Glc in *Synechocystis* sp. PCC 6803²⁶ and the fructose transporter FtrRABC in *Anabaena variabilis* ATCC 29413²⁷ are widely distributed.²⁸ Figure 1 summarizes various substrates that are imported and metabolized by cyanobacteria.

Synechocystis sp. PCC 6803 is capable of a particular form of chemoheterotrophy, as it requires 5 min of daily illumination for glucose-dependent dark growth. Nevertheless, this growth mode can be considered as a kind of chemoheterotrophic growth since it is strictly dependent on the presence of glucose. This modified growth mode is termed light-activated heterotrophic growth (LAHG).²⁹ A similar case is observed for *Synechococcus* sp. PCC 7002. This strain can grow in dim light (40 μW cm⁻²) in the presence of 55 mM glucose but not in total darkness at the same glucose concentration.³⁰ Since a light intensity of 40 μW cm⁻² itself does not induce (photoautotrophic) growth, it may be true (light-activated) heterotrophic growth.

Some strains of cyanobacteria are strictly light dependent but can also grow in the absence of photosystem II. If no water is split due to the lack of photosystem II, organic molecules serve as the electron source. Since photosystem I is still active, light can be considered as the energy source; however, the ATP originating from photosystem I is not sufficient for carbon dioxide fixation, and an organic electron source is additionally used for carbon assimilation. Therefore, this growth mode is termed photoorganoheterotrophic growth or, more briefly, photoheterotrophic growth.^{8,9,23} In cyanobacteria, photoheterotrophic conditions arise naturally through spontaneous mutations in genes encoding photosystem II subunits, but these mutations can also be engineered.^{31–33} Alternatively, photoheterotrophy can be induced by the herbicide 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), which blocks electron transfer from photosystem II to the quinone pool.^{34–37}

Some strains of cyanobacteria are unable to grow heterotrophically, so they cannot grow in the absence of light or either of their two photosystems; however, their photoautotrophic growth rate is further increased by the addition of organic molecules to their growth medium. This mode is called photomixotrophic growth. Although photomixotrophic growth is not a true form of heterotrophy, we will discuss it in this review because this growth mode is based on at least a partial growth dependence on organic molecules.

In all of the above cases, the appropriate substrate must be imported into the cell (for a review of the transport of organic substances across the cytoplasmic membrane, please see ref 38); however, strict photoautotrophy is not always caused by a lack of transport alone, as the entering molecule has to be metabolized in some way for the synthesis of ATP and NAD(P)H without accumulation of any toxic (by-)products.³⁹




Figure 1. Overview of uptake and metabolism of diverse organic molecules by a cyanobacterial cell. GlpK, glycerol kinase; XylA, xylose isomerase; XylB, xylulokinase; Glk, glucokinase; CscA/InvA, invertase; CscK, fructokinase (from *E. coli*). The “?” at the glycerol transporter implies that this transporter and the corresponding gene in *Cyanothece* sp. ATCC 51142 have not been identified yet. The “?” after the (native cyanobacterial) fructokinase indicates that no gene within the genome of either *Anabaena* sp. ATCC 29413 or *Nostoc* sp. ATCC 29133 has been annotated as fructokinase so far.²⁴

In this review, we summarize and analyze all available data on the different forms of heterotrophic (photomixotrophic, photoheterotrophic, or chemoheterotrophic) growth in cyanobacteria. In the past decades, there has been a strong focus on autotrophy, while little is known about facultative heterotrophy, as this topic has been rather neglected. The aim of this review is to expand knowledge about the possibility of heterotrophic cultivation of cyanobacterial strains that were previously considered strictly photoautotrophic in order to stimulate new research topics with cultivation forms that were previously not thought possible. Overall, the picture of cyanobacteria will change from purely photosynthetic organisms to highly flexible organisms that can adapt their metabolism to new circumstances when environmental conditions change.

2. CYANOBACTERIAL STRAINS CAPABLE OF HETEROTROPHIC GROWTH

The most comprehensive study of facultative photoheterotrophy among almost all cyanobacterial strains known at the time has been carried out by Rippka et al.²³ Here we discuss cyanobacterial strains in alphabetical order following the sections of Rippka et al.²³ since most of the physiological

studies on chemoheterotrophy were performed decades ago; however, we slightly modified the system as we have combined sections IV and V since they form one common phylogenetic group.⁴⁰ The strains described are mostly preserved in the Pasteur Culture Collection (PCC), the American Type Culture Collection (ATCC), the Culture Collection of Algae at the University of Texas (UTEX), or the Culture Collection of Algae and Protozoa (CCAP), and the underlying collection is part of the name of the strains (abbreviation followed by a number). Unless otherwise stated, all strains are cultivated in the BG11 medium (invented by Rippka et al.²³), containing 0.18 mM $K_2HPO_4 \cdot 3H_2O$, 0.3 mM $MgSO_4 \cdot 7H_2O$, 0.24 mM $CaCl_2 \cdot 2H_2O$, 17.6 mM $NaNO_3$, 2.8 μM $Na_2MgEDTA$, 22.8 μM ferric ammonium citrate, 31 μM citric acid, 0.19 mM Na_2CO_3 , 46 μM H_3BO_3 , 1.6 μM $Na_2MoO_4 \cdot 2H_2O$, 9.7 μM $MnCl_2 \cdot 4H_2O$, 0.77 μM $ZnSO_4 \cdot 7H_2O$, 0.32 μM $CuSO_4 \cdot 5H_2O$, and 0.17 μM $Co(NO_3)_2 \cdot 6H_2O$. Table S1 (in the Supporting Information) lists the strains known to date capable of heterotrophic growth and the organic substrate(s) used therefore, if known.

2.1. Section I: Chroococcales. Chroococcales are defined as unicellular cyanobacteria that divide by binary fission.²³

2.1.1. *Chamaesiphon*. PCC 7430 can grow photoheterotrophically on glucose, fructose, and sucrose.²³

2.1.2. *Cyanothece (Crocospaera subtropica)*. Reddy et al.⁴¹ tested possible substrates for supporting photoheterotrophic growth of two *Cyanothece* sp. strains BH63 and BH68, which have been assigned by the ATCC as *Cyanothece* sp. strains ATCC 51141 and ATCC 51142. While glucose, fructose, and sucrose do not exert any beneficial effect, 10 mM glycerol allows photoheterotrophic growth with 20 μ M DCMU in both the presence and absence of nitrate. Photoheterotrophic cultures in the absence of nitrate reach the stationary phase after 3 days, whereas they continue to grow at least for 6 days in the presence of nitrate.⁴¹ If the cells are incubated in diurnal 12 h light/12 h dark cycles, the nitrogenase was only active during the dark phase; however, even in continuous light its activity alternates in a 24 h cycle.^{41,42} In *Cyanothece* sp. ATCC 51142 the *nifHDK* genes encoding the subunits of the nitrogenase are exclusively expressed during the darkness in a 12 h light/12 h dark cycle or in regular intervals in a 24 h cycle when continuously illuminated. While the NifH subunit (Fe protein) is consequently degraded within this cycle regardless of illumination, the NifDK subunit (MoFe protein) is only completely degraded in 12 h light/12 h dark cycles but not in continuous light.⁴²

The genome of *Cyanothece* sp. ATCC 51142 contains many genes that are regulated within a diurnal cycle^{43,44} and genes involved in glucose and pyruvate metabolism.⁴⁵ Therefore, Feng et al.⁴⁶ checked photomixotrophic growth of *Cyanothece* sp. ATCC 51142 on glucose, pyruvate, and glycerol under continuous illumination. In the absence of bound nitrogen, only 54 mM glycerol significantly increased the growth rate compared to photolithoautotrophic conditions, while the positive effect of 26 mM glucose can be neglected. Low concentrations of sodium pyruvate (9 mM) do not change the growth behavior for the first 6 days and then exhibit a deleterious effect, whereas higher concentrations (64 mM) cause a growth inhibition from the beginning. Glycerol doubles the growth rate of *Cyanothece* sp. ATCC 51142 in light both in the presence and in the absence of 18 mM sodium nitrate. The addition of 54 mM glycerol or 26 mM glucose increases the nitrogen-dependent hydrogen production during the exponential growth phase in continuous light by a factor of 5.0 and 2.6, respectively. 11 mM sodium pyruvate on the other hand slightly reduces the hydrogen production by a factor of 0.8.⁴⁶

Isotope (¹³C) labeling reveals that under mixotrophic conditions alanine, histidine, and serine are exclusively built from glycerol as the only carbon source. The authors therefore call it photoheterotrophic metabolism since carbon dioxide does not contribute to the synthesis of the three amino acids in this experiment; however, we consider it to be photomixotrophic growth, as PSII has not been blocked (e.g., by DCMU). In the absence of sodium nitrate the portion of carbon derived from glycerol within alanine, histidine, and serine is dramatically reduced. Interestingly glucose and pyruvate also contribute to the formation of amino acids (i.e., they are metabolized within the cell), although they do not significantly increase the growth rate, although the substance does not support growth.⁴⁶ Recently the *Cyanothece* sp. ATCC 51142 has been renamed to *Crocospaera subtropica* ATCC 51142.⁴⁷

2.1.3. *Gloeocapsa*. *Gloeocapsa* sp. strain PCC 7428 grows photoheterotrophically on glucose, fructose, ribose, and sucrose, whereas *Gloeocapsa* sp. strain PCC 7501 can only do so on glucose.²³

2.1.4. *Synechococcus*. For a review of photomixotrophic growth of marine *Synechococcus* see Munoz-Marín et al.⁴⁸ Duhamel et al.⁴⁹ studied photomixotrophic metabolism in

marine undefined *Synechococcus* sp. strains that (along with *Prochlorococcus* sp.) contribute to most of the marine photosynthesis.^{50–52} *Synechococcus* sp. assimilates glucose, leucine, and ATP at low rates and molecules containing nitrogen and phosphorus at a higher rate. Molecular assimilation is higher in light under photomixotrophic conditions than in the dark under heterotrophic conditions or when photosystem II is inhibited.⁴⁹

Synechococcus sp. PCC 7002 (designated *Agmenellum quadruplicatum* PR-6 in the publication by Van Baalen⁵³) grows (light activated) heterotrophically in the presence of 44–55 mM (and to a lesser extent also at 11 mM) glucose, if weak light (40 μ W cm⁻²) is present; however, no growth occurs at the same glucose concentration in total darkness. Under high light conditions that allow for normal photoautotrophic growth, glucose exerts no stimulating effect. As a result, the contribution of carbon derived from glucose incorporated into amino acids is higher under dim light than under strong light. Acetate fails to induce the heterotrophic growth of *Agmenellum quadruplicatum* PR-6 in dim light.³⁰ In addition, *Synechococcus* sp. PCC 7002 grows photoheterotrophically in the presence of DCMU on glycerol and to a lesser extent also on glucose.²³ Lambert and Stevens⁵⁴ have also tested several substrates for photoheterotrophic growth. Fructose, sodium malate, sodium citrate, and glucose fail in this study (both at 10 and 50 mM), while growth occurs on heavily inoculated plates supplemented with glycerol (1–100 mM) and DCMU. Nevertheless, the pigmentation of the cells is reduced at 10 mM or higher concentrations of glycerol; however, dark green cells with normal pigments have formed on plates supplemented with 30 or 100 mM glycerol. This glycerol-tolerant strain named *Agmenellum quadruplicatum* PR-6G2 does not differ much under photoautotrophic or photomixotrophic conditions. PR-6 and PR-6G2 grow at generation times of 4.8 and 4.7 h (photoautotrophic conditions), 4.9 and 5.2 h (photomixotrophic conditions with 1 mM glycerol), and 4.6 and 4.8 h (photomixotrophic conditions with 30 mM glycerol), respectively. These data show that glycerol exerts an inhibitory effect on both strains at low concentrations and little inhibition on PR-6G2 at high concentrations, while for PR-6 high concentrations of glycerol stimulate the growth rate. Interestingly, strain PR-6 but not the glycerol-tolerant strain PR-6G2 can use glycerol for photomixotrophic growth. However, strain PR-6G2 grows under photoheterotrophic conditions (10 μ M DCMU, 1–30 mM glycerol) and faster in the presence of high glycerol concentration. The generation times of PR-6 and PR-6G2 are 19.6 and 17.8 h at 1 mM glycerol and 17.8 and 12.2 h at 30 mM glycerol, respectively.⁵⁴ Rippka et al.²³ have identified some other *Synechococcus* sp. strains capable of facultative photoheterotrophy. *Synechococcus* sp. PCC 7511 can grow on both glucose and sucrose. The strains *Synechococcus* sp. PCC 7003 (*Coccolithis elabens* 17a,⁵⁵) and PCC 73109 (*Agmenellum quadruplicatum* BG-1,⁵³) grow on glycerol and weakly on glucose, while the latter strain additionally exhibits weak growth on fructose. *Synechococcus* sp. PCC 7335 can only grow on fructose.²³

Kratz and Myers⁵⁶ observed an increased respiration in *Synechococcus* sp. PCC 6301 (named *Anacystis nidulans* in this publication) in the dark in the presence of 100 mM glucose or 100 mM fructose, although this organism and its closely related strain *Synechococcus* sp. PCC 7942⁵⁷ have been designated as strictly photoautotrophic.²³ When Zhang et al.³⁹ introduced the *Synechocystis* sp. PCC 6803 *gtr* gene coding for glucose permease²⁶ (Figure 1) on an autonomously replicating plasmid,




Figure 2. Growth (A) and glucose uptake (B) of *Synechococcus* PCC7942 WT and R2-1 cells incubated under the indicated trophic conditions. All cultures of R2-1 contained streptomycin. A: WT, photoautotrophy (\blacktriangle); R2-1, photoautotrophy (\square), transferred from photoautotrophy to photoheterotrophy (\blacksquare , lower line), maintained under photoheterotrophy (\blacksquare , upper line). B: WT (\blacktriangle); R2-1 pregrown under photoautotrophy (\square), adapted to photoheterotrophy (\blacksquare). WT *Synechocystis* PCC6803 grown under photoautotrophy (+). The growth curves shown represent one typical set of 3–4 repeats. Reprinted with permission from ref 39. Copyright 1998 Oxford University Press.




Figure 3. Installation of the glucose transporter to *S. elongatus*. (A) Schematic representation of the glucose transporter gene integration into the *S. elongatus* genome. (B) Growth curve of the *galP* strain (red) and wild type (blue) with and without 5 g/L glucose. (C) Growth curve of the *Glut1* (purple) and *glcP* (green) strains and the wild type. (D) Growth curve of the *galP* strain and the wild type with and without *glcC* deletion (*glcC* KO). For panels B, C, and D, empty symbols are samples in BG-11 medium without glucose, while solid symbols indicate results with BG-11 containing 5 g/L glucose. White and shaded areas indicate the light and dark cycles, respectively. All y axes denote OD₇₃₀, although the scales differ for visibility. Error bars represent standard deviations (in triplicate). NSI, neutral site 1. Reprinted with permission from ref 60. Copyright 2013 American Society for Microbiology.

the transgenic strain *Synechococcus* sp. PCC 7942 *gtr*⁺ becomes capable of photoheterotrophic growth in the presence of 56 mM

glucose (Figure 2), but the plasmid cannot be maintained stably. In contrast, the integration of *gtr* into the chromosome results in




Figure 4. Installation of the sucrose degradation pathway. (A) Schematic representation of integration of the sucrose degradation pathway into the *S. elongatus* chromosome. (B) Synthetic sucrose degradation pathway in *S. elongatus*. Red arrows indicate steps catalyzed by heterologous enzymes. PPP, pentose phosphate pathway. (C) Growth curves of the *cscB*–*cscK* strain (red) and the wild type (blue). Empty and filled symbols indicate growth without and with 5 g/L sucrose, respectively. White and shaded areas indicate light and dark cycles, respectively. Error bars represent standard deviations (in triplicate). Reprinted with permission from ref 60. Copyright 2013 American Society for Microbiology.

a strain that can no longer grow on 56 mM glucose. Niederholtmeyer et al.⁵⁸ introduced the *glf* gene from *Zymomonas mobilis*,⁵⁹ which encodes a glucose- and fructose-facilitated diffusion transporter into *Synechococcus* sp. PCC 7942, and the new transgenic strain can grow in the dark on both 500 μ M glucose and 500 μ M fructose.

McEwen et al.⁶⁰ investigated the ability of *Synechococcus* sp. PCC 7942 to grow on different substrates and find that glucose, sucrose, and xylose slightly increase growth under 12 h light/12 h dark cycles. In order to increase growth on the sugars mentioned above, various genes from other organisms that encode sugar transporters or enzymes involved in sugar metabolism are introduced. All experiments by McEwen et

al.⁶⁰ were performed in BG11 medium.²³ While acquisition of *Glut1* from human erythrocytes⁶¹ and *gtr* from *Synechocystis* sp. PCC 6803²⁶ reduce diurnal growth on 28 mM glucose compared to the wild type (Figure 3C), introduction of *galP* from *E. coli*,⁶² which encodes the glucose/galactose permease, leads to enormously increased growth at the same glucose concentration (Figure 3B). Transfer of *cscB* and *cscK* (Figure 1) from *E. coli*, which encode a sucrose transporter and a fructokinase (Figure 4B), significantly enhances growth on 15 mM sucrose under 12 h light/12 h dark conditions (Figure 4C). The introduction of only *xyIE* (Figure 1) from *E. coli*, which encodes a xylose transporter, does not only increase diurnal growth on 28 mM xylose but also abolishes any growth-




Figure 5. Installation of the xylose degradation pathway. (A) Schematic representation of integration of the xylose degradation pathway into the *S. elongatus* chromosome. (B) Synthetic xylose degradation pathway in *S. elongatus*. Red arrows indicate steps catalyzed by heterologous enzymes. (C) Growth curves of the *xyIE* and *xyIEAB* strains and wild type. Empty and filled symbols indicate growth without and with 5 g/L xylose, respectively. White and shaded areas indicate light and dark cycles. Error bars represent standard deviations (in triplicate). Reprinted with permission from ref 60. Copyright 2013 American Society for Microbiology.

stimulating effect of this sugar (Figure 5C). However, the cotransfer of *xyIA* and *xyIB* (Figure 1) from *E. coli*, which code for a xylose isomerase and a xylulokinase (Figure 5B), together with *xyIE* on an operon, leads to a greatly increased growth on 28 mM xylose under 12 h light/12 h dark conditions compared to

the wild type (Figure 5C). It is assumed that the inability of xylose to enhance the growth of transgenic PCC 7942 *xyIE*⁺ is caused by an accumulation of this sugar inside the cell due to its slow metabolism when genes such as *xyIA* and *xyIB*, which are important for xylose metabolism, are absent.⁶⁰ In contrast to

Synechocystis sp. PCC 6803,⁶³ the acquisition of *xylAB* alone without *xylE* in *Synechococcus* sp. PCC 7942 has not yet been tested. While Zhang et al.³⁹ reported a toxic effect of glucose when *gtr* was integrated into the chromosome, McEwen et al.⁶⁰ observed rapid growth (factor 1.5) within the first 12 h but then a decrease until 72 h with subsequent stagnation. Despite this, the cell density is still higher than that of cultures of the same strain in the absence of glucose. However, the effect of glucose on *Synechococcus* sp. PCC 7942 (wild type and all transgenic mutant strains created by Zhang et al.³⁹) was not tested for more than 96 h by McEwen et al.⁶⁰ Besides Zhang et al.³⁹ used the 2-fold glucose concentration for their experiments compared to McEwen et al.⁶⁰ (56 vs 28 mM).

2.1.5. *Synechocystis*. Rippka et al.²³ identified several *Synechocystis* sp. strains capable of photoheterotrophic growth dependent on glucose. *Synechocystis* sp. PCC 6702 (*Aphanocapsa* HD,⁵⁵), PCC 6805 (*Aphanocapsa* sp.,⁵⁵), PCC 6806 (*Aphanocapsa* sp.,⁵⁵), PCC 6905, and PCC 7201 can only grow on glucose. PCC 7509 also grows on sucrose, and PCC 6906 (also *Eucapsis* sp.) can grow well on glycerol but only weakly on glucose.²³ *Synechocystis* sp. PCC 6714 (*Aphanocapsa* sp.,⁵⁵) can grow both photoheterotrophically^{8,23} and chemoheterotrophically⁶⁴ dependent on glucose in the BG11 medium.²³ The closely related strain, *Synechocystis* sp. PCC 6803 (*Aphanocapsa* sp.,⁵⁵), also grows on 28 mM glucose under photoheterotrophic conditions,^{8,23,31} but contradicting results have been published for its growth in the dark. Rippka⁸ states that no chemoheterotrophic growth on glucose was observed. However, four later publications report the chemoheterotrophic growth of this strain under different glucose concentrations: Astier et al.,³¹ 111 mM, generation time: 19 h; Labarre et al.,⁶⁵ 56 mM, generation time: 22 h; Joset et al.,⁶⁶ 5 mM, generation time: 25 h; Jeanjean et al.,⁶⁷ 56 mM, generation time: 24 h. Under photomixotrophic conditions, 10 mM glucose is slightly toxic for *Synechocystis* sp. PCC 6803.⁶⁸ However, Williams⁶⁹ isolated a glucose-tolerant variant of this strain, which has since become the dominant variant for research, particularly when glucose is added to the growth medium.

Anderson and McIntosh²⁹ discovered that a glucose-tolerant strain of *Synechocystis* sp. PCC 6803⁶⁹ grows on glucose in the dark, when cells receive 5 min of daily illumination (white light, 40 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Since glucose is necessary and the growth rate depends on its concentration, this growth mode is named light-activated heterotrophic growth (LAHG). The different results obtained in the past on chemoheterotrophic growth are probably due to the fact that some scientists have inadvertently interrupted the darkness when measuring cell density, while others have maintained strict darkness. Anderson and McIntosh²⁹ tested different media for their experiments, including BG11²³ with 5 mM TES pH = 8.0 supplemented with 5 mM glucose and BG11 supplemented with 56 mM glucose and 6 mM *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES) at pH 7.5. The latter medium has been reported to be useful for photoheterotrophic growth.⁶⁵ LAHG is improved with the latter medium, and even slow growth occurs within the first 6 days under these conditions; however, a decline is noted thereafter. LAHG has also been observed in a mutant strain lacking all three *psbA* genes and thus a functional photosystem II.^{29,32} Later, Pils et al.⁵⁰ discover that a functional *coxBAC* locus encoding an *aa3* type cytochrome *c* oxidase (homologous to the cytochrome *c* oxidase in mitochondria) is necessary for the LAHG of *Synechocystis* sp. since a *coxBAC* deletion mutant strain loses the capacity of LAHG.

The *cytM* gene, which encodes a cytochrome *c*-type protein, significantly affects heterotrophic growth, for which no good explanation currently exists. Hiraide et al.⁷¹ investigated the capacity of a *Synechocystis* sp. PCC 6803 *cytM* knock out mutant strain under different growth modes and observed faster growth compared to the wild type on 5 mM glucose in both the light and the dark when illuminated for 15 min per day. Furthermore, *Synechocystis* sp. PCC 6803 *cytM*⁻ can grow in permanent darkness with a long generation time of 33 h. All the experiments were performed in BG11 medium²³ buffered with 20 mM HEPES–KOH pH = 8.2 and supplemented with 5 mM glucose as needed.⁶⁵

Lopo et al.⁷² studied the differential growth and behavior of *Synechocystis* sp. PCC 6803 in terms of temperature, pH, trophic mode, glucose, and nitrate concentrations. In these experiments, the presence (photomixotrophy) or absence (photoautotrophy) of glucose causes a significant difference in growth behavior. The growth of autotrophic cultures is largely influenced by pH and temperature, and higher growth is observed at higher pH. On the other hand, photomixotrophic cultures were fairly stable to these changes, although contrary to photomixotrophic cultures, they even grow slightly faster at lower pH.

Both *Synechocystis* sp. strains PCC 6714 and PCC 6803 are highly sensitive to fructose,^{66,68,73} and *Synechocystis* sp. PCC 6803 is killed even at a concentration of 10 mM fructose.⁶⁸ The mechanism of fructose toxicity is not well understood; however, both glucose and fructose enter the cell through the same mechanisms because mutating the gene *gtr*, which encodes the glucose permease²⁶ (Figure 1), abolishes the toxicity of fructose.⁶⁸ Since *Synechocystis* sp. PCC 6803 *gtr*⁻ can no longer use glucose for photoheterotrophic growth⁶⁸ this mutant strain is considered strictly photoautotrophic. Stebegg et al.⁷⁴ discovered that enormously high concentrations (50–200 mM) of fructose not only are harmless but can also support both photomixotrophic (Figure 6A) and photoheterotrophic growth (Figure 6B) of *Synechocystis* sp. PCC 6803 *gtr*⁻, while glucose, applied at the same concentration, does not.

Recently Rapp et al.⁷⁵ investigated the effect of 7-deoxysedoheptulose (7dSh) on several cyanobacteria. 7dSh is excreted by *Synechococcus* sp. PCC 7942 (and by non-cyanobacterial strains like *Streptomyces setonensis*) to inhibit the 3-dehydroquinate synthase (DHQS) of other organisms (including cyanobacteria), thereby preventing the growth of organisms of the same ecological niche. 7dSh is imported by *Synechocystis* sp. PCC 6803 via the Gtr permease since a spontaneous mutant strain resistant to 7dSh (at a concentration of 250 μM) contains a single-point mutation in the *gtr* gene. This strain has also lost the ability to grow on glucose as well as sensitivity to fructose. An artificially generated mutant strain whose *gtr* gene is replaced by a spectinomycin resistance cassette shows the same behavior toward 7dSh. In contrast to fructose, the toxicity of 7dSh is not alleviated by delivery of a 20-fold glucose concentration (5 mM).⁷⁵

Lee et al.⁶³ introduced the *xylA* and *xylB* genes encoding a xylose isomerase and a xylulokinase (Figure 1), respectively, on an operon from *E. coli* into *Synechocystis* sp. PCC 6803, resulting in a strain capable of growth on xylose. When *xylAB* is transferred into an ethylene-producing mutant strain of *Synechocystis* sp. PCC 6803, addition of xylose further increases the production of ethylene.⁶³ Finally, the introduction of *xylAB* into a strain whose *glgC* gene responsible for glycogen synthesis has been deleted^{76,77} results in higher keto acid production.⁶³ In the same year, Xiong et al.⁷⁸ report that this *glgC*⁻/*xylAB*⁺




Figure 6. Photomixotrophic (A) and photoheterotrophic (B) growths of *Synechocystis* sp. PCC 6803 *gtr*[−] dependent on (blue diamond) 0 mM, (red square) 50 mM, (green triangle) 100 mM, and (×) 200 mM fructose. Reprinted with permission from the authors of ref 74.

mutant strain⁶³ metabolizes xylose to acetate in the absence of bound nitrogen. Deletion of the *slr0453* gene, thought to encode a phosphoketolase, decreases acetate production in the light and eliminates it in the dark, demonstrating the importance of the phosphoketolase signaling pathway for heterotrophy in cyanobacteria.⁷⁸

2.1.6. *Thermosynechococcus elongatus*. Zilliges and Dau⁷⁹ investigated different mono- and disaccharides for their potential to induce photomixotrophic or photoheterotrophic growth of *Thermosynechococcus elongatus*. 50 mM glucose and 50 mM fructose and to a lesser extent 50 mM galactose, 35 mM lactose, and 35 mM sucrose act as substrates, while 75 mM arabinose, 35 mM maltose, and 35 mM trehalose do not significantly affect growth. 75 mM xylose even shows an inhibitory effect. Although 50 mM fructose is beneficial initially, it causes cell bleaching after 4 days. 50 mM fructose also supports chemoheterotrophic growth in total darkness or LAHG (at a daily illumination of 15 min).

2.2. Section II: Pleurocapsales. Pleurocapsales are defined as unicellular cyanobacteria that divide by multiple fission.²³ *Chroococidiopsis* is listed within the filamentous cyanobacteria due to phylogenetic reasons.⁴⁰

2.2.1. *Dermocarpa*. *Dermocarpa* sp. PCC 7301 (also *Dermocarpa violacea* in CCAP) can grow photoheterotrophically on glucose, fructose, and sucrose, while *Dermocarpa* sp. strains PCC 7304 and PCC 7437 (also *Chroococidiopsis cyanoasphaera* strain 1965/25⁸⁰) can only use glucose and sucrose for heterotrophy, and *Dermocarpa* sp. PCC 7438 (also *Chroococidiopsis cyanoasphaera* strain 1965/26⁸⁰) grows on glucose and fructose but not on sucrose.²³

2.2.2. *Dermocarpella*. *Dermocarpella* sp. PCC 7326 can grow heterotrophically on glucose, fructose, and sucrose.²³

2.2.3. *Myxosarcina*. *Myxosarcina* sp. strain PCC 7312 grows photoheterotrophically on glucose, while strain PCC 7325 is able to grow on glucose and sucrose. *Myxosarcina chroococcoides* CCAP 1451/1 (named *Chroococidiopsis* sp. PCC 7203 in ref 23) grows photoheterotrophically on glucose, fructose, and sucrose.²³

2.2.4. *Xenococcus*. *Xenococcus* sp. PCC 7307 grows photoheterotrophically on glucose and sucrose, while *Xenococcus* sp. strains 7305 and 7306 do not.²³

2.2.5. *Pleurocapsa* Group. This group consists of several strains within section II that cannot be assigned to any of the five genera mentioned. PCC 7310, PCC 7314, PCC 7319, PCC 7320, PCC 7320, PCC 7321, PCC 7322, PCC 7327 (also *Pleurocapsa minor* OH-69-pm⁸⁰), and PCC 7516 (also *Hyella caespitosa*⁸⁰) can grow photoheterotrophically on sucrose, while PCC 7314, PCC 7319, PCC 7320, and PCC 7322 grow photoheterotrophically on fructose. PCC 7317, PCC 7320, PCC 7322, and PCC 7506 grow photoheterotrophically on glucose.²³

2.3. Section III: Oscillatoriales. Oscillatoriales are defined as filamentous cyanobacteria that do not form heterocysts in the absence of bound nitrogen.²³

2.3.1. *Arthospira platensis*. Phycocyanin is produced in bioreactors from batch cultures of *Arthospira platensis*. Zhang et al.⁸¹ show that the addition of 11 mM glucose to a batch culture of *Arthospira platensis* (called *Spirulina platensis* by Zhang et al.⁸¹) in a growth medium according to Zarouk⁸² increases cell density and produced phycocyanin by a factor of 4.29 and 3.05, respectively.

2.3.2. *Lyngbya*. *Lyngbya lagerheimii* strain Montauk grows heterotrophically in the presence of 55 mM glucose under dim light (40 $\mu\text{W cm}^{-2}$), which does not allow autotrophic growth. In addition very little growth is observed on glucose in the dark. In contrast, at high light intensities that allow for autotrophic growth, glucose does not exert a growth-stimulating effect. Consequently, when *Lyngbya lagerheimii* is cultured under low light, the glucose taken up from the external medium contributes more to the cell's own amino acids compared with high light conditions. While sucrose also supports slight heterotrophic growth in low light, formate, acetate, pyruvate, and succinate do not.³⁰ Rippka et al.²³ tested the photoheterotrophic growth capacity (with DCMU) of *Lyngbya lagerheimii* and identified glucose, fructose, and sucrose as possible substrates (the strain is named PCC 7104 and placed into the LPP group by Rippka et al.²³), while *Lyngbya* sp. PCC 7419 (placed into the LPP group by Rippka et al.²³) grew only weakly on glucose.²³

2.3.3. LPP strains. This group was created by Rippka et al.²³ and includes the genera *Lyngbya*, *Phormidium*, and *Plectonema*. Many strains within the LPP group can use more than one particular organic compound as a carbon source.²³ In this paragraph, the strains that have not been assigned to any genus in 1979 are listed. Those strains capable of heterotrophy, and classified by Rippka et al.²³ as LPP strains but previously known by a different generic name, are discussed under their former name in this review. PCC 7410 and PCC 7505 grow photoheterotrophically on glucose, fructose, ribose, and sucrose. PCC 7123 grows on glucose, fructose, and sucrose. PCC 7114 grows on glucose and sucrose, while PCC 7113 grows photoheterotrophically only on sucrose.²³

2.3.4. *Microcoleus*. *Microcoleus vaginatus* (named LPP PCC 7407 in ref 23) grows on glucose, fructose, and ribose.²³

2.3.5. *Oscillatoria*. Rippka et al.²³ analyzed several *Oscillatoria* strains for their facultative photoheterotrophy. PCC 6602

and PCC 7515 grow on glucose and fructose. PCC 6412 (also *Lyngbya* sp. UTEX 1546) also grows on glucose and fructose but only weakly, while PCC 6407 and PCC 6506 (also *Lyngbya kuetzingii* UTEX 1547) grow weakly on glucose.²³ *Oscillatoria williamsii* MEV (LPP strain PCC 7105 in ref 23) grows on glycerol and weakly on glucose, fructose, and sucrose.²³ *Oscillatoria okeni* TISTR 8549 produces poly(3-hydroxybutyrate-co-3-hydroxyvalerate) (PHBV) in the absence of bound nitrogen. Addition of acetate to the growth medium increases PHBV production by *Oscillatoria okeni* TISTR 8549 under chemoheterotrophic conditions in the dark but not in the presence of light under photomixotrophic conditions.⁸³

2.3.6. Phormidium. *Phormidium* sp. UTEX 1540⁸⁴ grows photoheterotrophically on glucose and fructose.²³

2.3.7. Plectonema. White and Shilo⁸⁵ discover that *Plectonema boryanum* (UTEX 594⁸⁶) can use a variety of substrates for chemoheterotrophic growth in the dark, including ribose, sucrose, mannitol, maltose, glucose, fructose, and casamino acids. Out of all substances tested (each at a concentration of 10 mM), ribose supports the fastest growth in the dark (5 days doubling time) and also enables photoheterotrophic growth in the presence of moderate light and DCMU. The dark growth generation times on the other substances are as follows: sucrose or mannitol (6 days), maltose (10 days), glucose (12 days), and fructose (13 days). In contrast to the other substances, casamino acids alone do not cause dark growth, but in combination with ribose the doubling time is shortened by 50% from 5 days to 2.5 days.⁸⁵ The photoheterotrophic growth of *Plectonema boryanum* (referred to in ref 23 as LPP strain PCC 73110) on glucose, fructose, ribose, and sucrose is later confirmed by Rippka et al.²³ *Plectonema notatum* (described by Kenyon et al.,⁸⁷ also named *Plectonema boryanum* UTEX 581⁸⁶ or *P. boryanum* UTEX 1542,⁸⁴ or the LPP strain PCC 6306²³) and *Plectonema* sp. UTEX 1541 (described by Starr,⁸⁴ also named LPP strain PCC 6402²³) behaves in the same way.²³ Raboy et al.⁸⁸ observed that glucose-dependent dark growth of *Plectonema boryanum* starts after an adaptation phase required for glucose incorporation to start. Later Raboy and Padan⁸⁹ discovered that glucose is actively transported into the cells of this strain. Hiraide et al.⁷¹ identified a dark-adapted mutant strain *dg5* of *Leptolyngbya boryana* (Hiraide et al.⁷¹ call the strain *Plectonema boryanum*) that shows faster growth in the dark on 30 mM glucose in the BG11 medium²³ buffered with 20 mM HEPES–KOH at pH = 7.5. The genome analysis of *dg5* reveals a mutation in the *cytM* gene that encodes Cyt c_M .

2.3.8. Schizothrix. *Schizothrix calcicola* MAN⁸⁷ grows photoheterotrophically on glucose, fructose, sucrose, and ribose²³ (classified as LPP strain PCC 7004).

2.4. Heterocyst-Forming Cyanobacteria. In this subchapter we combine the former sections IV (Nostocales forming linear filaments)²³ and V (Stigonematales forming branching filaments),²³ as they form one monophyletic group.⁴⁰ Only Stigonematales, but not Nostocales, are considered to be of monophyletic origin.⁹⁰ For phylogenetic reasons,⁴⁰ *Chroococcidiopsis* is also mentioned in this chapter.

2.4.1. Anabaena. Sahu and Adhikary⁹¹ studied different organic carbon sources for an undefined *Anabaena* sp. strain isolated by Sahu et al.⁹² Fructose, but particularly lactose, enhances the growth of this strain in the light, while mannose, sodium acetate, and xylose inhibit the growth of the strain; however, none of the latter three substances completely inhibit the growth. In the dark, fructose, lactose, acetate, and mannose are stimulating, with acetate being beneficial only within the first

12 days. Lactose is the most effective substrate in both light and dark conditions, whereas xylose has the most negative effect on growth in light and no stimulating effect in the dark. Regarding nitrogen fixation in the light, acetate, mannose, and xylose all reduce the frequency of occurrence of heterocysts within a filament.⁹¹

Anabaena variabilis ATCC 29413 has been known for many years to grow in the dark at low (5 mM) fructose concentrations.⁹³ It was later shown that the same sugar decreases the doubling time from 24 to 8 h, and heterocyst frequency, nitrogenase activity, and dark respiration are increased by fructose.⁹⁴ The latter effect was reported decades ago, when Kratz and Myers⁵⁶ found increased dark respiration with 100 mM fructose, 100 mM sucrose, and 100 mM maltose but an even stronger effect with 100 mM glucose. Schmetterer et al.⁹⁵ reported that fructose-based dark growth is strictly dependent on an *aa₃*-type cytochrome *c* oxidase homologous to that in mitochondria, since the deletion of the *coxBAC* operon abolishes this ability. Ungerer et al.²⁷ discovered the genes responsible for fructose transport: the *frtABC* operon, which encodes the subunits of the transporter (Figure 1) and is under the control of a repressor encoded by the upstream *frtR* gene. Deletion of *frtR* abolishes dark growth on fructose, and fructose at a concentration greater than 1 mM becomes toxic to *frtR⁻* in the light.²⁷ Rapp et al.⁷⁵ demonstrated that the FrtABC fructose transporter also imports the bioactive inhibitory molecule 7dSh into the cell since a spontaneous mutant strain resistant toward 7dSh (25 μ M within the growth medium) lacks the whole *frtRABC* locus. Consequently the same mutant strain cannot use fructose for photomixotrophic or photoheterotrophic growth anymore; however, supplying a 40-fold molar concentration of fructose (1 mM) does not relieve the toxic effect of 7dSh on the wild type.⁷⁵

The closely related strain, *Anabaena* sp. PCC 7120, has been thought to be strictly photoautotrophic for decades²³ and lacks homologues to the *frtRABC* genes in its genome. Therefore, Ungerer et al.²⁷ introduced the fructose uptake operon from *Anabaena* sp. ATCC 29413 into *Anabaena* sp. PCC 7120, thus conferring to PCC 7120 the ability to grow at low fructose concentrations (5 mM) in the dark. Rapp et al.⁷⁵ demonstrated that the growth of a transgenic *Anabaena* sp. PCC 7120 strain containing the *frtRABC* locus is inhibited by 25 μ M 7dSh. In contrast, the wild type is not affected by the same 7dSh concentration, indicating once more that the FrtABC fructose transporter also imports 7dSh into the cells. Stebbeg et al.⁹⁶ later discovered the capacity of *Anabaena* sp. PCC 7120 wild-type strain to grow photomixotrophically, photoheterotrophically, and chemoheterotrophically at enormously high fructose concentrations (50–200 mM), without applying any genetic manipulation. Deletion of the *coxBAC1* locus, which encodes a cytochrome *c* oxidase similar to that found in the mitochondria, results in a strain unable to grow on fructose in the dark.⁹⁶ Since similar results have been observed for *Anabaena variabilis* ATCC 29413⁹⁵ and LAHG of *Synechocystis* sp. PCC 6803 on glucose,⁷⁰ the question is raised whether all chemoheterotrophic growth among cyanobacteria is dependent on a functional *aa₃*-type cytochrome *c* oxidase. Unfortunately, this has not been tested for other strains so far. The transfer of the *gtr* gene from *Synechocystis* sp. PCC 6803 on an autonomous replicating plasmid into PCC 7120 yields a strain sensitive to glucose concentrations of 5 mM or higher. Although the photoheterotrophic growth of PCC 7120 *gtr⁺* is increased at fructose concentrations of 10 mM up to 100 mM compared to the wild

type, the new transgenic strain loses the ability to grow in the dark on fructose, and 200 mM fructose is even toxic in the light.^{96,97} Unlike fructose, glucose can only support photomixotrophic growth.^{96,98,99} The five genes *glsC*, *glsP*, *glsD*, *glsQ*, and *glsR*, encoding putative components of one or more sugar transporters have been identified, and the five single knockout mutants generated result in decreased photomixotrophic, photoheterotrophic, and chemoheterotrophic growth on fructose.¹⁰⁰

2.4.2. *Anabaenopsis*. *Anabaenopsis circularis*¹⁰¹ grows and fixes nitrogen in the dark in the presence of glucose or fructose and to a lesser extent in the presence of sucrose or maltose, while xylose, mannose, ribose, arabinose, galactose, sorbose, rhamnose, and nonsugar compounds do not affect nitrogen fixation and growth in the dark.¹⁰² Rippka et al.²³ observed photoheterotrophic growth of *A. circularis* (CCAP 1402/1; named *Nostoc* sp. PCC 6720 in ref 23) on glucose and fructose but not on sucrose or ribose.

2.4.3. *Calothrix*. Kenyon et al.⁸⁷ discovered the glucose-based growth of *Calothrix parietina* 1018 (also named *Calothrix parietina* UTEX 484 by Starr⁸⁶) in both the dark and light in the presence of DCMU. Rippka et al.²³ could not reproduce growth on glucose but reported photoheterotrophic growth on sucrose (the strain was named *Calothrix* sp. PCC 6303 by Rippka et al.²³). Many other *Calothrix* sp. strains were identified by Rippka et al.²³ as facultative photoheterotrophs. *Calothrix* sp. PCC 7102 (also *Calothrix desertica*⁸⁷) grows on glucose and fructose. *Calothrix* sp. strains PCC 7103 (also *Nodularia spaerocarpa*, Koch,¹⁰³ or UTEX 583, Starr),⁸⁶ PCC 7111, PCC 7204, and PCC 7415 grow on glucose and fructose but also grow on sucrose, and the growth of *Calothrix* sp. strains PCC 7111 and PCC 7204 is low on fructose. *Calothrix* sp. PCC 7426 grows on ribose and sucrose and weakly on glucose and fructose. *Calothrix* sp. PCC 7116 shows growth behavior similar to that of PCC 7426, with the only exception that growth on ribose is also poor. *Calothrix* sp. PCC 7507 only grows on fructose.²³ The strains *Calothrix* sp. PCC 7101 and PCC 7504 are discussed in the *Tolypothrix* chapter.

Calothrix marchica Lemm. intermedia Rao is able to utilize many carbon sources for photomixotrophic, photoheterotrophic, and chemoheterotrophic growth, including glucose, fructose, and sucrose and to a lesser extent galactose, mannitol, and sorbitol. Continuous precultivation in the presence of sucrose increases both heterotrophic growth and nitrogen content of cultures.¹⁰⁴

2.4.4. *Chlorogloea* (*Chlorogloeopsis*). Fay and Fogg¹⁰⁵ observed that *Chlorogloea fritschii*^{106–108} can fix nitrogen not only in the light but also in the dark. Fay¹⁰⁹ tested various organic substances to see whether they can support the growth and nitrogen fixation of *Chlorogloea fritschii* in the dark. No growth is observed in the first month in the dark regardless of the substrate offered; however, after two or three months, the cultures begin to differ. Mannitol and glucose (each at a concentration of 10 mM) allow dark growth only in the presence of nitrate, while glutamine, glycine, maltose, and especially sucrose (concentration of 10 mM for each of them) induce chemoheterotrophic growth in both the presence and the absence of bound nitrogen. Sucrose, which is the most efficient substrate in the presence and absence of bound nitrogen, is tested at various concentrations (0.1–100 mM) for growth as well as nitrogen fixation in the dark, with 10 mM showing the greatest effect on both processes. Nitrate as a nitrogen source causes 4-fold dark growth at 10 mM sucrose compared to

diazotrophic conditions, while addition of ammonia halves the growth. However, glucose supports growth in the dark in the absence of bound nitrogen (but still less efficiently than 10 mM sucrose) when supplied at 10-fold concentration (i.e., 100 mM), and the addition of nitrate and ammonia increases dark growth only slightly. The growth of *Chlorogloea fritschii* on acetate, pyruvate, citrate, α -ketoglutarate, succinate, fumarate, malate, glycolate, arabinose, fructose, glutamate, and aspartate fails.¹⁰⁹ Rippka et al.²³ analyzed the facultative heterotrophy of *Chlorogloea fritschii* CCAP 1411/1 (also called SAUG 1411/1 by Koch),¹⁰³ *Chlorogloeopsis fritschii* by Mitra and Pandey,¹¹⁰ or *Chlorogloeopsis* sp. PCC 6912 by Rippka et al.²³) and *Chlorogloeopsis* sp. PCC 6718. Both strains grow on glucose, fructose, ribose, and especially on sucrose.²³

Carr¹¹¹ discovered that offering *Chlorogloea fritschii* acetate results in an accumulation of polyhydroxyalkanoates (PHA). Monshupanee et al.¹¹² observed that supplying the growth medium with acetate, pyruvate, citrate, glucose, and fructose increases the poly-3-hydroxybutyrate (PHB) production of *Chlorogloea fritschii* TISTR 8527 under photomixotrophic conditions, and PHB production is greater in the absence of nitrate than in its presence. Acetate, which has the strongest effect on PHB production in light, is also effective in the dark under chemoheterotrophic conditions, particularly in the absence of bound nitrogen and/or phosphorus. Pyruvate, citrate, glucose, and fructose are not tested in the dark. The highest PHB production occurs when cells are preincubated under photoautotrophy to increase biomass and then transferred in the dark under chemoheterotrophic conditions in the absence of bound nitrogen and/or phosphorus.¹¹²

2.4.5. *Chroococciopsis*. *Chroococciopsis* sp. strains PCC 6712 (also described as *Chlorogloea* sp.⁵⁵ or *Chlorogloea* sp. CCAP 1411/2), PCC 7203 (also *Myxosarcina chroococcoides* CCAP 1451/1), PCC 7431 (*Chroococciopsis thermalis* strain 1964/48⁸⁰), PCC 7432 (*Chroococciopsis thermalis* strain 1965/21⁸⁰), PCC 7433 (*Chroococciopsis thermalis* strain 1966/27⁸⁰), PCC 7436 (*Chroococciopsis cubana* strain 1965/108⁸⁰), and PCC 7439 (*Chroococciopsis doonensis* strain 1968/64⁸⁰) can grow photoheterotrophically on glucose, fructose, and sucrose. Apart from PCC 6712, the other strains are assumed by Rippka et al.²³ to be independent isolates from the same strain. *Chroococciopsis* sp. PCC 7434 (*C. cubana* strain 1965/19⁸⁰) can only grow on fructose but not on glucose or sucrose. *Chroococciopsis cyanospaera* strains 1965/25 and 1965/26 (named *Dermocarpa* sp. strains PCC 7437 and PCC 7438 in ref 23) grow on glucose and fructose.²³

2.4.6. *Cylindrospermum*. *Cylindrospermum* sp. PCC 7417 exhibits photoheterotrophic growth on fructose and sucrose.²³

2.4.7. *Fischerella*. Rippka et al.²³ tested several *Fischerella* strains for their ability for photoheterotrophic growth. Apart from *Fischerella* sp. PCC 73103, all other strains of the genus *Fischerella* sp. identified by Rippka et al.²³ have before been assigned to the genus *Mastigocladus*. All of these strains are capable of photoheterotrophy on glucose or fructose while differing according to their potential growth on ribose and sucrose. The *Fischerella* sp. strains PCC 7115 (also *Mastigocladus* sp. H₂), PCC 73103 (also *Fischerella muscicola*, Koch¹⁰³ or CCAP 1427/1 or UTEX 1301, Starr⁸⁴), PCC 7414 (also *Mastigocladus laminosus*), and PCC 7520 (also *Mastigocladus laminosus* I-Kris-m) grow on glucose, fructose, and sucrose. Both *Fischerella* sp. strains PCC 7521 (also *Mastigocladus laminosus* Y-16-m) and PCC 7523 (also *Mastigocladus laminosus* OH-CW-m) are able to grow on glucose, fructose, sucrose, and also on

ribose, but PCC 7521 only grows weakly on ribose. *Fischerella* sp. PCC 7522 (also *Mastigocladus laminosus* NZ-86-m) grows on glucose, fructose, and ribose but not on sucrose.²³

2.4.8. *Nodularia*. *Nodularia* sp. PCC 73104 grows photoheterotrophically on glucose and fructose and weakly on sucrose.²⁵

2.4.9. *Nostoc*. Long ago *Nostoc muscorum* was shown to exhibit slow growth and nitrogen fixation for months in the dark when glucose is supplied, and 56 mM glucose or 29 mM sucrose increases growth of *Nostoc muscorum* in the light.¹¹³ Lazaroff and Vishniac¹¹⁴ named this strain *Nostoc muscorum* A (as derived from Allison et al.¹¹³) and tested several organic molecules for dark growth. Only glucose, fructose, and sucrose act as substrates for chemoheterotrophic growth of *Nostoc muscorum* A, whereas maltose, lactose, glycerol, cellobiose, pyruvate, succinate, citrate, lactate, acetate, urea, and extracts from yeast, beef, or malt or from *Nostoc muscorum* cells fail to do so. Lazaroff and Vishniac¹¹⁴ have also checked the interrelationship of glucose and light on the growth of *Nostoc muscorum* A by measuring the dry weight after an incubation of 22 days. 56 mM glucose supports growth in total darkness; however, light intensities in the range of 1–450 ft candles further increase cell proliferation by reaching the highest dry weight at 80 ft candles. At any light intensity tested, the presence of 56 mM glucose leads to a higher cell mass compared to the same light intensity without glucose. While 56 mM glucose or 29 mM sucrose is sufficient for dark growth, higher concentrations of either sugar or addition of the other organic molecules listed above do not further increase the growth. Only an extract from light grown *Nostoc muscorum* A cells even causes a higher dark growth when added to 56 mM glucose or 29 mM sucrose. In continuous darkness, *Nostoc muscorum* changes its morphology and becomes a mass of large undifferentiated cells. Exposure to low light intensities restores the filamentous phenotype and enhances glucose- or sucrose-dependent growth. While glucose even in the light inhibits the formation of motile filaments, no such effect is observed with sucrose or fructose.¹¹⁴

Kratz and Myers⁵⁶ detected increased respiration in the dark when glucose, fructose, sucrose, or succinate were added. Rippka et al.²³ reported photoheterotrophic growth of *Nostoc* sp. PCC 6314 (described by Kenyon et al.;⁸⁷ also named *Nostoc muscorum* UTEX 1545 by Starr⁸⁴) on sucrose but not on glucose; however, growth on sucrose is sometimes absent and probably attributed to mutations. Vaishampayan¹¹⁵ evaluated all 20 proteinogenic amino acids plus citrulline for their potential to serve as a nitrogen and/or carbon source in a heterocystous but non-nitrogen-fixing mutant strain (*het*⁺/*nif*⁻₁₁) of *Nostoc muscorum*, cultured in the modified Chu 10 medium¹¹⁶ in all experiments. The modified Chu 10 medium contains 40 mg/L Ca(NO₃)₂, 10 mg/L K₂HPO₄, 25 mg/L MgSO₄·7H₂O, 20 mg/L Na₂CO₃, 25 mg/L Na₂SiO₃, 3 mg/L ferric citrate, and 3 mg/mL citric acid. Gerloff et al.¹¹⁶ modified the original Chu 10 medium¹¹⁷ by replacing ferric chloride with ferric citrate and citric acid and kept the K₂HPO₄ concentration at 10 mg/L. DCMU inhibits heterocyst formation in the absence of a suitable carbon source.^{118,119} In a positive control experiment, 3 mM glucose supports photoheterotrophic growth in the presence of DCMU.¹¹⁵ Amino acids that allow growth in a medium free of bound nitrogen are potential sources of nitrogen, while amino acids that allow the formation of heterocysts in the presence of DCMU are potential sources of carbon. Glutamine, histidine, asparagine, tryptophan, and serine are used only as a nitrogen source. Arginine, proline, and phenylalanine serve exclusively as

carbon sources, while leucine, isoleucine, lysine, methionine, valine, and citrulline serve as both carbon and nitrogen sources, as growth occurs in the presence of DCMU and in the absence of bound nitrogen. Aspartic acid, threonine, and glycine have no effect in this experiment, and glutamic acid, alanine, tyrosine, and cysteine are toxic even in the presence of ammonium and in the absence of DCMU.¹¹⁵

Hoare et al.¹²⁰ discovered that an axenic culture of *Nostoc* sp. strain MAC, which is naturally a symbiont on the coralloid roots of the cycad *Macrozamia lucida*, can use glucose, fructose, and sucrose for chemoheterotrophic growth over a pH range from 6 to 9. The growth is further enhanced by the addition of casamino acids. Harder¹²¹ reported that the endophyte can grow in the dark depending on glucose, galactose, sucrose, maltose, starch, insulin, and citric acid. Rippka et al.²³ confirmed photoheterotrophic growth of *Nostoc punctiforme* (also called *Nostoc* sp. ATCC 29133 by Rippka et al.²³) on glucose, fructose, ribose and weakly on sucrose. Summers et al.¹²² discovered both photoheterotrophic and chemoheterotrophic growth in the presence of 50 mM fructose and concomitantly added casamino acids and identified that the *zwf* gene encoding glucose-6-phosphate dehydrogenase is essential for chemoheterotrophy. Later, Ekman et al.²⁸ identified the gene *glcP* and the operon *frtA1A2BC*, which encode the transporters for both sugars and show a high degree of similarity to the corresponding genes in *Synechocystis* sp. PCC 6803 and *Anabaena* sp. ATCC 29413. While wild-type growth occurs at 5 mM fructose and to a lesser extent at 5 mM glucose (in BG11 medium where nitrate is substituted by 2.5 mM ammonium chloride), replacement of most of the *frtA1A2BC* operon with C.K3,¹²³ which encodes a neomycin phosphate transferase not only abolishes fructose-based chemoheterotrophic growth but also changes growth on glucose depending on the direction of insertion of C.K3 in *frtA1A2BC*. The strain CSME1A (C.K3 inserted in the opposite direction of the operon) hardly grows on glucose anymore, while the strain CSME1B (C.K3 inserted in the same direction of the operon) grows better on glucose than the wild type. In strain CSME11 both *glcP* and adjacent *oprB*, which encodes a putative carbohydrate porine, are deleted. CSME11 does not grow on glucose and growth dependent on fructose is drastically reduced.²⁸ Apart from the strains described above, Rippka et al.²³ observed photoheterotrophy in several *Nostoc* sp. strains. PCC 7107, PCC 7416, and PCC 7423 grow on fructose, while *Nostoc* sp. strains PCC 6705 and PCC 7524 grow on sucrose. The growth of strain *Nostoc* sp. PCC 6705 on glucose is variable due to mutations accumulated through culture over multiple cell cycles. Both *Nostoc* sp. strains PCC 6302 (described by Kenyon et al.,⁸⁷ also named *Anabaena* sp. UTEX 1551 by Starr)⁸⁴ and PCC 6310 (described by Kenyon et al.,⁸⁷ also named *Anabaena spiroides* UTEX 1552 by Starr⁸⁴) grow on glucose, fructose, ribose, and sucrose, but growth of the latter strain on ribose is low.²³ Later, Schmetterer and Flores¹²⁴ also reported growth of *Nostoc* sp. PCC 7107 (named *Nostoc* sp. ATCC 29150 in this work) on fructose in the dark while checking the fructose uptake of this strain. Several fructose concentrations (1–333 mM) are tested. The fastest growth is observed at 33 mM, while higher concentrations are less supportive.¹²⁴

2.4.10. *Scytonema*. Rippka et al.²³ reported photoheterotrophic growth of *Scytonema* sp. PCC 7110 on glucose, fructose, and sucrose. Similar results to those obtained for *Calothrix marchica* Lemm. Var. *intermedia* were discovered for *Scytonema schmidlei* de Toni indicating photomixotrophic, photoheterotrophic, and chemoheterotrophic growth on glucose, fructose,

and sucrose and to a lesser extent on galactose, mannitol, and sorbitol.¹⁰⁴

2.4.11. *Tolypothrix*. *Tolypothrix tenuis* can grow chemoheterotrophically on glucose in the dark only when ammonia is the nitrogen source and not nitrate as in most media. Ammonia and glucose enable dark growth most efficiently at a pH of 6.1. Casamino acids¹²⁵ even support dark growth in the absence of glucose for the first 2 weeks before the culture reaches the stationary phase. Hence casamino acids can be used as both a carbon and nitrogen source; however, coaddition of glucose as a more suitable carbon source increases growth rate further. Casamino acids are most beneficial at 30 mM, while higher concentrations are less supportive of *Tolypothrix tenuis* growth. In the case of glucose, on the other hand, 50 mM has the same stimulating effect as the higher concentrations. Different amino acids are tested as nitrogen sources on their own (in combination with glucose as a carbon source) for dark growth, and arginine and phenylalanine achieve about half of the growth rate compared to casamino acids when added at 28 mM, while other amino acids are less supportive. Fructose exerts an effect similar to that of glucose, while sucrose hardly elicits dark growth. Heterotrophy causes an increase of pigments like chlorophyll a, carotenoids, and phycocyanin within a week.¹²⁶ The growth rate on glucose is further enhanced when cultures are illuminated at low light intensities (less than 500 lx), insufficient for photoautotrophic growth. Although glucose consumption from the medium is even slightly higher in the dark than in the presence of regular light (6000 lx), much more glucose is incorporated into glycogen in the light than under dark conditions.¹²⁷ Rippka et al.²³ reported photoheterotrophic growth of *Tolypothrix tenuis* (referred to as *Calothrix* sp. PCC 7101 in Rippka et al.²³) on glucose, fructose, and ribose. The same phenotype was also detected for PCC 7505 by Rippka et al.,²³ who placed this strain in the genus *Calothrix* and assumed it just to be another isolate of PCC 7101.

2.4.12. *Westiellopsis*. *Westiellopsis prolifica* can fix nitrogen in light and in the dark. Fructose, lactose, sucrose, sorbose, galactose, glucose, sodium acetate, mannitol, sorbitol, and glycerol increase growth both in the dark and in the light. When incubated in the dark, the initial stimulating effect on growth is higher for all exogenous substrates mentioned, but these substances are utilized more quickly in the light. Mannose, xylose, acetic acid, propionic acid, fructose 1,6-bisphosphate, pyruvic acid, dihydroxy acetone, and succinic acid exert photogrowth inhibitory effects.¹²⁸

2.5. Prochlorophyta. In addition to the classic cyanobacteria (which, because of their phycocyanins, resemble the chloroplasts of red algae), there is a special group that (like green algae and embryophytes) has chlorophyll *b* instead of phycocyanins. Therefore, they were previously thought to be the actual ancestors of chloroplasts in green algae and land plants.¹²⁹ However, it is now assumed that classic cyanobacteria, prochlorophyta, and chloroplast of red algae, green algae, and embryophytes have developed differently from a common ancestor. This organism originally possessed both chlorophyll *b* and phycocyanins and later lost either one or both of them during evolution.^{130,131} Although these organisms do not form a monophyletic group, they are summarized as prochlorophyta.^{132,133} The genus *Prochlorococcus* is widespread in the oceans, where it (along with *Synechococcus*) is responsible for most of the marine photosynthesis.^{50–52} For a review of *Prochlorococcus* photomixotrophy see Munoz-Marin et al.⁴⁸ *Prochlorococcus* is reported to take up glucose, which increases the expression of

genes involved in glucose metabolism (e.g., *zwf*, *gnd*, *lld*).¹³⁴ It was later shown that the gene Pro1404 (although annotated as the putative melibiose/sodium symporter *melB*) encodes a glucose transporter since insertion of Pro1404 into the *asnS* gene of *Synechococcus* sp. PCC 7942 results in a strain that imports glucose into the cell. Nevertheless, the uptake rate of the new transgenic is still below the glucose uptake rate of *Synechocystis* sp. PCC 6803.¹³⁵ Finally, a positive correlation between Pro1404 expression and glucose levels is shown.¹³⁶ Duhamel et al.⁴⁹ studied the photomixotrophic growth of undefined *Prochlorococcus* sp. strains and observed a behavior similar to that of *Synechococcus* sp. The tested *Prochlorococcus* sp. strains also assimilated glucose, leucine, ATP and especially molecules containing nitrogen and phosphorus at a higher rate in the light compared to dark conditions or when photosystem II was inhibited.

3. CONCLUSION AND FUTURE RESEARCH

From the very beginning, cyanobacteria have mainly been presented as the inventors of plant oxygenic photosynthesis, which, however, accounts for only part of their complex metabolism. While even facultatively heterotrophic cyanobacteria show the highest growth rates under photoautotrophic conditions, the various forms of heterotrophy in this kingdom are more widespread than previously thought. In recent years, more and more strains have been either intentionally identified^{23,64,74,93,120,122} or identified by chance⁹⁶ as potential heterotrophs.

The vast majority of cyanobacteria strains is still considered to be strictly photoautotrophic²³ with the caveat that most of these strains have not been tested on all possible substrates. Therefore, we need to examine other potential substrates (fatty acids and amino acids) that have been rather neglected in the past. For example, Rippka et al.²³ tested only glucose, fructose, ribose, sucrose, and in some cases also glycerol for the ability to undergo heterotrophy. Although it is unlikely that strains that cannot grow photoheterotrophically have the potential for dark chemoheterotrophic growth, it is possible that some of these strains may use the substrates tested by Rippka et al.²³ or other compounds as growth-promoting substrates for photomixotrophy. There may also be strains classified as strictly photolithoautotrophic as no suitable conditions for heterotrophy have been identified so far. In some cases, classic substrates such as fructose can stimulate growth; however, exceptionally high concentrations not existing in nature must be offered as shown in *Anabaena* sp. PCC 7120⁹⁶ and *Synechocystis* sp. PCC 6803 *gtr*[−],⁷⁴ because the substrate would not enter the cell at naturally occurring external concentrations. Since no specific transporter has been identified so far, we have to assume that a transporter for a related substance (another sugar?) will import that molecule if its external concentration is high enough. Regarding the fact that photosynthesis and respiration are intimately linked in cyanobacteria,^{137,138} the question arises whether the facultative heterotrophy of some cyanobacteria strains is a relic of heterotrophic predators before the invention of oxygenic photosynthesis and was lost in other strains or whether the facultative heterotrophy was formed *de novo* as a secondary acquisition in some strains. To answer this question, knowledge about facultative (particularly photo) heterotrophy in anoxygenic phototrophs such as purple and green bacteria needs to be increased. Recently, Matheus Carnevali et al.¹³⁹ analyzed the genome sequences of melainobacteria and sericytochromatia, which are considered to be the organisms

most closely related to cyanobacteria, although they themselves do not carry out photosynthesis.¹⁴⁰ According to Matheus Carnevali et al.¹³⁹ the ancestors of cyanobacteria, melainabacteria, and seritochromatia are probably anaerobes living on fermentation and possessing various hydrogenases. Cyanobacteria therefore became aerobic after splitting off from melainabacteria and seritochromatia. Although the capacity for growth on external organic substances may have originated independently in various strains, we think that increasing the knowledge of heterotrophy among cyanobacteria will give new insights into evolutionary processes.

Cultivation under heterotrophic conditions without oxygenic photosynthesis can also promote research on critical cell parts (e.g., parts of photosystems), where otherwise erasing the gene information would be lethal. Heterotrophic cultivation can also allow growth when light must be avoided or is unavailable for long periods (e.g., transportation in space). Our picture of cyanobacteria needs to change from pure photoautotrophs to multitrophs capable of adapting to appropriate metabolic modes, depending on the current environmental conditions.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomega.3c02205>.

Table S1 that lists all cyanobacterial strains known by date to be capable of heterotrophic growth (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Ronald Stebegg – Universität Wien, Fakultät für Chemie, Institut für Biophysikalische Chemie, 1090 Wien, Austria; Email: ronald.stebegg@univie.ac.at

Annette Rompel – Universität Wien, Fakultät für Chemie, Institut für Biophysikalische Chemie, 1090 Wien, Austria; orcid.org/0000-0002-5919-0553; Email: annette.rompel@univie.ac.at

Author

Georg Schmetterer – Universität Wien, Fakultät für Chemie, Institut für Biophysikalische Chemie, 1090 Wien, Austria

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsomega.3c02205>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Nadiia Gumerova, PhD, for her great support designing Figure 1, the graphical abstract, and the supplementary cover. This work was supported by the project “Verbesserung des Verfahrens zur Herstellung sowie der Methoden der Qualitätskontrolle und Reinheitsprüfung von hochgereinigtem Chlorophyll a” of the Universität Wien [grant no. ET524001].

■ REFERENCES

(1) Schmetterer, G. Cyanobacterial respiration. In *The molecular biology of cyanobacteria*; Bryant, D.A., Ed.; Kluwer Academic Publishers, 1994; pp 409–435. DOI: 10.1007/978-94-011-0227-8_13.

(2) Schmetterer, G.; Pils, D. Cyanobacterial respiration. In *Respiration in archaea and bacteria*; Zannoni, D., Ed.; Diversity of prokaryotic respiratory systems, 2004; Vol. 16, pp 261–278.

(3) Golbeck, J. H. Photosystem I in cyanobacteria. In *The molecular biology of cyanobacteria*; Bryant, D. A., Ed.; Kluwer Academic Publishers, 1994; pp 319–360. DOI: 10.1007/978-94-011-0227-8_10.

(4) Barry, B. A.; Boerner, R. J.; de Paula, J. C. The use of cyanobacteria in the study of the structure and function of photosystem II. In *The molecular biology of cyanobacteria*; Bryant, D. A., Ed.; Kluwer Academic Publishers, 1994; pp 217–257. DOI: 10.1007/978-94-011-0227-8_8.

(5) Mereschkowski, C. Über Natur und Ursprung der Chromatophoren im Pflanzenreiche. *Biol. Centralbl.* 1905, 25, 593–604.

(6) Stanier, R. Y.; Cohen-Bazire, G. Phototrophic prokaryotes: the cyanobacteria. *Annu. Rev. Microbiol.* 1977, 31, 225–274.

(7) Khoja, T.; Whitton, B. A. Heterotrophic growth of blue-green algae. *Arch. Mikrobiol.* 1971, 79, 280–282.

(8) Rippka, R. Photoheterotrophy and chemoheterotrophy among unicellular blue-green algae. *Arch. Mikrobiol.* 1972, 87, 93–98.

(9) Smith, A. J. Modes of cyanobacterial carbon metabolism. In *The biology of cyanobacteria*; Carr, N. G., Whitton, B. A., Eds.; University of California Press: Berkeley, 1982; pp 47–86.

(10) Faust, M. A.; Gantt, E. Effect of light intensity and glycerol on the growth and pigment composition and ultrastructure of *Chroomonas* sp. *J. Phycol.* 1973, 9, 489–495.

(11) Lylis, J. C.; Trainor, F. R. The heterotrophic capabilities of *Cyclotella meneghiniana*. *J. Phycol.* 1973, 9, 365–369.

(12) Ellis, R. J. Heterotrophic nutrition and its effect on chlorophyll synthesis in *Golenkinia* (Chlorophyceae). *J. Phycol.* 1977, 13, 304–306.

(13) Danforth, W. F. Substrate assimilation and heterotrophy. In *Physiology and Biochemistry of Algae*; Lewin, R. A., Ed.; Academic Press: New York, 1962; pp 99–123.

(14) Hoare, D. S.; Hoare, S. L.; Smith, A. J. Heterotrophic potentialities of blue-green algae. In *Taxonomy and Biology of Blue-green Algae*; Desikachary, T. V., Ed.; University of Madras, India, 1970; pp 501–507.

(15) Pelroy, R. A.; Bassham, J. A. Photosynthetic and dark carbon metabolism in unicellular blue-green algae. *Archiv. Mikrobiol.* 1972, 86, 25–38.

(16) Fogg, G. E.; Stewart, W. D. P.; Fay, P.; Walsby, A. E. Heterotrophy and respiration. In *The Blue-green Algae*; Academic Press: New York and London, 1973; pp 155–180.

(17) Stanier, R. Y. Autotrophy and heterotrophy in unicellular blue-green algae. In *The Biology of blue-green Algae*; Carr, N. G., Whitton, B. A., Eds.; Alden Press: Oxford, 1973; pp 501–518.

(18) Smith, A. J. Synthesis of metabolic intermediates. In *The Biology of Blue-green algae*; Carr, N. G., Whitton, B. A., Eds.; Alden Press: Oxford, 1973; pp 1–38.

(19) Wolk, C. P. Physiology and cytological chemistry of blue-green algae. *Bacteriol. Rev.* 1973, 37, 32–101.

(20) Pope, D. H. Heterotrophic potential of *Phormidium* and other blue-green algae. *Canad. J. Bot.* 1974, 52, 2369–2374.

(21) Khoja, T.; Whitton, B. A. Heterotrophic growth of filamentous blue-green algae. *Br. Phycol. J.* 1975, 10, 139–148.

(22) Fay, P. Factors influencing dark nitrogen fixation in a blue-green alga. *Appl. Environ. Microbiol.* 1976, 31, 376–379.

(23) Rippka, R.; Derulles, J.; Waterbury, J. B.; Herdman, M.; Stanier, R. Y. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* 1979, 111, 1–61.

(24) Cyanobase: <https://genome.microbedb.jp/cyanobase/> (accessed June 15, 2023).

(25) Kowata, H.; Tochigi, S.; Takahashi, H.; Kojima, S. Outer Membrane Permeability of Cyanobacterium *Synechocystis* sp. strain PCC 6803: Studies of Passive Diffusion of Small Organic Nutrients Reveal the Absence of Classical Porins and Intrinsically Low Permeability. *J. Bacteriol.* 2017, 199, e00371–17.

(26) Schmetterer, G. R. Sequence conservation among the glucose transporter from the cyanobacterium *Synechocystis* sp. PCC 6803 and mammalian glucose transporters. *Plant. Mol. Biol.* 1990, 14, 697–706.

(27) Ungerer, J. L.; Pratte, B. S.; Thiel, T. Regulation of Fructose Transport and Its Effect on Fructose Toxicity in *Anabaena* spp. *J. Bacteriol.* 2008, 190, 8115–8125.

- (28) Ekman, M.; Picossi, S.; Campbell, E. L.; Meeks, J. C.; Flores, E. A. *Nostoc punctiforme* Sugar transporter Necessary to Establish a Cyanobacterium-Plant Symbiosis. *Plant Physiol.* **2013**, *161*, 1984–1992.
- (29) Anderson, S. L.; McIntosh, L. Light-activated heterotrophic growth of the cyanobacterium *Synechocystis* sp. strain PCC 6803: a blue-light-requiring process. *J. Bacteriol.* **1991**, *173*, 2761–2767.
- (30) Van Baalen, C.; Hoare, D. S.; Brandt, E. Heterotrophic growth of blue-green algae in dim light. *J. Bacteriol.* **1971**, *105*, 685–689.
- (31) Astier, C.; Elmorjani, K.; Meyer, I.; Joset, F.; Herdman, M. Photosynthetic mutants of the cyanobacteria *Synechocystis* sp. strains PCC 6714 and PCC 6803: sodium p-hydroxymercuribenzoate as selective agent. *J. Bacteriol.* **1984**, *158*, 659–664.
- (32) Jansson, C.; Debus, R. J.; Osiewicz, H. D.; Gurevitz, M.; McIntosh, L. Construction of an obligate photoheterotrophic mutant of the cyanobacterium *Synechocystis* 6803. *Plant Physiol.* **1987**, *85*, 1021–1025.
- (33) Diner, B. A.; Nixon, P. J.; Farchaus, J. W. Site-directed mutagenesis of photosynthetic reaction centers. *Curr. Opin. Struct. Biol.* **1991**, *1*, 546–554.
- (34) Bishop, N. I. The influence of the herbicide, DCMU, on the oxygen evolving system of photosynthesis. *Biochim. Biophys. Acta* **1958**, *27*, 205–206.
- (35) Hirschberg, J.; McIntosh, L. Molecular Basis of Herbicide Resistance in *Amaranthus hybridus*. *Science* **1983**, *222*, 1346–1349.
- (36) Golden, S. S.; Haselkorn, R. Mutation to herbicide resistance map within the *psbA* gene of *Anacystis nidulans* R2. *Science* **1985**, *229*, 1104–1107.
- (37) Metz, J. G.; Pakrasi, H. B.; Seibert, M.; Arntzer, C. J. Evidence for a dual function of the herbicide-binding D1 protein in photosystem II. *FEBS Lett.* **1986**, *205*, 269–274.
- (38) Stebbeg, R.; Schmetterer, G.; Rompel, A. Transport of organic substances through the cytoplasmic membrane of cyanobacteria. *Phytochem.* **2019**, *157*, 206–218.
- (39) Zhang, C. C.; Jeanjean, R.; Joset, F. Obligate phototrophy in cyanobacteria: more than a lack of sugar transport. *FEMS Microbiol. Lett.* **1998**, *161*, 285–292.
- (40) Fewer, D.; Friedl, T.; Büdel, B. *Chroococcidiopsis* and heterocyst-differentiating cyanobacteria are each other's closest living relatives. *Mol. Phylogenet. Evol.* **2002**, *23*, 82–90.
- (41) Reddy, K. J.; Haskell, J. B.; Sherman, D. M.; Sherman, L. A. Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus *Cyanothece*. *J. Bacteriol.* **1993**, *175*, 1284–1292.
- (42) Colon-Lopez, M. S.; Sherman, D. M.; Sherman, L. A. Transcriptional and translational regulation of nitrogenase in light-dark- and continuous-light-grown cultures of the unicellular cyanobacterium *Cyanothece* sp. strain ATCC 51142. *J. Bacteriol.* **1997**, *179*, 4319–4327.
- (43) Stöckel, J.; Welsh, E. A.; Liberton, M.; Kunnvakkam, R.; Aurora, R.; Pakrasi, H. B. Global transcriptomic analysis of *Cyanothece* 51142 reveals robust diurnal oscillation of central metabolic processes. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 6156–6161.
- (44) Toepel, J.; Welsh, E.; Summerfield, T. C.; Pakrasi, H. B.; Sherman, L. A. Differential transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during light-dark and continuous-light growth. *J. Bacteriol.* **2008**, *190*, 3904–3913.
- (45) Welsh, E. A.; Liberton, M.; Stöckel, J.; Loh, T.; Elvitigala, T.; Wang, C.; Wollam, A.; Fulton, R. S.; Clifton, S. W.; Jacobs, J. M.; Aurora, R.; Ghosh, B. K.; Sherman, L. A.; Smith, R. D.; Wilson, R. K.; Pakrasi, H. B. The genome of *Cyanothece* 51142, a unicellular diazotrophic cyanobacterium important in the marine nitrogen cycle. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 15094–15099.
- (46) Feng, X.; Bandyopadhyay, A.; Berla, B.; Page, L.; Wu, B.; Pakrasi, H. B.; Tang, Y. J. Mixotrophic and photoheterotrophic metabolism in *Cyanothece* sp. ATCC 51142 under continuous light. *Microbiol.* **2010**, *156*, 2566–2574.
- (47) Mares, J.; Johansen, J. R.; Hauer, T.; Zima, J., Jr; Ventura, S.; Cuzman, O.; Tiribilli, B.; Kastovsky, J. Taxonomic resolution of the genus *Cyanothece* (Chroococcales, cyanobacteria), with a treatment on *Gloeothece* and three new genera *Crocospaera*, *Rippkaea* and *Zehria*. *J. Phycol.* **2019**, *55*, 578–610.
- (48) Munoz-Marin, M. C.; Gomez-Baena, G.; Lopez-Lozano, A.; Moreno-Cabezuelo, J. A.; Diez, J.; Garcia-Fernandez, J. M. Mixotrophy in marine picocyanobacteria: use of organic compounds by *Prochlorococcus* and *Synechococcus*. *ISME J.* **2020**, *14*, 1065–1073.
- (49) Duhamel, S.; Van Wambeke, F.; Lefevre, D.; Benavides, M.; Bonnet, S. Mixotrophic metabolism by natural communities of unicellular cyanobacteria in the western tropical South Pacific Ocean. *Environ. Microbiol.* **2018**, *20*, 2743–2756.
- (50) Partensky, F.; Blanchot, J.; Vault, D. Differential distribution and ecology of *Prochlorococcus* and *Synechococcus* in oceanic waters: a review. In *Marine cyanobacteria*; Charpy, L., Larkum, A., Eds.; Musee Oceanographique: Monaco, 1999; pp 457–475.
- (51) Agawin, N. S. R.; Duarte, C. M.; Agusti, S. Nutrient and temperature control of the contribution of picoplankton biomass and production. *Limnol. Oceanogr.* **2000**, *45*, 591–600.
- (52) Garcia-Pichel, F.; Johnson, S. L.; Youngkin, D.; Belnap, J. Small scale vertical distribution of bacterial biomass and diversity in biological soil crusts from arid lands in the Colorado Plateau. *Microb. Ecol.* **2003**, *46*, 312–321.
- (53) Van Baalen, C. Studies on marine blue-green algae. *Bot. Mar.* **1962**, *4*, 129–139.
- (54) Lambert, D. H.; Stevens, S. E., Jr. Photoheterotrophic growth of *Agmenellum quadruplicatum* PR-6. *J. Bacteriol.* **1986**, *165*, 654–656.
- (55) Stanier, R. Y.; Kunisawa, R.; Mandel, M.; Cohen-Bazire, G. Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriol. Rev.* **1971**, *35*, 171–205.
- (56) Kratz, W. A.; Myers, J. Photosynthesis and respiration of three blue-green algae. *Pl. Physiol.* **1955**, *30*, 275–280.
- (57) Sugita, C.; Ogata, K.; Shikata, M.; Jikuya, H.; Takano, J.; Furumichi, M.; Kanehisa, M.; Omata, T.; Sugiura, M.; Sugita, M. Complete nucleotide sequence of the freshwater unicellular cyanobacterium *Synechococcus elongatus* PCC 6301 chromosome: gene content and organization. *Photosynth. Res.* **2007**, *93*, 55–67.
- (58) Niederholtmeyer, H.; Wolfstädter, B. T.; Savage, D. F.; Silver, P. A.; Way, J. C. Engineering Cyanobacteria To Synthesize and Export Hydrophilic Products. *Appl. Environ. Microbiol.* **2010**, *76*, 3462–3466.
- (59) Barnell, W. O. K.; Yi, K. C.; Conway, T. Sequence and genetic organization of a *Zymomonas mobilis* gene cluster that encodes several enzymes of glucose metabolism. *J. Bacteriol.* **1990**, *172*, 7227–7240.
- (60) McEwen, J. T.; Machado, I. M. P.; Connor, M. R.; Atsumi, S. Engineering *Synechococcus elongatus* PCC 7942 for Continuous Growth under Diurnal Conditions. *Appl. Environ. Microbiol.* **2013**, *79*, 1668–1675.
- (61) Mueckler, M.; Caruso, C.; Baldwin, S. A.; Panico, M.; Blench, I.; Morris, H. R.; Allard, W. J.; Lienhard, G. E.; Lodish, H. F. Sequence and structure of a human glucose transporter. *Science* **1985**, *229*, 941–945.
- (62) Hernandez-Montalvo, V.; Martinez, A.; Hernandez-Chavez, G.; Bolivar, F.; Valle, F.; Gosset, G. Expression of *galP* and *glk* in a *Escherichia coli* PTS mutant restores glucose transport and increases glycolytic flux to fermentation products. *Biotechnol. Bioeng.* **2003**, *83*, 687–694.
- (63) Lee, T. C.; Xiong, W.; Paddock, T.; Carrieri, D.; Chang, I. F.; Chiu, H. F.; Ungerer, J.; Hank Juo, S. H.; Maness, P. C.; Yu, J. Engineered xylose utilization enhances bio-products productivity in the cyanobacterium *Synechocystis* sp. PCC 6803. *Metab. Eng.* **2015**, *30*, 179–189.
- (64) Vernotte, C.; Picaud, M.; Kirilovsky, D.; Olive, J.; Ajlani, G.; Astier, C. Changes in the photosynthetic apparatus in the cyanobacterium *Synechocystis* sp. PCC 6714 following light-to-dark and dark-to-light transitions. *Photosynth. Res.* **1992**, *32*, 45–57.
- (65) Labarre, J.; Thuriaux, P.; Chauvat, F. Genetic analysis of amino acid transport in the facultatively heterotrophic cyanobacterium *Synechocystis* sp. strain 6803. *J. Bacteriol.* **1987**, *169*, 4668–4673.
- (66) Joset, F.; Buchou, T.; Zhang, C.-C.; Jeanjean, R. Physiological and genetic analysis of the glucose-fructose permeation system in two *Synechocystis* sp. species. *Arch. Microbiol.* **1988**, *149*, 417–421.

- (67) Jeanjean, R.; Onana, B.; Peschek, G. A.; Joset, F. Mutants of the cyanobacterium *Synechocystis* PCC 6803 impaired in respiration and unable to tolerate high salt concentrations. *FEMS Microbiol. Lett.* **1990**, *68*, 125–130.
- (68) Flores, E.; Schmetterer, G. Interaction of fructose with the glucose permease of the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **1986**, *166*, 693–696.
- (69) Williams, J. G. K. Construction of specific mutations in photosystem II photosynthetic reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol.* **1988**, *167*, 766–778.
- (70) Pils, D.; Gregor, W.; Schmetterer, G. Evidence for in vivo activity of three distinct respiratory terminal oxidases in the cyanobacterium *Synechocystis* sp. strain PCC6803. *FEMS Microbiol. Lett.* **1997**, *152*, 83–88.
- (71) Hiraide, Y.; Oshima, K.; Fujisawa, T.; Uesaka, K.; Hirose, Y.; Tsujimoto, R.; Yamamoto, H.; Okamoto, S.; Nakamura, Y.; Terauchi, K.; Omata, T.; Ihara, K.; Hattori, M.; Fujita, Y. Loss of cytochrome *cM* stimulates cyanobacterial heterotrophic growth in the dark. *Plant Cell Physiol.* **2015**, *56*, 334–345.
- (72) Lopo, M.; Montagud, A.; Navarro, E.; Cunha, I.; Zille, A.; de Cordoba, P. F.; Moradas-Ferreira, P.; Tamagnini, P.; Urchueguia, J. F. Experimental and modeling analysis of *Synechocystis* sp. PCC 6803 growth. *J. Mol. Microbiol. Biotechnol.* **2012**, *22*, 71–82.
- (73) Zhang, C. C.; Durand, M. C.; Jeanjean, R.; Joset, F. Molecular and genetical analysis of the fructose and glucose transport system in the cyanobacterium *Synechocystis* PCC6803. *Mol. Microbiol.* **1989**, *3*, 1221–1229.
- (74) Stebeegg, R.; Schmetterer, G.; Rompel, A. Photoheterotrophic growth of unicellular cyanobacterium *Synechocystis* sp. PCC 6803 *gtr*[−] dependent on fructose. *Monatsh. Chem.* **2019**, *150*, 1863–1868.
- (75) Rapp, J.; Wagner, B.; Brilisaue, K.; Forchhammer, K. *In vivo* inhibition of the 3-dehydroquinate synthase by 7-deoxysedoheptulose depends on promiscuous uptake by sugar transporters in cyanobacteria. *Front. Microbiol.* **2021**, *12*, No. 692986.
- (76) Carrieri, D.; Paddock, T.; Maness, P. C.; Seibert, M.; Yu, J. P. Photo-catalytic conversion of carbon dioxide to organic acids by a recombinant cyanobacterium incapable of glycogen storage. *Energy Environ. Sci.* **2012**, *5*, 9457–9461.
- (77) Grundel, M.; Scheunemann, R.; Lockau, W.; Zilliges, Y. Impaired glycogen synthesis causes metabolic overflow reactions and affects stress responses in the cyanobacterium *Synechocystis* sp. PCC 6803. *Microbiology* **2012**, *158*, 3032–3043.
- (78) Xiong, W.; Lee, T. C.; Rommelfanger, S.; Gjersing, E.; Cano, M.; Maness, P. C.; Ghirardi, M.; Yu, J. Phosphoketolase pathway contributes to carbon metabolism in cyanobacteria. *Nat. Plants* **2016**, *2*, 15187.
- (79) Zilliges, Y.; Dau, H. Unexpected capacity for organic carbon assimilation by *Thermosynechococcus elongatus*, a crucial photosynthetic organism. *FEBS Lett.* **2016**, *590*, 962–970.
- (80) Waterbury, J. B.; Stanier, R. Y. Patterns of growth and development in pleurocapsalean cyanobacteria. *Microbiol. Rev.* **1978**, *42*, 2–44.
- (81) Zhang, X. W.; Zhang, Y. M.; Chen, F. Kinetic models for phycocyanin production by high cell density mixotrophic culture of the microalga *Spirulina platensis*. *J. Ind. Microbiol. Biotechnol.* **1998**, *21*, 283–288.
- (82) Zarouk, C. Contribution a l'etude d'une cyanophycee de divers facteurs physique et chimiques sur la croissance et photosynthese de *Spirulina maxima*. Geitler, *PhD thesis*; University of Paris: France, 1966.
- (83) Taepucharoen, K.; Tarawat, S.; Puangcharoen, M.; Incharoensakdi, A.; Monshupanee, T. Production of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) under photoautotrophy and heterotrophy by non-heterocystous N₂-fixing cyanobacterium. *Bioresour. Technol.* **2017**, *239*, 523–527.
- (84) Starr, R. C. *Culture collection of algae*; Additions to the collection July 1, 1963–July 1, 1966. Privately printed. Department of Botany, Indiana University: Bloomington, Indiana, 1966.
- (85) White, A. W.; Shilo, M. Heterotrophic growth of the filamentous blue green alga *Plectonema boryanum*. *Arch. Microbiol.* **1975**, *102*, 123–127.
- (86) Starr, R. C. The culture collection of algae at Indiana University. Division Cyanophyta. *Am. J. Bot.* **1964**, *51*, 1013–1044.
- (87) Kenyon, C. N.; Rippka, R.; Stanier, R. Y. Fatty acid composition and physiological properties of some filamentous blue-green algae. *Arch. Mikrobiol.* **1972**, *83*, 216–236.
- (88) Raboy, B.; Padan, E.; Shilo, M. Heterotrophic capacities of *Plectonema boryanum*. *Arch. Microbiol.* **1976**, *110*, 77–85.
- (89) Raboy, B.; Padan, E. Active transport of glucose and α -methylglucoside in the cyanobacterium *Plectonema boryanum*. *J. Biol. Chem.* **1978**, *253*, 3287–3291.
- (90) Mishra, A. K.; Shukla, E. S.; Singh, S. S. Phylogenetic comparison among the heterocystous cyanobacteria based on a polyphasic approach. *Protoplasma* **2013**, *250*, 77–94.
- (91) Sahu, J.; Adhikary, S. P. Heterotrophic growth and nitrogen fixation in the filamentous blue-green alga *Anabaena* sp. *Z. Allg. Mikrobiol.* **1981**, *21*, 669–676.
- (92) Sahu, J. K.; Adhikary, S. P.; Pattnaik, H. Growth response of *Anabaena* sp., a blue-green alga to organic substrates in light and dark. *I. J. Expt. Biol.* **1979**, *17*, 1401–1402.
- (93) Wolk, C. P.; Shaffer, P. W. Heterotrophic micro- and macrocultures of a nitrogen-fixing cyanobacterium. *Arch. Microbiol.* **1976**, *110*, 145–147.
- (94) Haury, J. F.; Spiller, H. Fructose uptake and influence on growth of and nitrogen fixation by *Anabaena variabilis*. *J. Bacteriol.* **1981**, *147*, 227–235.
- (95) Schmetterer, G.; Valladares, A.; Pils, D.; Steinbach, S.; Pacher, M.; Muro-Pastor, A. M.; Flores, E.; Herrero, A. The *coxBAC* operon encodes a cytochrome *c* oxidase required for heterotrophic growth in the cyanobacterium *Anabaena variabilis* strain ATCC 29413. *J. Bacteriol.* **2001**, *183*, 6429–6434.
- (96) Stebeegg, R.; Wurzing, B.; Mikulic, M.; Schmetterer, G. Chemoheterotrophic growth of the cyanobacterium *Anabaena* sp. strain PCC 7120 dependent on a functional cytochrome *c* oxidase. *J. Bacteriol.* **2012**, *194*, 4601–4607.
- (97) Stebeegg, R. Heterotrophic growth of the cyanobacterium *Anabaena* (*Nostoc*) sp. strain PCC 7120 and its dependence on a functional *coxI* locus encoding cytochrome *c* oxidase. *Doctoral thesis*; University of Vienna: Austria, 2011.
- (98) Yu, G.; Shi, D.; Cai, Z.; Cong, W.; Ouyang, F. Growth and physiological features of cyanobacterium *Anabaena* sp. strain PCC 7120 in a glucose-mixotrophic culture. *Chin. J. Chem. Eng.* **2011**, *19*, 108–115.
- (99) Malatinszky, D.; Steuer, R.; Jones, P. R. A comprehensively curated genome-scale two-cell model for the heterocystous cyanobacterium *Anabaena* sp. PCC 7120. *Plant Physiol.* **2017**, *173*, 509–523.
- (100) Nieves-Morion, M.; Flores, E. Multiple ABC glucoside transporters mediate sugar-stimulated growth in the heterocyst-forming cyanobacterium *Anabaena* sp. strain PCC 7120. *Environ. Microbiol. Rep.* **2018**, *10*, 40–48.
- (101) Watanabe, A. Distribution of nitrogen-fixing blue green-algae in various areas of South and East Asia. *J. Gen. Appl. Microbiol.* **1959**, *5*, 21–29.
- (102) Watanabe, A.; Yamamoto, Y. Heterotrophic nitrogen fixation by the blue-green alga *Anabaenopsis circularis*. *Nature* **1967**, *214*, 738.
- (103) Koch, W. Verzeichnis der Sammlung von Algenkulturen am Pflanzenphysiologischen Institut der Universität Göttingen. *Arch. Mikrobiol.* **1964**, *47*, 402–432.
- (104) Adhikary, S. P.; Sahu, J. K. Utilization of organic substrates by two filamentous cyanobacteria under various growth conditions. *Acta Microbiol. Hung.* **1988**, *35*, 101–106.
- (105) Fay, P.; Fogg, G. E. Studies on nitrogen fixation by blue-green algae. III. Growth and nitrogen fixation in *Chlorogloea fritschii* Mitra. *Arch. Mikrobiol.* **1962**, *42*, 310–312.
- (106) Mitra, A. K. Two new algae from Indian soils. *Ann. Bot.* **1950**, *14*, 457–464.

- (107) Fay, P.; Kumar, H. D.; Fogg, G. E. Cellular factors affecting nitrogen fixation in the blue-green alga *Chlorogloea fritschii*. *J. Gen. Microbiol.* **1964**, *35*, 351–360.
- (108) Evans, E. H.; Foulds, I.; Carr, N. G. Environmental conditions and morphological variation of the blue-green alga *Chlorogloea fritschii*. *J. Gen. Microbiol.* **1976**, *92*, 147–155.
- (109) Fay, P. Heterotrophy and nitrogen fixation in *Chlorogloea fritschii*. *J. Gen. Microbiol.* **1965**, *39*, 11–20.
- (110) Mitra, A. K.; Pandey, D. C. On a new genus of the blue-green alga *Chlorogloeopsis* with remarks on the heterocysts in the alga. *Phykos* **1966**, *5*, 106–114.
- (111) Carr, N. G. The occurrence of poly- β -hydroxybutyrate in the blue-green alga *Chlorogloea fritschii*. *Biochem. Biophys. Acta* **1966**, *120*, 308–310.
- (112) Monshupanee, T.; Nimdach, P.; Incharoensakdi, A. Two-stage (photoautotrophy and heterotrophy) cultivation enables efficient production of Bioplastic poly-3-hydroxybutyrate in auto-sedimenting cyanobacterium. *Sci. Rep.* **2016**, *6*, 37121.
- (113) Allison, F. E.; Hoover, S. R.; Morris, H. J. Physiological studies with the nitrogen-fixing alga *Nostoc muscorum*. *Botanical Gazette* **1937**, *98*, 433–463.
- (114) Lazaroff, N.; Vishniac, W. The effect of light on the developmental cycle of *Nostoc muscorum*, a filamentous blue-green alga. *J. Gen. Microbiol.* **1961**, *25*, 365–374.
- (115) Vaishampayan, A. Amino acid nutrition in the blue-green alga *Nostoc muscorum*. *New Phytol.* **1982**, *90*, 545–549.
- (116) Gerloff, G. C.; Fitzgerald, G. P.; Skoog, F. The isolation, purification and culture of blue-green algae. *Am. J. Bot.* **1950**, *37*, 216–218.
- (117) Chu, S. P. The influence of the mineral composition of the medium on the growth of planktonic algae. I. Methods and culture media. *J. Ecol.* **1942**, *30*, 284–325.
- (118) Vaishampayan, A.; Singh, H. R.; Singh, H. N. Biological effects of rice-field herbicide stam *f-34* on various strains of the nitrogen-fixing blue-green alga *Nostoc muscorum*. *Biochemie und Physiologie der Pflanzen* **1978**, *173*, 410–419.
- (119) Singh, H. N.; Vaishampayan, A. Biological effects of the rice-field herbicide machete on various strains of the nitrogen-fixing blue-green alga *Nostoc muscorum*. *Env. Exp. Bot.* **1978**, *18*, 87–94.
- (120) Hoare, D. S.; Ingram, L. O.; Thurston, E. L.; Walkup, R. Dark heterotrophic growth of an endophytic blue-green alga. *Arch. Mikrobiol.* **1971**, *78*, 310–321.
- (121) Harder, R. Ernährungsphysiologische Untersuchungen an Cyanophyceen, hauptsächlich dem endophytischen *Nostoc punctiforme*. *Z. Bot.* **1917**, *9*, 149–242.
- (122) Summers, M. L.; Wallis, J. G.; Campbell, E. L.; Meeks, J. C. Genetic evidence of a major role for glucose-6-phosphate dehydrogenase in nitrogen fixation and dark growth of the cyanobacterium *Nostoc* sp. strain ATCC 29133. *J. Bacteriol.* **1995**, *177*, 6184–6194.
- (123) Elhai, J.; Wolk, C. P. A versatile class of positive-selection vectors based on the non viability of palindrome-containing plasmids that allows cloning into long polylinkers. *Gene* **1988**, *68*, 119–138.
- (124) Schmetterer, G.; Flores, E. Uptake of fructose by the cyanobacterium *Nostoc* sp. ATCC 29150. *Biochim. Biophys. Acta* **1988**, *942*, 33–37.
- (125) Mueller, J. H.; Johnson, E. R. Acid hydrolysates of casein to replace peptone in the preparation of bacteriological media. *J. Immunol.* **1941**, *40*, 33–38.
- (126) Kiyohara, T.; Fujita, Y.; Hattori, A.; Watanabe, A. Heterotrophic culture of a blue-green alga, *Tolypothrix tenuis* I. *J. Gen. Appl. Microbiol.* **1960**, *6*, 176–182.
- (127) Kiyohara, T.; Fujita, Y.; Hattori, A.; Watanabe, A. Effect of light on glucose assimilation in *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.* **1962**, *8*, 165–168.
- (128) Adhikary, S. P.; Pattnaik, H. Growth response of *Westiellopsis prolifica* Janet to organic substrates in light and dark. *Hydrobiol.* **1979**, *67*, 241–247.
- (129) Lewin, R. A. Prochloron and the theory of symbiogenesis. *Ann. N.Y. Acad. Sci.* **1981**, *361*, 325–329.
- (130) Pinevich, A. V.; Velichko, N.; Ivanikova, N. Cyanobacteria of the genus prochlorothrix. *Front. Microbiol.* **2012**, *3*, 173.
- (131) Pinevich, A. V. Chloroplast history clarified by the criterion of light-harvesting complex. *Biosystems* **2020**, *196*, No. 104173.
- (132) Palenik, B.; Haselkorn, R. Multiple evolutionary origins of prochlorophytes, the chlorophyll *b*-containing prokaryotes. *Nature* **1992**, *355*, 265–267.
- (133) Urbach, E.; Robertson, D. L.; Chisholm, S. W. Multiple evolutionary origins of prochlorophytes within the cyanobacterial radiation. *Nature* **1992**, *355*, 267–270.
- (134) Gomez-Baena, G.; Lopez-Lozano, A.; Gil-Martinez, J.; Lucena, J. M.; Diez, J.; Candau, P.; Garcia-Fernandez, J. M. Glucose uptake and its effect on gene expression in *Prochlorococcus*. *PLoS One* **2008**, *3*, No. e3416.
- (135) Munoz-Marin, M. C.; Luque, I.; Zubkov, M. V.; Hill, P. G.; Diez, J.; Garcia-Fernandez, J. M. *Prochlorococcus* can use the Pro1404 transporter to take up glucose at nanomolar concentrations in the atlantic ocean. *Proc. Natl. Acad. Sci. U.S.A.* **2013**, *110*, 8597–8602.
- (136) Munoz-Marin, M. d. C.; Gomez-Baena, G.; Diez, J.; Beynon, R. J.; Gonzalez-Ballester, D.; Zubkov, M. V.; Garcia-Fernandez, J. M. Glucose uptake in *Prochlorococcus*: Diversity of kinetics and effects on the metabolism. *Front. Microbiol.* **2017**, *8*, 327.
- (137) Scherer, S.; Almon, H.; Böger, P. Interaction of photosynthesis, respiration and nitrogen fixation in cyanobacteria. *Photosynth. Res.* **1988**, *15*, 95–114.
- (138) Scherer, S. Do photosynthetic and respiratory electron transport chains share redox proteins? *Trends Biochem. Sci.* **1990**, *15*, 458–462.
- (139) Matheus Carnevali, P. B.; Schulz, F.; Castelle, C. J.; Kantor, R. S.; Shih, P. M.; Sharon, I.; Santini, J. M.; Olm, M. R.; Amano, Y.; Thomas, B. C.; Anantharaman, K.; Burstein, D.; Becraft, E. D.; Stepanauskas, R.; Woyke, T.; Banfield, J. F. Hydrogen-based metabolism as an ancestral trait in lineages sibling to the cyanobacteria. *Nat. Commun.* **2019**, *10*, 463.
- (140) Di Rienzi, S. C.; Sharon, I.; Wrighton, K. C.; Koren, O.; Hug, L. A.; Thomas, B. C.; Goodrich, J. K.; Bell, J. T.; Spector, T. D.; Banfield, J. F.; Ley, R. E. The human gut and groundwater harbor non-photosynthetic bacteria belonging to a new candidate phylum sibling to cyanobacteria. *Elife*. **2013**, *2*, No. e01102.