

BOTANISCHES INSTITUT
der Universität Wien


Bibliothek

J.-Nr.

3449

Sign.

318/11



BOTANISCHES MUSEUM
der k. k. Universität.

J.N.º 3479

B

87/10

TABELLEN EN PLATEN

VOOR DE

JAVAANSCH E ORCHIDEËN,

DOOR

C. L. B L U M E.



BATAVIA,

TER LANDS DRUKKERIJ,

1825.



TABELLEN IN TAFELN

VOON 18

JAVANESCHES ORCHIDEEN

DOOR

G. A. RECHT

DEEL I

DEEL II

1888

CH. 113 CEMETERIAL OUCHIDE 114-1-1110301

CLAVIS GENERUM ORCHIDEARUM JAVANICARUM.

GRANULOSAE. POLLINIS MASSAE E PARTICULIS NUMEROSIS ANGULATIS V. SQUAMOSIS ELASTICE CONFLATAE. HERBAE TERRESTRES.

DIANDRAE. A. *Anthera loculis discretis.*

<i>Adfinitas.</i>	<i>Sepala.</i>	<i>Labellum.</i>	<i>Gynostemium.</i>	<i>Anthera.</i>	<i>Massæ Pollinis.</i>	<i>Habitus.</i>
HABENARIA. R. Br. . .	ringentia.	basi calcarat.; limb. par-tit. patens.	erectum.	antica, loculis lateralib. basi elongatis.	2, pedicellatæ, pedicellis singulis glandula suffult.	Terrest. caulescens. Rad. tubero-sæ. Flor. spicati.
PERISTYLUS. [Orchis.] .	præcedentis.	præcedentis.	erectum, ad apicem utroque latere glandula terminatum.	antica, loculis medio adnat., parallelis.	præcedentis.	præcedentis.
MECOSA. [Orchis.] . . .	præcedentis.	basi calcaratum; limb. in-divis., patens.	dilatatum, emarginat.	antica, loculis lateralib. basi non productis.	præcedentis.	præcedentis.

MONANDRAE. B. *Anthera unica, bilocularis.*

1. Pollinis massæ subpedicellatæ, glandula v. lamellula communi suffultæ.						
ANECOCYLUS. [Epi-pactis.] . . .	dorsali cum inter. forni-cat.; lateral. latior., patent.	inferne ventricos., intrus bicallos.; limb. elongat. di-latat. 2 lobus, patens.	breve, attenuato-emargin., utroque latere callos., an-tice appendic. bifid. instruct.	terminal., postic.	2, sub-2-lobæ, subpedi-cellat. pellicula comm. suffult.	Terrest. Caulis basi repens. Folia alternæ, ovata. Flor. spicat., sessiles.
CHEIROSTYLIS. [Epi-pactis.] . . .	ringent.; exterior. basi connat.	basi angustat.; limb. difa-tat., 2-lobus, patens.	breve, proboscideo-bifid.; laciniis dorso incrassat.	præcedentis.	præcedentis.	præcedentis.
ADENOSTYLIS. [Neottia.]	ringentia.	basi ventricos., gynostem. connat., intrus pubesc.; limb. spatulat., incrassat., patens.	breve, attenuato-emargi-nat., utroque latere glandu-loso-tumid.	præcedentis.	præcedentis.	Terrest. Caulis inferne repens. Fo- lia linearia. Spica tortilis.
ETABRIA. [Calochilus.] .	ringentia.	basi ventricosoproduct. in-trus biglandulos.; limbus for-nicat., incrassat., marginib. glandulos.	breve, attenuato-emargi-nat., utroque later. denie instruct.	præcedentis.	præcedentis.	præcedentis, sed folia ovato-lanceo-lat., nervosa. Flor. spicat., sessiles, non tortuosi.
ERYTHRODES. [Neottia.]	erecto-convivent.; lateral. labellum amplectent.	inferne scrobiculare-calca-rat.; limb. brevis, emargi-nat., patens.	breve, acuminato-emargi-natum, antice nudum.	præcedentis.	2, sub-2-lobæ, longitudi-naliter coalitæ, pedicello com-muni hamato suffult.	præcedentis.
NEOTTIA. Sw. . . .	ringent.; interior. angus-tiora, cum dorsali subformi-cata.	ventricos., intrus nudum aut barbat.; limbus brevis, attenuat., patens, rarius sublobat.	apice emarginatum.	præcedentis.	2, sub-2-lobæ, subpedicel-lat., stylo comm. suffult.	præcedentium.
SECT. 1. Gynostachis. . .	"	intrus eberbe, limb. indivis.	antice inappendiculatum.	"	"	Folia angusta. Spica tortilis.
* SECT. 2. Racodes. . .	"	intrus eberbe; limb. sublob.	" ad basin bilamellat.	"	"	" lata. Spica haud tortilis.
SECT. 3. Eriporchis. . .	"	intrus pubescens.; limb. in-divis.	" inappendiculatum.	"	"	" " "

2. Pollinis massæ pedicellatæ, pedicellis nudis, glandula v. lamellula destitutis.

GALERA. [Arethusa.] . .	convivent.; interior. la-tiora.	basi breviter obtuse-calca-rat., concav.; limb. indivis.	breve, subforficatum.	terminal., antica, carinosa	2, sub-2-lobæ, pedicel-lis arcuatis instruct.	Terrest. Rad. tuberosa. Caulis aphyll. Flor. spicat., pedicellat.
EUCOSIA. [Eriochilus.] . .	convivent.; interior. an-gustiora.	sepalis exter. conforme.	breve, obtusum.	denti dorsal. affixa.	2, 2-lobæ, pedicellis apice connexis.	Terrest. caulesc. Folia ovata, membranac. Spica pauciflora.

3. Pollinis massæ non pedicellatæ, margini stigmatis adglutinata.

CORDYLA.	convivent.; interior. mi-nora.	calcarat., basi concav., limbus sublobat.	clavatum.	terminal., carinosa.	2, sub-2-lobæ.	Terrest. Rad. tuberosa. Folia mem-branac., cordat. Flores spicati.
------------------	--------------------------------	---	-----------	----------------------	----------------	--

PULVEREA. POLLINIS MASSAE IN PULVEREM FACILE SOLVENDAE. HERBAE TERRESTRES, PAUCAE PARASITICAE.

A. *Monantherae. Anthera unica.*

1. Massæ pollinis lobatæ, longitudinaliter dehiscentes, intus pulveræ, glandula v. dente communi suffultæ.						
ZOSTEROSTYLIS. [Cranichis.] .	patentia, linearia.	basi ventricos., gynostem. amplect.; limb. indivis., pa-tens.	breve, dorso cavitate an-therifera repanda instruct.	dorsalis.	2, 2-lobæ, glandulae com-mun. insidentes.	Terrest. Rad. fasciculat. Folia ra-dical. Flor. spicati.
CALCEARIA. [Satyrium.]	ringent., basi coalit.; dor-sale fornicat.; reliq. angus-tiss., patentibus.	inferne scrotiforme, gynostemium amplect.; limb. patens, 2-lobus.	breve.	terminalis.	2, denti gynostemii ad-hærent.	Terrest. Rad. tuberosa. Fol. ra-dicalia, cordat. Flor. solitar.

2. Massæ pollinis pulveræ, non lobatim conflata.

EPIPHANES. [Gastrodia.]	monophyll., antice fissum ore 5-fido.	mediocre, ungui gynostem. incurv. affixum.	apice utroque latere dente instructum.	terminalis.	pulveræ.	Terrest. Radic. tuberos. Scapus aphyll. Flores spicati.
VANILLA. Sw.	patent., subæqual.	gynostem. inferne adnat.; limb. sublobat., patens.	antice ad apicem cavitate antherifer. instruct.	terminal., antica.	pulveræ.	Parasit. vel rupestr. Caulis scan-dentes. Flor. subspicati.
CHLOROSA. [Cranichis.]	erecto-patent.; inter. angustiora	sepalis exteriorib. confor-me, gynostem. amplectens.	breve, crassum.	terminalis.	pulveræ.	Parasit. Rad. articulat. Folia ra-dicalia, membranacea. Flor. spicati.
THELYMITRA. Sw. . . .	erecto-patent., subæqual.	sepalis conforme.	erectum, superne cucul-lato-trifidum.	antice infra apicem cucul-lo inscrt.	pulveræ.	Terrest. Rad. bulbos. Scapus in-ferne foliat. Flor. spicati.

APOSTASIA. [Cyripedium.]

	erecto-patent., æqualia.	sepalis conforme.	erectum, versus apicem process. dorsal. et inferne 2 lateralib. antheriferis instruct.	2, oblongæ, 2-loculares.	pulveræ.	Terrest. Rad. fibros. Caulis erect. Fol. linearilanceolat. Spica termi-nalis.
--	--------------------------	-------------------	--	--------------------------	----------	---

Continuatio CLAVIS GENERUM ORCHIDEARUM JAVANICARUM.

CEREACEAE. POLLINIS MASSAE DETERMINATAE, FARINOSO-PULPOSAE AUT CEREACEAE.

A. GYNOSTEMIUM OVARIO CONTINUUM.

* * * * *

1. *Labellum basi nullo modo productum.*

a. *GYNOSTEMIUM LIBERUM.*

* Gynostemium indivisum.

<i>Adfinitas.</i>	<i>Sepala.</i>	<i>Labellum.</i>	<i>Gynostemium.</i>	<i>Anthera.</i>	<i>Massa pollinis.</i>	<i>Habitus.</i>
MALAXIS. Sw.	erecto-patent. v. reflex; int. angustior.	inferne concav.; limb. patens.	obtusum.	terminal., bilocul.	2, 2-partibiles.	Terrest. v. parasit. Folia membr. v. subcarnea.
Sect. Leptorchis.	erecto-patentia.	"	semiteretes.	"	"	Terrest. Folia membranac., ovato-lanceol.; bulbis suffult. Spica panicul. Flor. purpureascent.
* " Platystylis.	patentia.	"	dilatatum.	"	"	Parasit. Folia membranac. v. subcarnea. Spica paucifl. Flores flavescens. v. purpur.
" Liparis.	reflexa.	"	inferne gibbosum.	"	"	Parasit. Folia subcarnea., lineari-lanceol.; bulbis imposit. Spica multifl. Flor. flavi.
* " Ensifera. [Cymb. Sw.]	patent. v. reflexa.	"	breve, crassum.	"	"	Parasit., ebulb. Folia distich., ensat., subcarnea. Spica multiflor., tortilis. Flor. flavi.
CYRTOSIA. [Vanilla.]	conviventia.	concav., int. tumidum; limb. erect., rotundat. v. sublobat.	apice subdilat. v. subalatum.	terminal., 2-loc., carnos.	2, subarcuatae.	Terrest. Rad. tuberos. Caul. aphyll. Flor. spicat. v. paniculat.
CYMBIDIUM. Sw.	patent., subaeq., linear.	concav.; limb. lobat., patens.	obtusum.	terminal., simi-2-loc.	2, 2-lob., juxtaposit., elastic. tice lamin. comm. semilunat. suffult.	Parasit. Folia bulbis saepe suffult. Flores racemos.
SECT. 1.					angulatae, angulo ext. attenuat.	Folia radical., linear. Peduncul. multiflor.
SECT. 2.					ovales v. subglobose.	Folia bulb. suffult., lanceolat. Peduncul. paucifl.
GRAMMATOPHYLLUM. [Cymb. Sw.]	patent., subaequal., suprae latiora.	concav., gynostemii callo elastic affix.; limb. lobat., patens.	obtusum, basi antice processu calloso instruct.	praecedent.	2, globose, basi foveat., lamin. comm. arcuat. a se invicem disjunct. insident.	Parasit., caulescens. Folia linearia. Flores racemos.
CHELONANTHERA. [Cymb. Sw.]	patent. v. erecto-convivent.; int. angusta, saepe reflexa.	basi concav.; limb. sublobat., patens.	superne dilatata.	inferne infra apicem affixa, 2-loc., 2-valv.	4, basi foveat., elastic. membran. comm. insident.	Parasit. Folia bulb. suffult. Flor. spicati, distichi.
SECT. 1.	"	utroque latere dente autem; limb. emarginat.	supra antheram petaloideo dilatatum.	inferne medio affixa.	"	"
SECT. 2.	"	intus cristat.; limb. semitrilobus.	superne dilatata.	inferne versus apicem gynostemio insidens.	"	"
SECT. 3.	"	praecedentis sed limbus indivisus.	praecedentis.	praecedentis.	"	"
ACANTHOCLOSSUM. [Chelonanth.]	praecedentis.	basi ventricos.; limb. patent., in. bicallos. (unde nomen.)	membranaceo-dilatata.	inferne infra apicem locata, 2-valv., 2-locul.	4, elastic. pedicello comm. hamato suffult.	praecedentis.
ADENONCOS.	erectiuscula, subaequal.	concav., carnos., indivis., int. glanduloso-tumidum.	breve, obtusum.	terminal., 2-locul.	4, subglobos., elastic. pedicello hamato affixe.	Parasit., subcaulesc. Folia lineari-subulat., carnos. Pedunculi paucifl., oppositifolii.
ARUNDINA. [Bletia.]	erecto patent.; subaequalia.	concav. intus cristat.; limb. lobat., patens.	erecta., superne subdilatata.	terminal., antica, 8-locul.	8, compr., elastic. cohaerent.	Terrest., caulesc. Caul. nodosi. Folia lineari-lanceolat. Spica terminal.
SPATHOGLOTTIS.	praecedentis.	basi dilatato-bilob., supra basin callos, limb. spatulat.	subincurv., superne dilatata.	terminal., 2-locul.	8, filis elastic. coherent.	Terrest. Folia membranac., nervos. Scapus radical., multifl.
TUHELASIS.	convivent., subaequal.; ext. basi conat.	sepalis conforme.	breve, attenuato-emarginat.	dorsal., 2-locular.	8, pedicello hamato suffult.	Parasit. Folia plerumq. bulbis suffult., subcarnea. Flores spicati.

** Gynostemium divisum.

DENDROCHILUM. [Oncid. Sw.]	patent., subaequal.	basi concav., integrum; limb. erecto-patens.	apice dilatato-3-fidum.	inferne infra apicem affixa, bivalv.	4, elastic. glandul. comm. suffultae.	Parasit. Folia e bulb. erumpent. Flor. spicat., distichi.
SECT.		parvum, elastic affix.	gynostemii lacin. media attenuata, emarginata.			Bulbi remoti. Spica laterales.
SECT.		sepalis subaequal., concav.	gynostemii lacin. media truncato-emarginata.			Bulbi consociat. Spicae ex apice bulbos. ortae.

b. *Gynostemium cum labello connatum.*

* Gynostemium indivisum.

PLOCOGLOTTIS.	tingent.; ext. basi levit. coalita.	inferne plicis inflexis gynost. junct.; limb. indivis., patens.	superne liberum, obtusum.	inferne ad apicem locata, 2-locul.	4, compress., per paria stylis connexae, ope glandul. comm. affix.	Terrest. Rad. fibros. Folia solitaria., membranac. Scapus radical., multifl.
GASTROGLOTTIS [Oncid. thidium.]	patent.; lateral latior., labello supposit., inferne conat.	basi ventricos., inferne gynost. adnat.; limb. semi-3-fidus.	superne liberum, antice attenuatum.	terminal., postica, 2-locular.	4, per paria filis elastic. junctae	Terrest. Rad. fibros. Caul. simpl. Folia membranac. Spica terminalis.

** Gynostemium divisum.

ACRIOPSIS.	quatuor, uniseriata, patent.	inferne cum gynost. tubulat.; limb. patens, callosus.	apice cucullat., utroque latere processu auctum.	inferne infra apicem, 2-locular.	2, pedicello comm. ad apic. glandulif. suffultae.	Parasit. Folia bulb. insid. Flores paniculati.
CREPIDIUM. [Malaxis. Sw.]	reflexa; int. angustiora.	basi concav.; cum gynost. coalit.; limb. semilunat., gynost. amplexens.	brevisimum, biauriculatum.	terminal., postica, 2-locular.	2, 2-partibiles.	Terrest. Rad. fibros. Folia membranac. Spica terminal., multiflora.

Continuatio CLAVIS GENERUM ORCHIDEARUM JAVANICARUM.

CEREACEAE. POLLINIS MASSAE DETERMINATAE, FARINOSO-PULPOSAE AUT CEREACEAE.

A. GYNOSTEMIUM OVARIO CONTINUUM.

* * * * *

2. *Labellum basi saccatum vel calcaratum.*

A. GYNOSTEMIUM LIBERUM.

* Gynostemium indivisum.

<i>Adfinitas.</i>	<i>Sepala.</i>	<i>Labellum.</i>	<i>Gynostemium.</i>	<i>Anthera.</i>	<i>Massae Pollinis.</i>	<i>Habitus.</i>
AERIDES. Sw.	lineari-oblong., patent.	parvum, saccatum; limb. patens.	semicylindric., obtusum, ad apicem interne emarginat.	terminal., sub-2-locul.	2, lobulo dorsali auct., pedicello comm. ad basin pel-tato. affix.	
ARACHNIS. [Aerides Sw.]	praeceidentis.	subsaccato-concav., 3-lob.; lobo medio carnos.	praeceidentis.	praeceidentis.	2, 2-lobae, lamina triangula-ri ad basin peltat. suffult.	Paras. rarius rupestr. Caulis sub-ramos. Folia distich., linearia v. lan-ceolat., coriacea v. carnos. Spic. multifl., saepe paniculat.
CLEISOSTOMA. [Aerides Sw.]	dorsale fornicat.; reliquis patentibus; lateral. labello supposit.	basi obtuso-calcarat.; cal-car demum clausum; limbus erectus, indivis.	interne ad apicem rostello attenuato instructum.	praeceidentis.	2, lobulo dorsali auct., pedicello comm. hamato in-tinct.	
ECHIOGLOSSUM. [Aerides Sw.]	praeceidentis.	praeceidentis, sed limbus elongatus, difform., callosus, in ligulam bifid. pro-ductus.	semicylindricum., obtu-sum, antice ad apicem emar-ginat.	praeceidentium.	2, lobulo auct., pedicel-lo comm. ad basin arcuato-dilat. suff.	
PACHYSTOMA.	erecta; lateral. label. in-feme amplectent.; int. an-gustiora.	inferne calcar breve ob-tus. formans; limb. erect., semilobat., intus tumidus.	incurvum, clavatum.	terminal., sub-8-locellat.	8, elasticae cohaerentes.	Terrestr. Rad. tuberosa. Caulis aphyll., paleis vaginatus. Flor. ter-minat., nutantes.
LIMATODIS. [Bletia.] . . .	erecto-patent., subaequal.	concav., in calcar breve acut. product.; limb. erecto-patens, indivis.	superne dilat., infra api-cem rostello deflexo instruct.	gynostemium antice ter-minans, 8-locellat.	praeceidentis.	Terrestr. Rad. fibrose. Caulis sim-pl. Folia membranac. Pedunculi la-teral., paucifl.
NEPHELAPHYLLUM. . . .	reflexa, subaeq.	inferne in calcar breve scrobiculare product.; lim-bus erect.	latum.	terminal., 2-locul., suban-gulata.	praeceidentis.	Terrestr., caulescens. Folia pe-tiolata, ovata, nebulosa (unde no-men). Flor. terminal., spicati.
GLOMERA.	subrigit., lateral. infer-ne amplect.	inferne obtuse calcar.; lim-bus indivis., patens, callos.	breve, dilatatum.	terminal., dorsalis, sub-8-locellat.	praeceidentis.	Paras. Caulis simpl. Fol. coria-cea. Flores dense capitat.

** Gynostemium divisum.

SCHOENORCHIS. [Aerides Sw.]	erecto-convivent.; int. mi-nora.	saccat. v. calcarat.; limb. incrassat., erect., v. patens.	antice bicornutum (rostel-lum nempe bifidum).	terminal., 2-locul., ligu-lata.	2, bipartibil., elasticae pe-dicello comm. hamato affix.	
Sect. 1.	"	calcaratum; limbus patens.	mediocre.	"	"	Paras. Caulis simpl. v. subramos. Folia angusta, subcarnos. Spic. sim-pl. v. pauciculata.
" 2. Catastylis.	"	saccatum; limb. conicus, erect.	brevissimum.	"	"	

B. GYNOSTEMIUM CUM LABELLO CONNATUM.

CERATOCHYLUS. . . .	patent., subaeq., superne latiora.	parvum, inferne in sac-cum compress. product.; limb. styliformis, patens.	obtus., apice interne emar-ginatum.	terminal., semi-2-loc.	2, lobo dorsali auct., clas-tic. pedicello communi pel-tato suffult.	Paras. Caulis simpl. Fol. distich., carnos., subtriquetr. Flor. solitar.
OMOEA. [Ceratophilus.] .	praeceidentis.	parvum, inferne in calcar compress. emarginat. pro-duct.; limb. 3-fid.; lacin. media erect., incrassat.	praeceidentis.	praeceidentis.	praeceidentis.	praeceidentis sed caules subramosi.
MICROSACCUS.	erecto-patent., linearia, subaeq.	inferne in saccum depress. elongat.; limb. abbreviat., obtusus.	breve, latum, obtusum, antice sub apice emargina-tum.	terminal., semi-2-locul.	4, elasticae margini stig-matis insert.	Paras. Caulis simpl. Fol. distich., compressa, carnos., equitant. Pe-dunculi inrafoliac., pauciflori.
TRICHOGLOTTIS. [Epi-dendr. Sw. Dipodium. Brwn.]	patent.; lateral. labello su-persita.	inferne in calcar obtus. pro-duct.; limb. patens v. for-nicat., callos.	basi antice processu et me-dio ad apicem rostello in-struct.	terminal., 2-locul.	2, lobulo dorsali auct., pedicello comm. hamato suf-fult.	Paras. Caulis radicans. Folia li-neari-lanceolat. Pedunculi paucifl.
Sect. 1.	"	intus pubescens, ad mar-ginem utrinq. dente inflexo instruct.; limb. patens.	ad apicem subbicuspidat.	praeceidentis.	praeceidentis.	praeceidentis sed caules subramosi.
Sect. 2.	"	intus eberbe, dentibus la-teralibus obtusis; limb. ab-breviat., subfornicat.	obtusum.	"	praeceidentis.	
LIMODORUM. Sw. . . .	patent., subaeq.	inferne breviter calcarat., ba-si gynost. adnat., concav.; limb. sublobat., patens.	superne liberum, subdi-lat., antice infra apicem ros-tello lamellat. praedit.	ad apicem interne locat., 4-locellat.	8, angulat., filis elastic. cohaerent.	Terrestr. Rad. fibrose. Fol. mem-branac., petiolat. Scapus multifo-rus.
AMBLYGLOTTIS. [Limod. Sw.]	praeceidentis.	praeceident. sed secund. tor-longitudin. gynost. adnat.; limb. sublobat., patens.	breve, dilat., antice ad apicem rostello attenuat. in-struct.	terminal., 8-locellat.	8, clav., elasticae pedi-cello comm. suffult.	
AGROSTOPHYLLUM. . . .	patent.; exter. latiora.	saccat., gynost. inferne ad-nat., medio construct.; limb. truncat., cerosus.	superne liberum, interne ad apicem attenuato-rostel-lat.	terminal., 4-locellat.	8, elasticae pelliculo comm. suffult.	Paras. Caulis simpl. Folia distich., angusta, membranac. Flores dense capitat.

3. *Labellum sepalis lateralibus exterioribus sesserium.*

COLLABIUM. [Malaxis. Sw.]	patent.; lateral. inferne in calcar breve coalita.	unguiculat., int. callos.; limb. semilunat., patens.	obtusum.	terminal., 2-locul.	2, filis elastic. margini stig-matis aff.	Terrestr. Folia radical., membra-nac. Spica multifl.
TÆNIOPHYLLUM.	erecta; exterior. inferne connat.; lateral. calcar bre-ve æmulantia.	sepalis aequale.	apice interne rostello semi-bifid.	terminal., 2-valv.	4, elasticae ope pharetræ pelliculo stigmat. communi affixæ.	Paras. Folia radical., fasciculat., vermicularia. Flor. spicati.

Continuatio CLAVIS GENERUM ORCHIDEARUM JAVANICARUM.

CEREACEAE. POLLINIS MASSAE DETERMINATAE, FARINOSO-PULPOSAE AUT CEREACEAE.

B. GYNOSTEMIUM OVARIO INCUMBENS.

1. *Labellum articulatione distincta (emarginata) cum ungue gynostemii junctum.*

<i>Affinitas.</i>	<i>Sepala.</i>	<i>Labellum.</i>	<i>Gynostemium.</i>	<i>Anthera.</i>	<i>Masse pollinis.</i>	<i>Habitus.</i>
EPICRANTHES.	tria, aequalia, patent.	parv., patulum, glanduloso-tumidum.	basi intus processib. antenniformibus pendulis cinctum.	terminal., semi-2-loc.	2, margini stigmat. juxta-positae.	Parasit. Fol. bulb. suffult. Pedunculi e paleis rumpentes, 1-flori.
EPHIPPIUM. [Dulbophylli Spec. Th.]	ringent., ineq.; lateral. ungui gynost. insert.	parvum, crassum, indivis. (sellaform.), cum ungue gynost. arcuato articulatum.	anice ad apicem bicuspidatum v. biconatum.	praeceidentis.	2, bipartibiles margini stigmat. approximatae.	Parasit. Fol. bulb. suffult. Pedunculi pauci. aut multi-flori.
OSYRICERA.	subringent.; exter. majora; lateral. usque ad apicem fere coadunata.	concaevum, indivis., cum ungue gynost. calloso articulatum; limb. tumido-glandulosus.	breve, apice bialatum; alis emargin.	terminal., 2-loc., anice in ligulam glandulosam producta.	2, margini stigmat. juxta-positae.	Parasit. Fol. bulb. imposit. Flor. spicati, sessiles.
CALLOSTYLIS.	erecto-patent.; lateral. sub-tus connata.	convexum, indivis., ungue gynost. incurvo calloso articulat.	superne incurv., clavatum.	terminal., 3-locellat.	3, elastice margini stigmat. adherentes.	Parasit. Folia bulb. suffult. Pedunculi ad basin folior. erumpent, multiflor. Flor. pedicellati.

2. *Labellum articulatione spuria (at tamen in nonnullis elastica) aut nulla cum ungue gynostemii junctum.*

a. Pollinis massae duae, rarius solitariae, plerumque bipartibiles et nuda.

COCHLIA. [Oxyricera.]	erecta; ext. latiora, basi subcoalit.; later. ungui gynost. inserta.	basi angustat., superne concav., in limbum semilunatum revolut. expansum, ascendens.	apice interne bialat.; alis emarginatis.	terminal., semi-2-loc.	2, juxtapositae.	Parasit. Caules vaginat. Folia cochleata (unde nomen). Flores capitati.
DIPHYES [Ephippium.]	erecto-patent. v. connivent.; ext. basi subcoalit.; lateral. ungui gynost. oblique inserta; inter. minuta.	parvum, elastice gynost. ungui incurvo adherens.	breve, apice semibifid. rarius indivisum.	terminal., 1 v. sub-2-locularis.	1-2, saepe bipartibiles.	Parasit. Folia plerumq. bulb. suff. Flor. solitarii v. conferti v. spicati. (Herbae pusillae.)
SECT. Monanthes.	"	"	2-fidum.	2-locularis.	2, globose.	Folia bulbis hand imposita, disticha. Pedunculi breves, 1-flori.
" Diptychantes.	"	"	2-cuspidatum.	1-locularis.	1, bipartibilis.	Folia bulbis non imposita. Pedunculi breves pauciflori.
" Stachysanthides.	"	"	2-fidum.	1-locularis.	1, bipartibilis.	Caules radiceiformes, foliis bulb. suff. Flores laxae spicati.
" Desmosaundes.	"	"	semi-2-fidum.	2-locularis.	2, sulco dorsali inscriptae.	Caules radiceiformes, foliis bulb. suff. Flores capitati.
" Corymbosia.	"	"	semi-2-fidum.	2-locularis.	2, subglobosae, dorso sublobat.	Caules radiceiformes, foliis bulb. suff. Flores corymbosi.
" Polymeres.	"	"	2-cuspidatum rarius indivisum.	2-locularis.	2, subbipartibiles.	Caules radiceiformes, foliis bulb. suff. Pedunculi conferti, 1-flori.
" Diosperia.	"	"	2-fidum.	1-locularis.	1, bipartibilis.	Caules radiceiformes, foliis bulb. suff. Pedunculi subsolitarii, 1-flori.
" Macrocaulia.	"	"	semi-2-fidum v. integrum.	2-locularis.	2.	Caules radiceiformes, foliis bulb. suff. Pedunculi solitarii, elongati, 1-flori.
DENDROM.	erecto-patent.; lateral. circa basin labelli oblique producta, calcar v. saccum referentia.	sepalis lateralib. inferne inclusum, gynost. accumbens; limb. divis. v. indivis.	dorso ad apicem dense antherifero et utroque latere dense plerumque oblitterato instructum.	terminal., 2-locularis.	2, bipartibiles, ovato-v. ovali oblongae, pulposae-rentes.	Parasit. caulescens, floribus solitariis, fasciculatis aut racemosis.
SECT. Pedilonum.	patent. v. erectiuscula; lateral. reliquis latiora, in sacculum v. calcar interne connata.	angustum, gynostemio accumbens; limb. subintegerrim.	anice ad apicem rostellum lamellato praeditum.	depressa.	2, bipartibiles, nuda.	Parasit. Caules simplices, articulati. Racemi oppositifol., v. subterminal., pauci- aut multi-flori.
a	sepalis lateralibus exteriora saccata.	liberum, sepal. lateral. hand connata.	"	"	"	
b	sepalis lateralibus exteriora calcarata.	inferne sepalis lateral. plerumque adunatum.	"	"	"	
SECT. Onychium.	patent. v. erect.; exterior. basi subcoalit.; lateral. calcar referent.	unguiculat., interne saepe dense instruct.; limb. sublobat., patens.	praeceidentis.	praeceidentis.	praeceidentis.	Parasit. Caules articulati. Pedunculi racemosi. multiflori aut uniflori, oppositifolii v. subterminales.
a	patent.; interiora dorsali latiora.	unguis labelli interne denticulo instructus.	anice pubescens.	"	"	Caules inferne ebulbi. Flores dense racemosi.
b	erectiuscula; interiora dorsali subaequalia.	unguis labelli interne aut nudus aut denticulo munitus.	" nudum.	"	"	Caules inferne subbulbosi. Flores laxae racemosi v. remoti, paleis cincti.
SECT. Desmotrichum.	patent. v. erectiuscula, longitudine subaequalia; exter. basi subcoalit.; lateral. inferne oblique producta.	basi concavum, intus lineat.; limb. elongatus, sublobatus, patens.	Pediloni.	convexuscula.	praeceidentis.	Parasit. Caules radiceiformes. Folia bulbis suffulta, coriacea. Pedunculi ad basin foliorum orti, fasciculati, subracemosi v. solitarii, e paleis rumpentes.
SECT. Graetidum.	erecta, longitudine subaeq.; lateralibus calcar emulanti.	sepalis exterioribus subaequalibus, intus lineatum.	praeceidentis.	praeceidentis.	praeceidentis.	Parasit. Caules simplices, ebulbi. Folia submembranacea. Flores suboppositifol., e paleis cornicis erumpentes.
SECT. Aporum.	lateral. erect. patent., labellum inferne amplectent.; reliquis angustiorib., erectis.	basi concavum; limbus dilatatus, crista transversali instructus.	praeceidentis.	praeceidentis.	praeceidentis.	Parasit. Caules foliosi, ebulbi. Folia equitant., ensiformia, compressa, rigida. Flor. intraxilliaci, inferne paleis cincti.
SECT. Macrostromium.	reflexa; lateral. inferne in calcar breve connata.	angustat.; limbus elongatus, spatulat., concav., nudus.	praeceidentis.	praeceidentis.	2, subglobosae, inpartibiles.	
* SECT. Oxystophyllum.	erecto-patent.; exter. inferne subcoalta; lateralibus latiora, labellum inferne amplectentia.	ungui gynost. lato elastice affix., indivisum; limbus intricatus, externe tumidus.	praeceidentis.	praeceidentis.		
DIGLYPHOSA. [Oncidium. Sw.]	subringentia.	basi concav., elastice affixum; limb. indivisus, patens.	incurvum, obtusum.	terminal., 2-locul., anice bifida.	2, margini stigmat. insidentes.	Terrestr. Petioli basi subbulbosi, 1-phylli. Folia membranacea. Scapi radicales, multiflori.
CHRYSOGLOSSUM. [Trichoceras Humbd. Oncid. Sw.]	subringentia.	basi concav., elastice affixum; limb. lobat., patens.	utroque latere process. instruct.; unguis basi cucullatus.	terminal., 2-locul.	praeceidentis.	Parasit. Caules simplices, radican-tes, ebulbi. Flor. paniculati.
PHALAEENOPSIS. [Epidendr. Sw.]	patentia, inaeq.; interiora latiora.	cum ungue gynost. coarctum, basi callos.; limb. difform., 3 lobus.	ad apicem anice rostellum emarginato instruct.	gynostemium anice terminans, 2-locul.	2, dorso foveate, pedicello comm ad basin petiati suffult.	

Continuatio CLAVIS GENERUM ORCHIDEARUM JAVANICARUM.

CERACEAE. POLLINIS MASSAE DETERMINATAE, FARINOSO-PULPOSAE AUT CERACEAE.

B. GYNOSTEMIUM OVARIO INCUMBENS.


B. POLLINIS MASSAE 4-8, PLENUMQUE FILIS AUT STYLIS SUFFULTAE.

<i>Adfinitas.</i>	<i>Sepala.</i>	<i>Labellum.</i>	<i>Gynostemium.</i>	<i>Anthera.</i>	<i>Massae Pollinis.</i>	<i>Habitus.</i>
CERATOSTYLIS.	erectiuscula; ext. inferne subcohaerit; lateral. calcar emulans.	subinclusum, unguiculat., versus limbum incrassat., concav.	breve, latum, subbicornutum.	gynostem. postice termin., 8-locellat.	8, elastice membrana commun. suffultae.	Parasit., subcaulescens. Pedunculi breves, 1-2 flori, ad basin foliorum orti, paleis cincti.
APPENDICULA. [Octomeria Brwn.]	erecto-patent.; dorsali subfornicato; ext. basi subcohaerit; lateral. calcar obtus. referent.	concav., basi margine cristato cinctum (unde nonen); limbus patens, integerrim. v. sublobatus.	ad apicem in rostell. emarginat. attenuatum.	dorsalis, sub-8-locellat.	8, (nonnul. interdum steril.), clavace, glandula v. membranacea commun. suffultae.	Parasit. Caulis simpl. v. subramosus. Folia disticha, plerumque membranacea. Flores spicati.
SECT. 1.	sepala lateral. cum ungue labelli inferne rarius connata.	limbus integ.	"	"	"	Flores spicati.
SECT. 2.	sepala lateral. cum ungue labelli ad basin plerumque connata.	limbus sublobatus.	"	"	"	Flores capitati v. dense spicati.
CRYPTOGLOTTIS.	erecto-convivent.; ext. basi connat.; lateral. cum ungue emulans.	angustum, inclusum, basi extus bifidum, ope appendiculi unguis gynost. adherens.	parvum, apice rostell. emarginato.	dorsal., 2-locularis, locul. approxim.	4, ineq., styl. commun. suffult.	Caulis ebulbi. Spicae axillares v. terminales. Flores minui.
PODOCHILUS.	convivent.; ext. usque ad medium connat.; lateral. inferne venricosa.	concav., basi appendicib. 2-3-nerosis instruct. unguis gynost. elastice adherens.	parvum, apice 3-fidum; lacinia intermedia (rostellum Rich.) tricuspid.	praecedentis.	praecedentis.	praecedentis, sed flores solitarii, subterminales.
PLATYSMA.	Podochili, sed sepala ext. minus cohaerent., erecto-patentia; lateral. calcar referent.	Podochili, sed inferne inappendiculatum.	parvum, margine indivisum, apice 3-cuspe, antice ad apicem lamellatum.	praecedentis.	praecedentis.	Podochili.
APISTA.	Podochili.	Podochili.	latum, utrinque emarginatum.	dorsal.; 2-loc., loculis discretis.	4, per paria invicem disjunctae et stylis 2-suffult.	
CRINONIA.	erecto-patent.; lateral. labello supposita.	concav., basi obuse unguiculat.; limb. indivisus, patens.	utrinque dente laterali instructum.	dorsal., 4-locellat.	4, per paria filis junctae, margini stigmat. affixae.	Parasit. Folia bulbis suffult. Spicae ad basin bulborum orti, inferne vaginat. Bractee imbricat., caducae.
DENDROLIRIUM. [Brassavola Brwn.]	erecto-patent.; lateral. ext. unguis gynost. oblique imposit., inferne subcohaerit, sepe calcar obtusum referentia.	basi concav., unguis gynost. incumbens; limbus indivisus, v. sublobatus.	obtusum.	terminal., sub-8-locellat.	8, filis elastic. cohaerentes.	Parasit. Folia disticha, plerumque bulbis suffult. Spica multiflora.
SECT. 1. Cosmium.	lateral. ext. labello inferne amplectent.	gynost. longitudine superans, inferne callos. v. crispatum; limbus lobatus.	"	"	"	Spica plerumque bracteis coloratis ornata. Flores sat magni, rarius parvi. Foliorum bulbis compressi.
" 2. Brachystomium.	lateral. ext. unguis labelli ad basin subcohaerent.	abbreviat., inferne cucullat.; limbus indivisus.	"	"	"	praecedentis, sed floribus et bracteis minutis.
" 3. Odontium.	praecedentis.	inferne cucullatum et dentibus 2 lateralibus auctum; limbus indivisus.	"	"	"	praecedentis, sed bulbis elongatis, cauliformibus.
" 4. Rhizophyllum.	lateral. ext. labello supposita, erectiuscula, calcar non emulans.	praecedentis, sed edentulum, inferne concavum.	unguis gynostemii abbreviatus.	"	8, elastice pedicello commun. hamato suffult.	Bractee et flores minutissimi. Folia radicalia, disticha, basi vaginantia.
" 5. Cronmophyllum.	praecedentis.	praecedentis.	praecedentis.	"	praecedentis.	Bractee et flores praecedentis. Folia bulbis suffulta.
" 6. Anacamptium.	lateral. ext. inferne labello amplect., calcar referent.	basi angustum; limbus appendiculat.	unguis gynost. recurvus.	"	praecedentis.	
MYCARANTHES.	patentia; lateral. unguis gynost. oblique insert., reliquis lateralibus.	trilobum, elastice affix., basi et versus apicem callo instructum.	obtusum.	gynostem. postice termin., 8-locellat.	8, filis elasticis margini stigmat. affixae.	Parasit. Caulis simpl., ebulbi. Spica terminalis, multifi.
TRICHOTOSIA.	erecto-patent.; lateral. inferne producta, calcar emulans.	basi angustata, gynostemio accumbens; limb. sublobat. v. integerrimus.	obtusum.	terminal., 8-locellat., dentati dorsali affixae.	8, ovatae, compressae, filis elasticis cohaerentes.	Parasit. Caulis ebulbi. Pedunculi oppositifolii. (Herbe pilosae).
CERATIUM.	lateral. unguis gynost. inserta, erecto-patent., reliquis conviventibus lateralibus.	concav., inferne cristat., unguis gynost. elastice affix.; limb. sublobat., fornicatus.	apice cavitate pollinifera instructum.	terminal., 4-locellat., dentati dorsali inserta.	4, oblongae, lobulis dorsalis auctae et filis elastic. junct.	Praecedentis sed glaber.
SARCOSTOMA. [Ceratostylis]	erecto-patent.; lateral. oblique product., calcar referent.	unguiculat., superne gynost. amplectens; limbi lobo medio carnosus.	superne crassum, antice ad apicem rostellatum.	terminal., postica, 4-locellat.	4, obovat., per paria filis cohaerentes.	Parasit., ebulb., subcaulesc. Folia linearia, carnosae. Flores terminal., subsolitarii.
ACANTHOPHIPIUM.	venricosum-convivent.; lateral. latior, cum ungue gynost. connat.	parvum, carnosum, inuspectum, unguis gynost. geniculato elastice junctum.	obtusum.	antice ad apicem locata, 2-locular.	8, ineq., membran. elast. bifidae insidentes.	Terrestr. Caulis inferne bulbosi. Folia membranacea. Scapus radicalis, vaginatus, pauciflorus.
TAINIA. [Nephelaphyllum.]	ringent., subaequal.; lateral. unguis gynost. affix.	concav., indivisus; limb. acuminatus.	semiteres.	terminal., sub-8-locul., angulat.	8, ineq., membran. elast. cohaerent.	Terrestr. Caul. radicans. Fol. petiol., membranac. Scapus lateral., spicato-multifi.


3. Labellum cum ungue gynostemii inferne in saccum v. calcar connatum.

BACCOLABIUM. [Aërides Sw.; Cleisostoma.]	patentia; lateral. latiora.	inferne in saccum obtusum elongat.; limb. truncatus, tumidus.	ad apicem antice rostellum instructum.	terminal., semi-2-locul.	2, globose, pedicello communi ad basin hamato affixae.	Parasit. Caulis simplices, compressi. Folia disticha. Flores axillares, solitarii.
RHYNCHOSTYLIS. [Aërides Sw.]	patent.; lateral. inferne producta.	inferne in saccum compress. product.; limb. ovatus, patens.	apice rostellum elongato praeditum.	terminal., semi-2-locul.	2, dorso sulco inscriptae, pedicello longissimo ad basin appendiculato suffultae.	Parasit. Caulis simplices. Flores dense spicati.
DENDROCOLLA. [Aërides Sw.; Cleisostoma.]	patentia; lateral. inferne latiora.	inferne saccat. v. calcarat.; limbus cucullatus, subulatus, callosus aut fornicatus.	obtusum, antice ad apicem emarginatum.	terminal., 2-locular.	2, dorso sublobat., pedicello comm. ad basin petalato affixae.	Parasit. Caulis radicans. Pedunculi multiflori.
SECT. Cucula.	"	limbus cucullat. rarius expansus.	"	"	"	Pedunculi bracteis pectinatis instructi.
" Tubera.	"	limbus abbreviatus, callosus.	"	"	"	Pedunculi bracteis scrobiculatis coalitis muniti.
" Fornicaria.	"	limbus subfornicatus.	"	"	"	Pedunculi bracteis subulatis confertis praediti.
CISTELLA. [Maxalis Willd.]	erecta; lateral. labello supposita, carinata.	erect., inferne saccatum; limbus indivisus.	superne liberum, latum.	gynost. antice termin., 2-locular.	2, basi foveatae, juxtaposita.	Terrestr. Folia subradicalia, membranacea. Spica multiflora.


I




II




III




IV




V




VI




VII

Chrysoglossum villosum


VIII

Trichoglottis retusa


IX

Epicranthes Javanica


X


Dendrobium lamellatum

XI

Trichosia pauciflora


XII

Podochilus lucescens




XIII

Habenaria Salaccensis




XIV

Elaeria Oblongifolia




XV

Anecochilus Setaceus




XVI

Cheirostylis Montana




XVII

Adenostylis Integerrima




XVIII

Eucosia Carneae




XIX

Cymbidium Aloefolium




Cymbidium Tavanicum




XX

Grammatophyllum Speciosum.




XXI

Plocoglottis Javanica




XXII

Nephetaphyllum Pulchrum




X XIII


Schoenorchis Funicifolia




XXIV

Acerides Matutina


XXV

Ceratochilus Biglandulosus


XXVI

Arachnis Moschifera

XXVII

Gleisostoma Sagittata


XXVIII

Echioglossum Javanicum

Léon Del.

XXIX

Pachystoma pubescens




XXX

Peristylis Grandis




XXXI

Chlorosa latifolia




XXXII

Zosterostylis Arachnites



XXXIII

Calcearia fornicata



Calcearia picta




XXXIV

Vanilla Albida




Latour del⁴




XXXV





Desmotrichum Angulatum*Desmotrichum pusillum*

XXXVI

Pedilonum undulatum


XXXVII

Macrostomium alceifoliumXXXVIII *Oxystophyllum Carnosum*XXXIX *Aporum indivisum*




XL

Appendicula alba




Appendicula cristata




XLl

Crinonia carnea




XLII

Cryptoglottis serpillifolia




XLIII

Platysma gracilis




XLIII

Phalaenopsis amabilis




XLV

Sarcostoma javanica




XLVI

Ceratium compressum




XLVII

Acanthophippium javanicum




XLIX

Rhynchostylis retusa



XLVIII




Tainia speciosa



L


Saccolabium pusillum





LI


dg sect. 1. *Chelonanthera ventricosa*



sect. 2 *Chelonanthera speciosa*




sect. 3 *Chelonanthera longifolia*




LII

sect 1. *Dendrochilum pallide flavens*





sect 2. *Dendrochilum simile*




LIII

Agrostophyllum javanicum







Malaxis Rheduvaret




Malaxis compressa




Malaxis affinis




Malaxis decurrens




Cistella cernua



Ceratostylis graminifolia



Mycaranthes latifolia



LVIII

Oxyricera crassifolia

The illustration consists of several numbered parts: 1. A small detail of a leaf or petal. 2. A long, narrow, pointed structure, possibly a seed or fruit, with labels 'sd', 'si', 'l', 'sl', and '2'. 3. A complex structure showing a flower or fruit with labels 'a', 'g', 'cg', 'l', 'lg', and '3'. 4. A structure with two long, curved, pointed lobes, labeled 'cg', 'ca', 'g', and '4'. 5. A structure with two long, curved, pointed lobes, labeled 'cg', 'ca', 'mp', 'cg', 's', 'cg', and '5'. 6. A structure with two long, curved, pointed lobes, labeled 'cg', 'ds', 'o', and '6'. 7. A structure with two long, curved, pointed lobes, labeled 'cg', 'ds', 'o', and '7'. 8. A structure with two long, curved, pointed lobes, labeled 'cg', 'ds', 'o', and '8'.

LIX

Cochlia violacea

Figure 1: Lateral view of the head and thorax. Labels: *si*, *sl*, *sd.*, *o.*

Figure 2: Lateral view of the head and thorax. Labels: *lg*, *cg*, *la.*, *g.*, *o.*, *u g*

Figure 3: Dorsal view of the head and thorax. Labels: *a*, *cg*, *ea*, *s*, *l*

Small heart-shaped structure to the right of Figure 3.

LX *Diglyphosa latifolia*

1 2 3 4 5

[illegible]



LXII

Limatodis pauciflora

This botanical illustration depicts the morphology of *Limatodis pauciflora*. It includes several numbered figures: 1. A flowering branch with leaves and a terminal inflorescence, labeled with 'sl' (sepal), 'sd' (stamen), 'cl' (calyx), and 'b' (bract). 2. A detailed view of a flower, labeled with 'a' (anther), 'g' (gynoecium), 'l' (lobes), 'o' (ovary), and 'cl' (calyx). 3. A longitudinal section of the ovary, labeled with 'mp' (marginal process), 'r' (receptacle), 's' (stigma), 'cl' (calyx), and 'b' (bract). 4. A longitudinal section of the ovary, labeled with 'ca' (carpel), 'r' (receptacle), 's' (stigma), 'ds' (dorsal side), 'cl' (calyx), and 'b' (bract). 5. A longitudinal section of the ovary, labeled with 'o' (ovary) and 'cl' (calyx). 6. A longitudinal section of the ovary, labeled with 'o' (ovary) and 'cl' (calyx). 7. A longitudinal section of the ovary, labeled with 'o' (ovary) and 'cl' (calyx). The artist's signature 'Latour Del.' is at the bottom left.


LXIII

Crepidium Rhedii




LXIV

Amblyglottis flava




LXV

Ephippium ciliatum




Ephippium cornutum




LXVI


Diphrys gibbosa. Sect. *Stachysanth.*




Diphrys tenella. Sect. *Macrocaul.*




Diphrys bicolor. Sect. *Polymer.*




Diphrys crassifolia. Sect. *Desmosanth.*






LXVII

Dendrocolla arachnites Sect. 1




Dendrocolla emarginata Sect. 2.




LXVIII

Glomera erythrosma





LXIX


Dendrolirium flavescens Sect. 1.




Dendrolirium acuminatum Sect. 2.



Dendrolirium pusillum Sect. 4




Dendrolirium appendiculatum Sect. 6.




Latour, Del.


LXX *Tacniophyllum fornicatum*



LXXI. *Acriopsis javanica*



LXXII. *Erythroides latifolia*



LXXIII. *Arundina speciosa*



UB WIEN



+AM469936807

