

Research Article

SPECIAL ISSUE: Island Plant Biology—Celebrating Carlquist's Legacy

Genetic consequences of cladogenetic vs. anagenetic speciation in endemic plants of oceanic islands

Koji Takayama¹, Patricio López-Sepúlveda², Josef Greimler³, Daniel J. Crawford⁴, Patricio Peñailillo⁵, Marcelo Baeza², Eduardo Ruiz², Gudrun Kohl³, Karin Tremetsberger⁶, Alejandro Gatica⁷, Luis Letelier⁸, Patricio Novoa⁹, Johannes Novak¹⁰ and Tod F. Stuessy^{3,11*}

¹ Museum of Natural and Environmental History, Shizuoka, Oya 5762, Suruga-ku, Shizuoka-shi, Shizuoka 422-8017, Japan

² Departamento de Botánica, Universidad de Concepción, Casilla 160-C, Concepción, Chile

³ Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, A-1030 Vienna, Austria

⁴ Department of Ecology and Evolutionary Biology and the Biodiversity Institute, University of Kansas, Lawrence, KS 60045, USA

⁵ Instituto de Ciencias Biológicas, Universidad de Talca, 2 Norte 685, Talca, Chile

⁶ Institute of Botany, Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences, Gregor Mendel Straße 33, A-1180 Vienna, Austria

⁷ Bioma Consultores S.A., Mariano Sanchez Fontecilla No. 396, Las Condes, Santiago, Chile

⁸ Universidad Bernardo O'Higgins, Centro de Investigaciones en Recursos Naturales y Sustentabilidad, General Gana 1702, Santiago, Chile

⁹ Jardín Botánico de Viña del Mar, Corporación Nacional Forestal, Camino El Olivar 305, Viña del Mar, Chile

¹⁰ Institute for Applied Botany and Pharmacognosy, University of Veterinary Medicine, Veterinärplatz 1, A-1210 Vienna, Austria

¹¹ Herbarium, Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, 1315 Kinnear Road, Columbus, OH 43212, USA

Received: 17 January 2015; **Accepted:** 27 July 2015; **Published:** 26 August 2015

Guest Editor: Clifford Morden

Citation: Takayama K, López-Sepúlveda P, Greimler J, Crawford DJ, Peñailillo P, Baeza M, Ruiz E, Kohl G, Tremetsberger K, Gatica A, Letelier L, Novoa P, Novak J, Stuessy TF. 2015. Genetic consequences of cladogenetic vs. anagenetic speciation in endemic plants of oceanic islands. *AoB PLANTS* 7: plv102; doi:10.1093/aobpla/plv102

Abstract. Adaptive radiation is a common mode of speciation among plants endemic to oceanic islands. This pattern is one of cladogenesis, or splitting of the founder population, into diverse lineages in divergent habitats. In contrast, endemic species have also evolved primarily by simple transformations from progenitors in source regions. This is anagenesis, whereby the founding population changes genetically and morphologically over time primarily through mutation and recombination. Gene flow among populations is maintained in a homogeneous environment with no splitting events. Genetic consequences of these modes of speciation have been examined in the Juan Fernández Archipelago, which contains two principal islands of differing geological ages. This article summarizes population genetic results (nearly 4000 analyses) from examination of 15 endemic species, involving 1716 and 1870 individuals in 162 and 163 populations (with amplified fragment length polymorphisms and simple sequence repeats, respectively) in the following genera: *Drimys* (Winteraceae), *Myrceugenia* (Myrtaceae), *Rhaphithamnus* (Verbenaceae), *Robinsonia* (Asteraceae, Senecioneae) and *Erigeron* (Asteraceae, Astereae). The results indicate that species originating

* Corresponding author's e-mail address: stuessy.1@osu.edu

Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

anagenetically show high levels of genetic variation within the island population and no geographic genetic partitioning. This contrasts with cladogenetic species that show less genetic diversity within and among populations. Species that have been derived anagenetically on the younger island (1–2 Ma) contain less genetic variation than those that have anagenetically speciated on the older island (4 Ma). Genetic distinctness among cladogenetically derived species on the older island is greater than among similarly derived species on the younger island. An important point is that the total genetic variation within each genus analysed is comparable, regardless of whether adaptive divergence occurs.

Keywords: Adaptive radiation; anagenesis; cladogenesis; genetic diversity; phyletic speciation; Robinson Crusoe Islands.

Introduction

Oceanic islands have long stimulated biologists to investigate patterns and processes of evolution (e.g. Darwin 1842; Wallace 1881; Whittaker and Fernández-Palacios 2007; Bramwell and Caujapé-Castells 2011). These isolated land masses, far from continental source areas, offer opportunities for determining origins of immigrants and their evolutionary history after establishment. The low probability of long-distance dispersal and successful colonization, the reduction of genetic variation in founding populations and the challenges of adaptation to new environments are all features that combine to affect processes of evolution in island archipelagos, particularly speciation.

One dimension of speciation in island plants that has received considerable attention is adaptive radiation (Carlquist 1974; Whittaker and Fernández-Palacios 2007; Rundell and Price 2009). This is a process that begins with dispersal from the original immigrant population into different habitats on the same or neighbouring island. This isolation leads to divergence of the new segregate populations, each becoming rapidly adapted to divergent habitats (Schluter 2001), such that eventually new species are recognized taxonomically. This general process of speciation is usually diagrammed (Fig. 1) as splitting events or cladogenesis (Rensch 1959). A number of dramatic species complexes have developed in oceanic islands through adaptive radiation, such as illustrated by the lobelioids (Givnish et al. 2009) and silverswords (Carlquist et al. 2003) in Hawaii, *Aeonium* (Liu 1989; Jorgensen and Olesen 2001) and *Echium* (Böhle et al. 1996) in the Canary Islands and *Scalesia* (Eliasson 1974) in the Galapagos archipelago.

In addition to speciation via adaptive radiation (involving cladogenesis), another process, anagenesis (Fig. 1), has recently been emphasized (Stuessy et al. 1990, 2006; Whittaker et al. 2008). Some immigrant populations, especially when arriving on an island with limited ecological opportunity, proliferate in size and accumulate genetic diversity mainly through mutation and recombination. After many generations (perhaps over a million or more years), genetic changes result in different morphology

that may be treated as a distinct species. This process has been labelled anagenetic speciation (Stuessy et al. 2006), being one type of progenitor-derivative speciation (Crawford 2010). It has been estimated that at least one-quarter of all endemic plant species of oceanic islands have originated via anagenesis (Stuessy et al. 2006).

Some studies have been published on the genetic consequences of cladogenesis in endemic plants of different archipelagos. Böhle et al. (1996) examined chloroplast sequence variation among endemic species of *Echium* (Boraginaceae) of the Canary Islands, showing very little nucleotide divergence even though the morphological variation is striking. Likewise, Baldwin (2003) examined internal transcribed spacer regions of nuclear ribosomal DNA (ITS) variation among species of the Hawaiian silverswords (Asteraceae) and again, limited sequence variation was seen. The general result from these, and other studies, is that during cladogenesis, the immigrant population becomes fragmented, with each segment containing a limited range of genetic variation in comparison with the continental progenitor population (Baldwin et al. 1998). Maximum morphological divergence occurs but with low levels of observable genetic diversity (Frankham 1997). There is some evidence (Perugganan et al. 2003) that the genetic changes responsible for the morphological adaptations involve alterations in regulatory rather than structural genes.

Results so far with anagenesis show a strikingly different pattern. Most of the investigations have been done on endemic species of Ullung Island, in which at least 88 % of the endemic species have originated anagenetically (Stuessy et al. 2006). The island is young (1.8 Ma; Kim 1985), of low elevation (<1000 m) and relatively ecologically uniform (Yim et al. 1981). Pfosser et al. (2005), using amplified fragment length polymorphisms (AFLPs), examined island and Japanese populations of *Dystaenia takesimana* and *D. ibukiensis*, respectively, and the results showed high levels of genetic variation within *D. takesimana* in comparison with *D. ibukiensis*. Similar results have been obtained in assessing the origin of *Acer takesimensis* and *A. okomotoanum* (Takayama




Figure 1. Diagram of the two principal modes of speciation in oceanic islands.

et al. 2012, 2013a). Because there is no partitioning of the immigrant population, it survives and proliferates, during which time it accumulates genetic variation through mutation and recombination. Eventually, the level of genetic diversity may even equal (or surpass) that observed in parental source populations (Stuessy 2007).

Because the above studies have been done on different genera in different island archipelagos, it would be useful to compare the genetic consequences of both types of speciation within groups of the same island system, preferably within the same island. In this fashion, more direct comparisons can be made because the general environment is the same. Important, obviously, is to locate plant groups that have originated via both anagenesis and cladogenesis within the same archipelago. A good choice for examining the genetic consequences of anagenesis and cladogenesis in endemic plants of oceanic islands is the Juan Fernández Archipelago, Chile. Approximately 64 % of the species have originated by cladogenesis and 36 % by anagenesis (Stuessy et al. 2006). From another perspective, it is estimated that 70 % of the colonists to the islands have diverged anagenetically, in contrast to only 30 % that have diverged via adaptive radiation (Stuessy et al. 1990).

The Juan Fernández Archipelago consists of two major islands (Fig. 2): Robinson Crusoe (= Masatierra), located 667 km west of continental Chile at 33°S latitude, and Alejandro Selkirk (= Masafuera) situated 181 km further westward into the Pacific Ocean. The former is known to be ~4 million years old and the latter 1–2 million years old (Stuessy et al. 1984). At present, these two islands are approximately the same size of 50 km² (Stuessy 1995). The flora is small, containing 78 native and 135 endemic vascular plant species (Danton et al. 2006). From a biogeographic standpoint, this setting is particularly favourable for generating initial hypotheses, because the near island (Robinson Crusoe) is also the older one, making it highly probable as the initial site for colonization of most groups. Furthermore, the older island is hypothesized to have been much larger when formed (Stuessy et al. 1998), making it a bigger target for dispersal from the mainland.




Figure 2. Location of the Juan Fernández Archipelago and its two major islands, Alejandro Selkirk (= Masafuera) and Robinson Crusoe (= Masatierra).

Numerous molecular markers now exist for assessing genetic variation within and among populations (Lowe et al. 2004). Amplified fragment length polymorphisms (Vos et al. 1995) have been used effectively to provide an overall evaluation of population genetic diversity (Tremetsberger et al. 2003; López-Sepúlveda et al. 2013a). These are treated as dominant markers and hence cannot be employed to determine allelic frequencies. An appropriate co-dominant and polymorphic marker that does allow allelic calculations are nuclear microsatellites or simple sequence repeats (SSRs). The challenge with this marker is to develop primers for locating sequences within the genome for comparison. Next-generation sequencing (NGS) methods are now available that allow this to be done much more easily and at reasonable cost (Takayama et al. 2011, 2013b). Numerous successful applications of SSRs have shown their efficacy to reveal genetic variation at the population level (Gleiser et al. 2008; Kikuchi et al. 2009; López-Sepúlveda et al. 2013b).

Studies using AFLPs and SSRs have already been published on a number of endemic taxa of the Juan Fernández Archipelago, representing groups that have undergone speciation via cladogenesis and anagenesis. The largest (and endemic) genus that has been investigated is *Robinsonia*

(Asteraceae; Takayama et al. 2015), which has seven species on Robinson Crusoe Island that have originated cladogenetically and one on Alejandro Selkirk Island that has evolved anagenetically. The genus *Erigeron* (Asteraceae; López-Sepúlveda et al. 2015) has six species that evolved cladogenetically on the younger island, Alejandro Selkirk. These two genera were selected because *Robinsonia* has speciated primarily via cladogenesis on the older island, and *Erigeron* has done so on the younger island. Regarding anagenesis, studies have been completed on *Drimys confertifolia* (Winteraceae; López-Sepúlveda et al. 2014) and *Rhaphithamnus venustus* (Verbenaceae; P. López-Sepúlveda, K. Takayama, D. J. Crawford, J. Greimler, P. Peñailillo, M. Baeza, E. Ruiz, G. Kohl, K. Tremetsberger, A. Gatica, L. Letelier, P. Novoa, J. Novak, T. F. Stuessy, submitted for publication), which occur on both islands of the archipelago. Investigations have also been completed on *Myrceugenia* (Myrtaceae; López-Sepúlveda et al. 2013b), which contains one endemic species on each of the islands. The available genetic data to date, therefore, come from 15 endemic species, plus 4 close continental relatives, summing to 1870 individuals in 163 populations.

The purposes of this article are to (i) summarize published data from AFLP and SSR investigations on endemic species of the genera *Drimys*, *Myrceugenia*, *Rhaphithamnus*, *Robinsonia* and *Erigeron*; (ii) compare and contrast differences in genetic diversity in groups that have undergone anagenetic or cladogenetic speciation and (iii) discuss the importance of considering modes of speciation for understanding levels of genetic diversity within endemic species of oceanic archipelagos.

Methods

The data summarized here (Table 1) provide the first comprehensive genetic comparisons (with AFLPs and SSRs) in the Juan Fernández Archipelago of species that have evolved by anagenesis and cladogenesis, based on consistent samplings, laboratory methods and modes of analysis. A number of earlier studies utilizing isozymes and DNA sequences have examined genetic variation in endemic species of these islands (e.g. Crawford et al. 1998, 2001a), but these investigations were not focussed on comparing modes of speciation. Genera in the present studies were selected for their representation of anagenesis and cladogenesis and for their occurrence on the two islands of different geological ages. The samples were collected during expeditions in February 2010 and 2011 from 1870 individuals in 163 populations in 15 endemic species, hence representing 14 % of the endemic angiosperms in the archipelago. The samples provide very good geographic coverage of populations over the landscape in

both islands. The term population, as used here in the sense of sampling, refers to groups of individuals that were clearly delimited spatially in the field. The number of individuals analysed per population ranged from 1 to 31. The voucher data for these samples and details of data gathering and analysis are given in the respective publications.

Briefly, the following approaches were used for AFLPs. Four or six selective primer combinations were chosen. Numerous (24–85) primer trials were run with each genus to determine the best combination of primers for good resolution of individuals and populations. Data were obtained on an automated DNA sequencer (ABI 3130xl, Applied Biosystems, Waltham, MA, USA). Scoring was done using GeneMarker ver. 1.85 (SoftGenetics, State College, PA, USA). For analysis of AFLP data, the programs Arlequin 3.5.1.2 (Excoffier et al. 2005), FAMD ver. 1.25 (Schlüter and Harris 2006), R-Script AFLPdat (Ehrich 2006) and SPSS ver. 15.0 (SPSS; IBM, Armonk, NY, USA) were employed to determine total number of fragments (TNB), percentage of polymorphic fragments (PPB), Shannon Diversity Index (SDI), average gene diversity over loci (AGDOL) and rarity index (RI).

For SSRs, NGS methods (Takayama et al. 2011) were used to generate 6–12 loci, selected for their repeatability and scoring convenience. Polymerase chain reaction-amplified fragments were also run on the same automated sequencer and scored with GeneMarker ver. 1.85. Data analysis involved using GENEPOP 4.0 (Raymond and Rousset 1995), Micro-Checker 2.2.3 (van Oosterhout et al. 2004), FSTAT 2.9.3.2 and GENALEX 6 (Peakall and Smouse 2006). These allow analyses for observed proportion of heterozygotes (H_O), expected proportion of heterozygotes (H_E), number of alleles per locus (N_A), inbreeding coefficient (F_{IS}) and allelic richness standardized by five individuals (A_{R5}).

The overall pattern of higher genetic diversities in anagenetically derived species in comparison with cladogenetically derived ones was examined by a Student's *t*-test (average TNB, PPB, SDI, AGDOL and RI in AFLPs, and H_O , H_E , N_A and A_{R5} in SSRs) and shown in Table 2. To improve normality of H_O and H_E , a square-root transformation was applied. The overall patterns of higher genetic diversities in Robinson Crusoe Island (old) than Alejandro Selkirk Island (new) were also examined in the same way. The effects of two factors (speciation mode and island) and their interaction were analysed in a two-way ANOVA in R version 3.0.0 (R Core Team 2013) and shown in Table 3.

Data from both AFLPs and microsatellites were further analysed by assessing genetic distance (Nei et al. 1983) with the NeighborNet algorithm (Bryant and Moulton 2004) implemented by SplitsTree4 ver. 4.10 (Huson and Bryant 2006) and Population 1.2.30 (Langella 1999), respectively.

Table 1. Summary of measures of genetic diversity in endemic species of the Juan Fernández Archipelago that have originated by anagenesis or cladogenesis. All average values. Data from López-Sepúlveda *et al.* (2013a, b, 2014), Takayama *et al.* (2015) and P. López-Sepúlveda, K. Takayama, D. J. Crawford, J. Greimler, P. Peñailillo, M. Baeza, E. Ruiz, G. Kohl, K. Tremetsberger, A. Gatica, L. Letelier, P. Novoa, J. Novak, T. F. Stuessy, submitted for publication. TNB, total number of bands (fragments); PPB, percentage of polymorphic bands; SDI, Shannon Diversity Index; AGDOL, average gene diversity over loci; RI, rarity index; H_O , observed proportion of heterozygotes; H_E , expected proportion of heterozygotes; N_A , number of alleles per locus; F_{IS} , inbreeding coefficient; A_{R5} , allelic richness standardized by five individuals; RC, Robinson Crusoe Island; AS, Alejandro Selkirk Island.

Species	AFLPs							Microsatellites (SSRs)						
	No. of pops.	No. of plants	TNB	PPB	SDI	AGDOL	RI	No. of pops.	No. of plants	H_O	H_E	N_A	F_{IS}	A_{R5}
Anagenesis														
<i>D. confertifolia</i> (RC)	16	183	557	96.5	125.3	0.26	1.96	16	181	0.48	0.68	9.00	0.29	4.12
<i>D. confertifolia</i> (AS)	15	96	538	96.5	114.3	0.23	2.26	15	80	0.35	0.51	6.38	0.26	3.24
<i>D. confertifolia</i> (combined RC and AS)	31	279	576	100	134.7	0.28	2.06	31	261	0.44	0.68	9.88	0.33	4.13
<i>M. fernandeziana</i> (RC)	18	211	371	100	74.6	0.23	1.76	18	231	0.38	0.49	10.08	0.19	3.38
<i>M. schulzei</i> (AS)	13	129	417	100	96.2	0.28	3.39	13	155	0.39	0.61	10.33	0.35	3.79
<i>R. venustus</i> (RC)	20	143	440	99.3	96.4	0.25	2.80	20	140	0.17	0.23	4.22	0.31	1.83
<i>R. venustus</i> (AS)	4	18	271	57.3	60.8	0.18	2.34	4	11	0.30	0.34	2.33	0.13	2.12
<i>R. venustus</i> (combined RC and AS)	24	161	443	100	98.7	0.26	2.75	24	151	0.18	0.28	4.56	0.40	2.04
<i>R. masafuerae</i> (AS)	5	9	344	41.4	84.1	0.15	2.90	5	7	0.36	0.43	3.50	0.17	3.08
Cladogenesis														
<i>Robinsonia gayana</i> (RC)	10	123	592	77.2	111.0	0.16	2.39	10	134	0.34	0.42	6.30	0.28	3.04
<i>R. gracilis</i> (RC)	5	75	515	63.2	97.3	0.15	2.68	5	87	0.28	0.39	3.50	0.24	2.26
<i>R. evenia</i> (RC)	6	73	586	73.4	112.0	0.17	3.18	6	86	0.21	0.26	2.80	0.21	1.87
<i>R. saxatilis</i> (RC)	1	5	267	29.0	67.0	0.14	1.99	1	5	0.30	0.26	2.10	-0.22	2.10
<i>Robinsonia</i> (combined all RC species)	22	276	765	100	183.7	0.26	2.77	22	312	0.28	0.66	8.40	0.61	3.97
<i>Robinsonia</i> (combined all species)	27	285	766	100	265.0	0.26	2.68	27	319	0.29	0.67	8.70	0.61	4.02
<i>E. fernandezianus</i> (RC)	13	240	403	90.3	70.7	0.20	0.58	13	271	0.21	0.29	4.20	0.31	2.17
<i>E. fernandezianus</i> (AS)	19	172	426	95.3	81.1	0.23	0.81	19	200	0.17	0.50	7.50	0.72	3.27
<i>E. fernandezianus</i> (combined RC and AS)	32	412	433	97.5	81.7	0.23	0.68	32	471	0.20	0.40	8.00	0.64	2.86
<i>E. ingae</i> (AS)	2	21	315	61.3	62.0	0.18	0.62	2	25	0.20	0.34	2.90	0.55	2.04
<i>E. luteoviridis</i> (AS)	2	25	334	61.5	60.2	0.18	0.99	2	25	0.05	0.31	3.10	0.72	2.19
<i>E. rupicola</i> (AS)	9	175	377	81.8	69.5	0.20	0.67	9	211	0.17	0.36	4.40	0.57	2.43
<i>E. turricola</i> (AS)	3	10	269	49.3	57.6	0.19	0.50	3	10	0.24	0.53	3.40	0.57	2.94
<i>E. stuessyi</i> (AS)	1	8	306	66.7	82.4	0.28	0.81	2	11	0.20	0.25	2.10	0.53	1.89

Continued

Table 1. Continued

Species	AFLPs		Microsatellites (SSRs)											
	No. of pops.	No. of plants	TNB	PPB	SDI	AGDOL	RI	No. of pops.	No. of plants	H _O	H _E	N _A	F _{IS}	A _{RS}
<i>Erigeron</i> (combined all AS species)	36	411	443	100	95.1	0.26	0.74	37	482	0.17	0.62	9.20	0.76	2.85
<i>Erigeron</i> (combined all species)	49	651	444	100	94.2	0.26	0.68	50	753	0.18	0.56	9.50	0.73	3.46
Total and averages														
Anagenesis	91	789	419.7	84.4	93.1	0.23	2.49	91	805	0.35	0.47	6.55	0.24	3.08
Cladogenesis	71	927	399.1	68.1	79.2	0.19	1.38	72	1065	0.2	0.4	3.8	0.41	2.38
Robinson Crusoe	89	1053	466.4	78.6	94.3	0.19	2.17	89	1135	0.30	0.38	5.28	0.20	2.60
Alejandro Serkirk	73	663	359.7	71.1	76.8	0.21	1.53	74	735	0.24	0.42	4.59	0.46	2.70
Anagenesis (RC)	54	537	456.0	98.6	98.8	0.24	2.17	54	552	0.34	0.47	7.77	0.26	3.11
Anagenesis (AS)	37	252	392.5	73.8	88.9	0.21	2.72	37	253	0.35	0.47	5.64	0.23	3.06
Cladogenesis (RC)	35	516	472.6	66.6	91.6	0.16	2.16	35	583	0.27	0.32	3.78	0.17	2.29
Cladogenesis (AS)	36	411	337.8	69.3	68.8	0.21	0.73	37	482	0.17	0.38	3.90	0.61	2.46

For this article, to allow ease of visual comparisons of results among the species, emphasis has been placed on selected graphic presentations. SplitsTree Neighbor-Net was employed with the AFLP data, and the results are given in a series of graphs (Fig. 3). Neighbour-joining based on genetic distance was used for analysis of the SSRs, and simplified networks were used to show relationships among the populations (Fig. 4). For summary comparisons of genetic diversity among species, AGDOL was used with the AFLP data (Fig. 5). Not all calculated values for all original populations are presented or discussed in this review. The reader is referred to the original publications for additional methods and data.

Results

The results from the AFLP and SSR data analyses are given in Tables 1–4 and shown graphically in Figs 3–5. In general, the results from the two sources of genetic data are similar, with some exceptions, reinforcing confidence in the patterns seen. These data will be presented in context of the two modes of speciation, anagenesis and cladogenesis, but with attention also to the different ages of the islands. Robinson Crusoe Island is ~4 million years old and Alejandro Selkirk 1–2 million (Stuessy et al. 1984).

Anagenesis

The results from analysis of species that have evolved anagenetically include those from *Myrceugenia fernandeziana*, *M. schulzei*, *Robinsonia masafuerae*, *D. confertifolia* and *R. venustus*. The first species occurs only on the older island, the second and third species only on the younger island and the last two on both islands. A number of points seem evident. First, all anagenetically derived species show considerable levels of genetic diversity (Table 1, and Figs 3 and 5), and none of them shows geographic patterns over the island landscape (López-Sepúlveda et al. 2013b, 2014, P. López-Sepúlveda, K. Takayama, D. J. Crawford, J. Greimler, P. Peñailillo, M. Baeza, E. Ruiz, G. Kohl, K. Tremetsberger, A. Gatica, L. Letelier, P. Novoa, J. Novak, T. F. Stuessy, submitted for publication). This is what might be expected from the predictions regarding anagenesis based on previous studies. Even more interesting, perhaps, is that the amount of genetic diversity differs in species on the two islands of different ages. In *D. confertifolia*, and *R. venustus*, which occur on both islands, one sees in both cases more genetic diversity (SDI) in populations on the older island than on the younger island except for estimates of SSRs in *R. venustus* (Table 1). The explanation of these data may relate to the time available for a genetic change to take place. Because Alejandro Selkirk Island is no more than 1–2 million years old, this must be the maximum

Table 2. Summary of statistical tests based on Table 1. TNB, total number of bands (fragments); PPB, percentage of polymorphic bands; SDI, Shannon Diversity Index; AGDOL, average gene diversity over loci; RI, rarity index; H_O , observed proportion of heterozygotes; H_E , expected proportion of heterozygotes; N_A , number of alleles per locus; A_{R5} , allelic richness standardized by five individuals. Bold font indicates significant values ($P < 0.05$).

	High genetic diversity in anagenetically derived species	High genetic diversity in Robinson Crusoe Island species
AFLPs		
TNB	0.351	0.024
PPB	0.086	0.235
SDI	0.101	0.045
AGDOL	0.050	0.227
RI	0.004	0.085
SSRs		
H_O	0.006	0.132
H_E	0.061	0.236
N_A	0.040	0.308
A_{R5}	0.038	0.388

time available for population divergence to take place. With anagenetically evolved species, all factors being equal, genetic variation increases through time, and this can be seen in the species investigated.

One case of anagenesis in the archipelago also merits comment. *Robinsonia masafuerae* is a species that appears to have speciated from *R. evenia*, with which it has been closely associated in all studies so far (Crawford et al. 1993a; Sang et al. 1995; Takayama et al. 2015). Previous investigations on ITS 1 and 2 in *Robinsonia* (Sang et al. 1995) have shown sequence divergence between *R. evenia* and *R. masafuerae* as only 0.0063 (two base substitutions). Although one cannot place an absolute time on this divergence, it is the lowest level among any pair of species in the genus, which correlates well with the youthful geological age of Alejandro Selkirk Island. Genetic variation in *R. masafuerae* is much lower from AFLP data than in *R. evenia* from Robinson Crusoe (Table 1 and Fig. 5), but in SSRs, the pattern reverses with the anagenetically derived species, *R. masafuerae*, showing more variation than any single one of the cladogenetically originated species on Robinson Crusoe (Table 1).

It is also possible to make comparisons between populations of continental progenitors with endemic island derivatives. In the case of *Myrceugenia schulzei*, the closest continental congener is *M. colchaguensis* (Landrum 1981a, b; Ruiz et al. 2004). Although the sampling of populations on the continent is limited to two populations,

Table 3. Summary of two-way ANOVA based on Table 1. TNB, total number of bands (fragments); PPB, percentage of polymorphic bands; SDI, Shannon Diversity Index; AGDOL, average gene diversity over loci; RI, rarity index; H_O , observed proportion of heterozygotes; H_E , expected proportion of heterozygotes; N_A , number of alleles per locus; A_{R5} , allelic richness standardized by five individuals. For all F -values, the degree of freedom was 1. Bold font indicates significant values ($P < 0.05$).

	Factor	F-value	P-value
AFLPs			
TNB	Island	4.78	0.046
	Speciation mode	0.22	0.645
	Island vs. speciation mode	0.51	0.489
PPB	Island	0.67	0.427
	Speciation mode	2.60	0.129
	Island vs. speciation mode	2.05	0.174
SDI	Island	3.61	0.078
	Speciation mode	2.36	0.147
	Island vs. speciation mode	0.47	0.504
AGDOL	Island	0.85	0.372
	Speciation mode	4.09	0.063
	Island vs. speciation mode	4.67	0.048
RI	Island	4.63	0.049
	Speciation mode	13.71	0.002
	Island vs. speciation mode	10.53	0.006
SSRs			
H_O	Island	2.03	0.176
	Speciation mode	11.65	0.004
	Island vs. speciation mode	1.64	0.221
H_E	Island	0.47	0.502
	Speciation mode	3.44	0.085
	Island vs. speciation mode	0.19	0.671
N_A	Island	0.47	0.502
	Speciation mode	3.44	0.085
	Island vs. speciation mode	0.19	0.671
A_{R5}	Island	0.10	0.752
	Speciation mode	4.54	0.051
	Island vs. speciation mode	0.11	0.744

the amount of genetic diversity is particularly low as shown by AFLP data, although somewhat higher with SSRs (López-Sepúlveda et al. 2013b). Although *M. schulzei* is known only on the younger island, it did not diverge from *M. fernandeziana* on the older island because the two are unrelated (Murillo-Aldana et al. 2012), so much so that the latter has now been transferred to another genus (*Nothomyrcia*; Murillo-Aldana and Ruiz 2011). With *D. confertifolia*, comparisons with *D. winteri* and *D. andina* show less genetic variation in the two latter species as




Figure 3. SplitsTree NeighborNet showing genetic relationships based on AFLPs among individuals in endemic species of *Myrceugenia* (A and B), *Drimys* (C), *Rhapsithamnus* (D), *Robinsonia* (E) and *Erigeron* (F) in the Juan Fernández Archipelago. Closely related continental relatives are also shown in A, C and D. Orange = species and populations on Robinson Crusoe Island; blue = on Alejandro Selkirk Island and black = on the or islands continent.

seen from AFLPs and SSRs (López-Sepúlveda et al. 2014). In *R. venustus*, which is a congener of *R. spinosus* (the only other known species in the genus; Moldenke 1937; Crawford et al. 1993b), the amount of genetic diversity is again greater in the population on Robinson Crusoe Island

than documented on the continent, although considerably lower in the population on Alejandro Selkirk (P. López-Sepúlveda, K. Takayama, D. J. Crawford, J. Greimler, P. Peñailillo, M. Baeza, E. Ruiz, G. Kohl, K. Tremetsberger, A. Gatica, L. Letelier, P. Novoa, J. Novak, T. F. Stuessy,




Figure 4. Neighbour-joining tree showing genetic relationships based on SSRs among populations in endemic species of *Myrceugenia* (A and B), *Drimys* (C), *Rhabithamnus* (D), *Robinsonia* (E) and *Erigeron* (F) in the Juan Fernández Archipelago. Closely related continental relatives are also shown in A, C and D. Orange = species and populations on Robinson Crusoe Island; blue = on Alejandro Selkirk Island and black = on the continent.

submitted for publication). These results support the concept that over time, considerable genetic variation can accumulate in anagenetically derived populations, so much so that the degree of variation can approximate and even surpass that in the progenitor source populations.

Cladogenesis

Two of the largest genera of the archipelago are *Robinsonia* with eight endemic species and *Erigeron* with six. Both are in Asteraceae, although unrelated and placed in different tribes (Senecioneae vs. Astereae, respectively).

Robinsonia has adaptively radiated on Robinson Crusoe Island during the past 4 million years (maximum value) and *Erigeron* has done so on Alejandro Selkirk Island in the past 1–2 million years.

Robinsonia is the second largest genus in the archipelago. The largest is *Dendroseris*, also of Asteraceae but from still another tribe (Cichorieae). This latter genus is of interest as it has derived cladogenetically on the older island with three independent dispersals to the younger island and three anagenetic speciations there (Sanders et al. 1987; Pacheco et al. 1991; Sang et al. 1994). Most of




Figure 5. Summary of genetic diversities, AGDOL, within the endemic species of *Drimys*, *Myrceugenia* and *Rhaphithamnus* having originated by anagenesis, and *Robinsonia* and *Erigeron* having been derived through cladogenesis. *Robinsonia masafuerae* from the younger island is also an anagenetic derivative from the cladogenetic complex of *Robinsonia* on the older island. AS, Alejandro Selkirk Island; RC, Robinson Crusoe Island. White bar indicates an endemic species in RC, grey bar an endemic species in AS and black bar multiple species or islands combined.

these species are quite rare now, however, which precluded our being able to obtain sufficient population data for genetic evaluation. *Robinsonia* has eight species, but two are presumed extinct (*R. berteroi* and *R. megacephala*; Danton et al. 2006). Our studies have focussed on five species having originated cladogenetically on the older island. Comments have already been made regarding the one anagenetically derived species (*R. masafuerae*) on Alejandro Selkirk Island. The results from AFLP data are shown in Fig. 3 and from SSRs in Fig. 4. Most notable from the SplitsTree graph in Fig. 3 is that the different species of *Robinsonia* are very distinct genetically. Divergence has obviously taken place during adaptive radiation and also during a maximum time available of 4 million years. The species *R. gayana*, *R. thurifera* and *R. saxatilis* form an evolutionary complex, which taxonomically has been regarded as sect. *Robinsonia* (Skottsberg 1922, as sect. *Symphylepis*; Takayama et al. 2015). *Robinsonia gracilis* ties with *R. evenia* and its close anagenetic relative *R. masafuerae* in sect. *Eleutherolepis* (Skottsberg 1922). With SSR data (Fig. 4), the species are also very distinct and genetically more cohesive, with the anagenetic species *R. masafuerae* showing the greatest genetic diversity.

Another important point seen clearly in Figs 3 and 4 is that the range of genetic diversity within each of these cladogenetic species is limited in comparison with the anagenetically derived species discussed above.

Although *Erigeron* is not an endemic genus in the archipelago, six endemic species occur there having evolved via cladogenesis and adaptive radiation. The origin of this complex is unusual in that the colonist(s) presumably arrived directly to the younger island (Valdebenito et al. 1992). Amplified fragment length polymorphism and SSR data (Figs 3 and 4) reveal considerable genetic diversity within these endemic species, and each species is reasonably distinct. An exception is the *Erigeron ingae* complex consisting of *E. ingae*, *E. luteoviridis* and *E. turricola*. These species are sometimes difficult to distinguish morphologically. Solbrig (1962) and Marticorena et al. (1998), for example, placed *E. turricola* into synonymy with *E. ingae*, but Danton et al. (2006) kept them distinct. The molecular data parallel this morphological inconsistency. This may be a population complex in early stages of speciation, now undergoing divergence from within a pool of morphological and genetic variation. All of these species grow in the ‘alpine zone’ on the younger island (Skottsberg 1922),

Table 4. Generalized comparison of the levels of genetic diversity obtained with AFLPs and SSRs from species that have originated via anagenesis and cladogenesis on the two islands of the Juan Fernández Archipelago. See Table 1 for the actual data. RC, Robinson Crusoe Island; AS, Alejandro Selkirk Island.

	Anagenesis		Cladogenesis	
	RC	AS	RC	AS
AFLPs				
Total number of bands (TNB)	High	Medium	High	Low
Percentage of polymorphic bands (PPB)	High	Low	Low	Low
Shannon Diversity Index (SDI)	High	Medium high	Medium high	Low
Average gene diversity over loci (AGDOL)	High	Medium high	Low	Medium high
Rarity index (RI)	Medium	High	Medium	Very low
Microsatellites (SSRs)				
Observed proportion of heterozygotes (H_O)	High	High	Medium	Low
Expected proportion of heterozygotes (H_E)	High	High	Medium	Medium
Number of alleles per locus (N_A)	High	Medium	Low	Low
Inbreeding coefficient (F_{IS})	Low	Low	Low	High
Allelic richness (A_{RS})	High	High	Low	Low

and we have not noticed any clear habitat differences among them. The species *E. rupicola* is confined to coastal rocks along the sea and also penetrates into the quebradas (ravines); its close relative, *E. stuessyi*, is also found on rocky ledges but residing inside the cool and deep ravines. *Erigeron fernandezianus* occurs in a broad altitudinal range (100–1200 m), and it inhabits mainly rocky areas in middle elevation plains, quebradas and ridges. This species also occurs on the older island, but it is found there in many plant communities and especially in disturbed sites. It appears, therefore, to be an example of back migration from the younger to the older island (Valdebenito et al. 1992; López-Sepúlveda et al. 2015).

Although most species of *Erigeron* on the younger island are distinct genetically, the degree of distinctness is much less than observed among species of *Robinsonia* on the older island (Figs 3 and 4). It may be that these species of *Erigeron* have had less time to diverge in comparison with those of *Robinsonia*. With the passage of time, therefore, the genetic profiles of species undergoing adaptive radiation may remain narrow due to strong directional selection in each different habitat. In both *Erigeron* and *Robinsonia*, however, the range of genetic variation seen is less than that in the anagenetically derived species.

Discussion

Comparison of anagenesis and cladogenesis

Predictions from theory (Stuessy 2007) would suggest that higher levels of genetic diversity should be found

within the anagenetically derived species. This is because the founding population increases in size over time, accumulating genetic diversity mainly through mutation and recombination. One would expect no (or very little) geographic partitioning over the landscape. Likewise, due to a lack of strong selection, one would not expect to find high levels of private alleles or bands, nor a high RI. With cladogenetic speciation, on the other hand, one would expect less overall genetic diversity within each species, but with more private alleles due to strong directional selection. As for impact from the age of the islands, one would predict less total genetic diversity within anagenetically derived species on the younger island because diversity increases through time. As for the cladogenetic species, one would predict less genetic divergence (distinctness) on the younger island in comparison with species on the older island, because directional selection continues over time and refines the genetic profile of each species as it adapts to the particular ecological zone.

Results from genetic analyses of 5 anagenetic species and 10 cladogenetic species allow comparisons between the two modes of speciation and the two islands of differing ages (Tables 1–3). A number of general points can be observed (Table 4 and Fig. 5). First, in anagenetic species, the level of genetic diversity tends to be higher per species than in the cladogenetic species, especially on Robinson Crusoe Island. This can be seen in percentage of polymorphic bands, SDI, AGDOL, observed proportion of heterozygotes, expected proportion of heterozygotes, number of alleles per locus and allelic richness. Second, in the

anagenetic species, the individuals on each island behave genetically as one large population, showing no genetic pattern over the landscape (López-Sepúlveda et al. 2013b, 2014; Takayama et al. 2015; P. López-Sepúlveda, K. Takayama, D. J. Crawford, J. Greimler, P. Peñailillo, M. Baeza, E. Ruiz, G. Kohl, K. Tremetsberger, A. Gatica, L. Letelier, P. Novoa, J. Novak, T. F. Stuessy, submitted for publication). This is true on both islands of differing ages. This suggests that this pattern can develop easily within 1–2 million years and that it can persist for up to 4 million. This is consistent with the results reported for Ullung Island, Korea, which is known to be 1.8 million years old (Pfosser et al. 2005; Takayama et al. 2012, 2013a). Third, the ability of an immigrant population to radiate adaptively has much to do with the properties of the colonists (and progenitors) and less with differences of habitat. Some colonists remain as a single larger population and are not responsive to adaptive change in different ecological zones, whereas others disperse well to micro-zones and quickly become modified morphologically and genetically. Fourth, perhaps most importantly, the total amount of genetic diversity within an anagenetically derived species in comparison with an entire adaptively radiating lineage is approximately the same (Fig. 5).

Genetics of speciation in endemic plants of oceanic islands

A number of previous studies have assessed levels of genetic variation within and among populations of endemic species of the Juan Fernández Archipelago with other markers such as isozymes, random amplified polymorphic DNA (RAPDs) and inter simple sequence repeats (ISSRs). Isozymes have been analysed in *Dendroseris* (Crawford et al. 1987), *Chenopodium sanctae-clarae* (Crawford et al. 1988), *Wahlenbergia* (Crawford et al. 1990), *Robinsonia* (Crawford et al. 1992), *Lactoris* (Crawford et al. 1994) and *Myrceugenia* (Jensen et al. 2002). RAPDs have been investigated in *Dendroseris* (Esselman et al. 2000) and *Lactoris* (Brauner et al. 1992), and ISSRs also in *Lactoris* (Crawford et al. 2001b).

Crawford et al. (2001a) summarized the results from isozyme studies on 29 endemic species of the Juan Fernández Archipelago, and this represents the best set of observations to compare with the AFLP and SSR data summarized here. The most conspicuous result is that the mean genetic diversities at the species level are low ($H_{es} = 0.065$). Higher levels of diversity were seen in larger populations or in many small populations and also in out-crossing species in contrast to selfers. Of relevance for comparisons to the present study, isozymes have been analysed from four species of *Robinsonia* and in *M. fernandeziana*, *E. fernandezianus* and *R. venustus*. It is difficult to compare the results of the isozymes because they provide

less detailed genetic information than from AFLPs and SSRs. Isozyme studies on the endemic *Lactoris fernandezianus*, for example Crawford et al. (1994), revealed virtually no variation, but ISSRs showed considerable variation within and among populations (Crawford et al. 2001b). Studies on isozymes (Crawford et al. 1987) and RAPDs (Esselman et al. 2000) from *Dendroseris* showed greater resolution of relationships from the latter. The isozyme data for the four cladogenetically derived species of *Robinsonia* show higher levels of genetic variation than in the anagenetic *R. venustus* (Crawford et al. 1993b) and *Myrceugenia* (Jensen et al. 2002), which would be in contrast to the trends documented here. It is important, therefore, that for questions involving population genetics in endemic plants of oceanic islands, rapidly evolving markers need to be used.

The employment of AFLPs and SSRs in the present study from 15 species of the Juan Fernández Archipelago, therefore, does provide detailed genetic data at the population level for purposes of comparing consequences of different modes of speciation. A general review has recently been published on the general topic of interpretation of genetic variation within endemic species of oceanic islands (Stuessy et al. 2014), and the present data corroborate ideas summarized there. Clearly, the alternative modes of speciation, anagenesis and cladogenesis result in different genetic consequences. Interpretation of the evolutionary significance of levels of genetic diversity, therefore, must be done in context of type of speciation. As can be seen in the results of adaptive radiation in *Erigeron* and *Robinsonia*, on the young and older islands, respectively, the geological age of the island also matters, as this provides the time dimension in which the evolutionary processes unfold.

Another very significant impact on levels of genetic variation in populations of endemic plants of oceanic islands is that from human activity. Because oceanic islands often have agreeable climates and attractive beaches, people have come to live, play and build homes and apartments, all of which have caused pressures on the native vegetation. In the Juan Fernández Archipelago, for example, people have been living continuously on Robinson Crusoe Island for >300 years (Woodward 1969; Wester 1991). It is not impossible that the species of *Robinsonia* on the older island have suffered some genetic loss due to human activity. Although these species occur either on high ridges or in deep forests, far removed from most persons who live at sea level in the village (San Juan Bautista), incursions into the native forest must have taken place and some plants destroyed. It is known that two species of *Robinsonia*, both on Robinson Crusoe Island, are now extinct (*R. berteroi* and *R. megacephala*; Danton and Perrier 2005; Danton et al.

2006). Assessing the level of human impact on the vegetation of an oceanic island, therefore, is challenging. At least in the Juan Fernández Archipelago, there were no aboriginal peoples, and human activity could only have begun with discovery by Europeans (Juan Fernández; Medina 1974) at the end of the 16th century. Since that time, however, considerable negative impact from human activity has been documented in the archipelago (Wester 1991; Matthei et al. 1993; Stuessy et al. 1997; Swenson et al. 1997; Cuevas and Leersum 2001; Greimler et al. 2002; Dirnböck et al. 2003; Cuevas et al. 2004; Ricci 2006; Vargas et al. 2011), especially from introduced animals, such as rats, rabbits and goats (e.g. Camus et al. 2008). These combined activities have surely had some impact on the levels of genetic variation within and among populations.

Sources of Funding

This work was supported by an FWF (Austrian Science Fund) grant (P21723-B16) to T.F.S. and a Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowship for Research Abroad (grant 526) to Ko.T.

Contributions by the Authors

Ko.T. conceived the idea behind the article; all authors participated in the field work except G.K. and Ka.T.; J.N., P.L.-S., G.K. and Ko.T. completed the laboratory work; J.N. coordinated the NGS data acquisition; T.F.S. and Ko.T. wrote the initial draft and all authors contributed to subsequent drafts and offered comments for improvement.

Conflict of Interest Statement

None declared.

Acknowledgements

We very much appreciate the generous logistic and facility support of Sr Iván Leiva, Chief of the Robinson Crusoe Islands national park, administered by the Corporación Nacional Forestal (CONAF); the help and cooperation in fieldwork from the CONAF guides, especially Jorge Angulo, Danilo Arredondo, Danilo Arredondo, Jr, Oscar Chamorro, Michael González, Bernardo López, Eduardo Paredes, Ramon Schiller and Manuel Tobar; and the Armada de Chile for logistic support in transporting supplies from the continent to the islands. The results presented in this paper form part of an Open Partnership Joint Project of the JSPS Bilateral Joint Research program.

Literature Cited

- Baldwin BG. 2003. A phylogenetic perspective on the origin and evolution of Madiinae. In: Carlquist S, Baldwin BG, Carr GD, eds. *Tarweeds & silverswords: evolution of the Madiinae (Asteraceae)*. St. Louis: Missouri Botanical Garden Press, 193–228.
- Baldwin BG, Crawford DJ, Francisco-Ortega J, Kim S-C, Sang T, Stuessy TF. 1998. Molecular phylogenetic insights on the origin and evolution of oceanic island plants. In: Soltis DE, Soltis PS, Doyle JJ, eds. *Molecular systematics of plants II: DNA sequencing*. Boston: Kluwer, 410–441.
- Böhle U-R, Hilger HH, Martin WF. 1996. Island colonization and evolution of the insular woody habit in *Echium* L. (Boraginaceae). *Proceedings of the National Academy of Sciences of the USA* **93**: 11740–11745.
- Bramwell D, Caujapé-Castells J. 2011. *The biology of island floras*. Cambridge: Cambridge University Press.
- Brauner S, Crawford DJ, Stuessy TF. 1992. Ribosomal DNA and RAPD variation in the rare plant family Lactoridaceae. *American Journal of Botany* **79**:1436–1439.
- Bryant D, Moulton V. 2004. Neighbor-Net: an agglomerative method for the construction of phylogenetic networks. *Molecular Biology and Evolution* **21**:255–265.
- Camus P, Castro S, Jaksic F. 2008. El conejo europeo en Chile: Historia de una invasión histórica. *Historia* **41**:305–339.
- Carlquist S. 1974. *Island biology*. New York: Columbia University Press.
- Carlquist S, Baldwin BG, Carr GD, eds. 2003. *Tarweeds & silverswords: evolution of the Madiinae (Asteraceae)*. St. Louis: Missouri Botanical Garden.
- Crawford DJ. 2010. Progenitor-derivative species pairs and plant speciation. *Taxon* **59**:1413–1423.
- Crawford DJ, Stuessy TF, Silva M. 1987. Allozyme divergence and the evolution of *Dendroseris* (Compositae: Lactuceae) on the Juan Fernandez Islands. *Systematic Botany* **12**:435–443.
- Crawford DJ, Stuessy TF, Silva M. 1988. Allozyme variation in *Chenopodium sanctae-clarae*, an endemic species of the Juan Fernandez Islands, Chile. *Biochemical Systematics and Ecology* **16**: 279–284.
- Crawford DJ, Stuessy TF, Lammers TG, Silva M, Pacheco P. 1990. Allozyme variation and evolutionary relationships among three species of *Wahlenbergia* (Campanulaceae) in the Juan Fernandez Islands. *Botanical Gazette* **151**:119–124.
- Crawford DJ, Stuessy TF, Haines DW, Cosner MB, Silva M, Lopez P. 1992. Allozyme diversity within and divergence among four species of *Robinsonia* (Asteraceae: Senecioneae), a genus endemic to the Juan Fernandez Islands, Chile. *American Journal of Botany* **79**:962–966.
- Crawford DJ, Stuessy TF, Cosner MB, Haines DW, Silva M. 1993a. Ribosomal and chloroplast DNA restriction site mutations and the radiation of *Robinsonia* (Asteraceae: Senecioneae) on the Juan Fernandez Islands. *Plant Systematics and Evolution* **184**: 233–239.
- Crawford DJ, Stuessy TF, Rodriguez R, Rondinelli M. 1993b. Genetic diversity in *Rhaphithamnus venustus* (Verbenaceae), a species endemic to the Juan Fernandez Islands. *Bulletin of the Torrey Botanical Club* **120**:23–28.
- Crawford DJ, Stuessy TF, Cosner MB, Haines DW, Wiens D, Peñaillo P. 1994. *Lactoris fernandeziana* (Lactonaceae) on the Juan

- Fernandez Islands: allozyme uniformity and field observations. *Conservation Biology* **8**:277–280.
- Crawford DJ, Sang T, Stuessy TF, Kim S-C, Silva M. 1998. *Dendroseris* (Asteraceae: Lactuceae) and *Robinsonia* (Asteraceae: Senecioideae) on the Juan Fernandez Islands: similarities and differences in biology and phylogeny. In: Stuessy TF, Ono M, eds. *Evolution and speciation of island plants*. Cambridge: Cambridge University Press, 97–119.
- Crawford DJ, Ruiz E, Stuessy TF, Tepe E, Aqueveque P, González F, Jensen RJ, Anderson GJ, Bernardello G, Baeza CM, Swenson U, Silva M. 2001a. Allozyme diversity in endemic flowering plant species of the Juan Fernandez Archipelago, Chile: ecological and historical factors with implications for conservation. *American Journal of Botany* **88**:2195–2203.
- Crawford DJ, Tago-Nakazawa M, Stuessy TF, Anderson GJ, Bernardello G, Ruiz E, Jensen RJ, Baeza C, Wolfe AD, Silva M. 2001b. Intersimple sequence repeat (ISSR) variation in *Lactoris fernandeziana* (Lactoridaceae), a rare endemic of the Juan Fernández Archipelago, Chile. *Plant Species Biology* **16**:185–192.
- Cuevas JG, van Leersum G. 2001. Project “Conservation, restoration, and development of the Juan Fernández Islands, Chile”. *Revista Chilena de Historia Natural* **74**:899–910.
- Cuevas JG, Marticorena A, Cavieres LA. 2004. New additions to the introduced flora of the Juan Fernández Islands: origin, distribution, life history traits, and potential of invasion. *Revista Chilena de Historia Natural* **77**:523–538.
- Danton P, Perrier C. 2005. Note sur la disparition d’une espèce emblématique, *Robinsonia berteroi* (DC.) Sanders, Stuessy & Martic. (Asteraceae) dans l’île Robinson Crusoe, archipel Juan Fernández (Chili). *Journal de Botanique de la Société Botanique de France* **31**: 1–6.
- Danton P, Perrier C, Martinez Reyes G. 2006. Nouveau catalogue de la flore vasculaire de l’archipel Juan Fernández (Chili). *Nuevo catálogo de la flora vascular del Archipiélago Juan Fernández (Chile)*. *Acta Botanica Gallica* **153**:399–587.
- Darwin C. 1842. *The structure and distribution of coral reefs*. London: Smith, Elder and Co.
- Dirnböck T, Greimler J, López P, Stuessy TF. 2003. Predicting future threats to the native vegetation of Robinson Crusoe Island, Juan Fernandez Archipelago, Chile. *Conservation Biology* **17**: 1650–1659.
- Ehrich D. 2006. AFLPdat: a collection of R functions for convenient handling of AFLP data. *Molecular Ecology Notes* **6**:603–604.
- Eliasson U. 1974. Studies in Galapagos Plants XIV. The genus *Scalesia* Arn. *Opera Botanica* **36**:1–117.
- Esselman EJ, Crawford DJ, Brauner S, Stuessy TF, Anderson GJ, Silva M. 2000. RAPD marker diversity within and divergence among species of *Dendroseris* (Asteraceae: Lactuceae). *American Journal of Botany* **87**:591–596.
- Excoffier L, Laval G, Schneider S. 2005. Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1**:47–50.
- Frankham R. 1997. Do island populations have less genetic variation than mainland populations? *Heredity* **78**:311–327.
- Givnish TJ, Millam KC, Mast AR, Paterson TB, Theim TJ, Hipp AL, Henss JM, Smith JF, Wood KR, Sytsma KJ. 2009. Origin, adaptive radiation and diversification of the Hawaiian lobeliads (Asterales: Campanulaceae). *Proceedings of the Royal Society B Biological Sciences* **276**:407–416.
- Gleiser G, Verdú M, Segarra-Moragues JG, González-Martínez SC, Pannell JR. 2008. Disassortative mating, sexual specialization, and the evolution of gender dimorphism in heterodichogamous *Acer opalus*. *Evolution* **62**:1676–1688.
- Greimler J, Stuessy TF, Swenson U, Baeza CM, Matthei O. 2002. Plant invasions on an oceanic archipelago. *Biological Invasions* **4**: 73–85.
- Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution* **23**:254–267.
- Jensen RJ, Schwoyer M, Crawford DJ, Stuessy TF, Anderson GJ, Baeza CM, Silva M, Ruiz E. 2002. Patterns of morphological and genetic variation among populations of *Myrceugenia fernandeziana* (Myrtaceae) on Masatierra Island: implications for conservation. *Systematic Botany* **27**:534–547.
- Jorgensen TH, Olesen JM. 2001. Adaptive radiation of island plants: evidence from *Aeonium* (Crassulaceae) of the Canary Islands. *Perspectives in Plant Ecology, Evolution and Systematics* **4**:29–42.
- Kikuchi S, Shibata M, Tanaka H, Yoshimaru H, Niiyama K. 2009. Analysis of the disassortative mating pattern in a heterodichogamous plant, *Acer mono* Maxim. using microsatellite markers. *Plant Ecology* **204**:43–54.
- Kim YK. 1985. Petrology of Ulreung volcanic island, Korea—Part 1. Geology. *Journal of the Japanese Association of Mineralogists, Petrologists and Economic Geologists* **80**:128–135.
- Landrum L. 1981a. A monograph of the genus *Myrceugenia* (Myrtaceae). Bronx, NY: New York Botanical Garden Press.
- Landrum LR. 1981b. The phylogeny and geography of *Myrceugenia* (Myrtaceae). *Brittonia* **33**:105–129.
- Langella O. 1999. *Populations*, 1.2.30. <http://www.bioinformatics.org/~tryphon/populations/> (15 October 2014).
- Liu H-Y. 1989. *Systematics of Aeonium (Crassulaceae)*. Special Publications of the National Museum of Natural Science (Taichung), No. 3, 1–102.
- López-Sepúlveda P, Tremetsberger K, Ortiz MA, Baeza CM, Peñailillo P, Stuessy TF. 2013a. Radiation of the *Hypochaeris apargioides* complex (Asteraceae: Cichorieae) of southern South America. *Taxon* **62**:550–564.
- López-Sepúlveda P, Takayama K, Greimler J, Peñailillo P, Crawford DJ, Baeza M, Ruiz E, Kohl G, Tremetsberger K, Gatica A, Letelier L, Novoa P, Novak J, Stuessy TF. 2013b. Genetic variation (AFLPs and nuclear microsatellites) in two anagenetically derived endemic species of *Myrceugenia* (Myrtaceae) on the Juan Fernández Islands, Chile. *American Journal of Botany* **100**:722–734.
- López-Sepúlveda P, Takayama K, Greimler J, Crawford DJ, Peñailillo P, Baeza M, Ruiz E, Kohl G, Tremetsberger K, Gatica A, Letelier L, Novoa P, Novak J, Stuessy TF. 2014. Progressive migration and anagenesis in *Drimys confertifolia* of the Juan Fernández Archipelago, Chile. *Journal of Plant Research* **128**:73–90.
- López-Sepúlveda P, Takayama K, Crawford DJ, Greimler J, Peñailillo P, Baeza M, Ruiz E, Kohl G, Tremetsberger K, Gatica A, Letelier L, Novoa P, Novak J, Stuessy TF. 2015. Speciation and biogeography of *Erigeron* (Asteraceae) endemic to the Juan Fernández Archipelago, Chile, based on AFLPs and SSRs. *Systematic Botany*, **40**: doi: 10.1600/036364415X689311.
- Lowe A, Harris S, Ashton P. 2004. *Ecological genetics: design, analysis, and application*. Oxford: Blackwell.
- Marticorena C, Stuessy TF, Baeza M. 1998. Catálogo de la flora vascular del Archipiélago de Juan Fernández, Chile. *Gayana Botánica* **55**:187–211.

- Matthei O, Marticorena C, Stuessy TF. 1993. La flora adventicia del archipiélago de Juan Fernández. *Gayana Botánica* **50**:69–102.
- Medina JT. 1974. *El piloto Juan Fernández, descubridor de las islas que llevan su nombre, y Juan Jufre, armador de la expedición que hizo en busca de otras en el Mar del Sur*. Santiago: Editora Nacional Gabriela Mistral.
- Moldenke HN. 1937. A monograph of the genus *Rhaphithamnus*. *Repertorium Novarum Specierum Regni Vegetabilis* **42**:62–82.
- Murillo-Aldana J, Ruiz E. 2011. Revalidación de *Nothomyrcia* (Myrtaceae), un género endémico del Archipiélago de Juan Fernández. *Gayana Botánica* **68**:129–134.
- Murillo-Aldana J, Ruiz-P E, Landrum LR, Stuessy TF, Barfuss MHJ. 2012. Phylogenetic relationships in *Myrceogenia* (Myrtaceae) based on plastid and nuclear DNA sequences. *Molecular Phylogenetics and Evolution* **62**:764–776.
- Nei M, Tajima F, Tatenos Y. 1983. Accuracy of estimated phylogenetic trees from molecular data II. Gene frequency data. *Journal of Molecular Evolution* **19**:153–170.
- Pacheco P, Crawford DJ, Stuessy TF, Silva OM. 1991. Flavonoid evolution in *Dendroseris* (Compositae, Lactuceae) from the Juan Fernandez Islands, Chile. *American Journal of Botany* **78**:534–543.
- Peakall R, Smouse PE. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* **6**:288–295.
- Perugganan MD, Remington DL, Robichaux RH. 2003. Molecular evolution of regulatory genes in the silversword alliance. In: Carlquist S, Baldwin BG, Carr GD, eds. *Tarweeds & silverswords: evolution of the Madiinae (Asteraceae)*. St. Louis: Missouri Botanical Garden Press, 171–182.
- Pfossner M, Jakubowsky G, Schlüter PM, Fer T, Kato H, Stuessy TF, Sun B-Y. 2005. Evolution of *Dystaenia takesimana* (Apiaceae), endemic to Ullung Island, Korea. *Plant Systematics and Evolution* **256**:159–170.
- Raymond M, Rousset F. 1995. GENEPOP (version 1.2): population genetics software for exact test and ecumenicism. *Journal of Heredity* **86**:248–249.
- R Core Team. 2013. *A language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.
- Rensch B. 1959. *Evolution above the species level*. New York: Columbia University Press.
- Ricci M. 2006. Conservation status and *ex situ* cultivation efforts of endemic flora of the Juan Fernández Archipelago. *Biodiversity and Conservation* **15**:3111–3130.
- Ruiz E, Crawford DJ, Stuessy TF, González F, Samuel R, Becerra J, Silva M. 2004. Phylogenetic relationships and genetic divergence among endemic species of *Berberis*, *Gunnera*, *Myrceogenia* and *Sophora* of the Juan Fernández Islands (Chile) and their continental progenitors based on isozymes and nrITS sequences. *Taxon* **53**:321–332.
- Rundell RJ, Price TD. 2009. Adaptive radiation, nonadaptive radiation, ecological speciation and nonecological speciation. *Trends in Ecology and Evolution* **24**:394–399.
- Sanders RW, Stuessy TF, Marticorena C, Silva M. 1987. Phytogeography and evolution of *Dendroseris* and *Robinsonia*, tree-Compositae of the Juan Fernandez Islands. *Opera Botanica* **92**:195–215.
- Sang T, Crawford DJ, Kim S-C, Stuessy TF. 1994. Radiation of the endemic genus *Dendroseris* (Asteraceae) on the Juan Fernandez Islands: evidence from sequences of the ITS regions of nuclear ribosomal DNA. *American Journal of Botany* **81**:1494–1501.
- Sang T, Crawford DJ, Stuessy TF, Silva M. 1995. ITS sequences and the phylogeny of the genus *Robinsonia* (Asteraceae). *Systematic Botany* **20**:55–64.
- Schluter D. 2001. Ecology and the origin of species. *Trends in Ecology and Evolution* **16**:372–380.
- Schlüter PM, Harris SA. 2006. Analysis of multilocus fingerprinting data sets containing missing data. *Molecular Ecology Notes* **6**:569–572.
- Skottsberg C. 1922. The phanerogams of the Juan Fernandez Islands. In: Skottsberg C, ed. *The natural history of Juan Fernandez and Easter Island*, Vol. 2. Uppsala: Almqvist & Wiksells, 95–240.
- Solbrig OT. 1962. The South American species of *Erigeron*. *Contributions from the Gray Herbarium of Harvard University* **191**:3–79.
- Stuessy TF. 1995. Juan Fernández Islands. In: Davis SD, Heywood VH, Hamilton AC, eds. *Centres of plant diversity: a guide and strategy of their conservation*. Cambridge: IUCN Publications Unit, 565–568.
- Stuessy TF. 2007. Evolution of specific and genetic diversity during ontogeny of island floras: the importance of understanding process for interpreting island biogeographic patterns. In: Ebach MC, Tangney RS, eds. *Biogeography in a changing world*. Boca Raton: CRC Press, 117–133.
- Stuessy TF, Foland KA, Sutter JF, Sanders RW, Silva M. 1984. Botanical and geological significance of potassium-argon dates from the Juan Fernandez Islands. *Science* **225**:49–51.
- Stuessy TF, Crawford DJ, Marticorena C. 1990. Patterns of phylogeny in the endemic vascular flora of the Juan Fernandez Islands, Chile. *Systematic Botany* **15**:338–346.
- Stuessy TF, Swenson U, Marticorena C, Mathei O, Crawford DJ. 1997. Loss of plant diversity and extinction on Robinson Crusoe Island, Chile. In: Peng C-I, ed. *Rare, threatened, and endangered floras of Asia and the Pacific Rim*. Taipei: Academia Sinica (Monograph Series No. 16), 147–257.
- Stuessy TF, Crawford DJ, Marticorena C, Rodriguez R. 1998. Island biogeography of angiosperms of the Juan Fernandez archipelago. In: Stuessy TF, Ono M, eds. *Evolution and speciation of island plants*. Cambridge: Cambridge University Press, 121–138.
- Stuessy TF, Jakubowsky G, Gómez RS, Pfossner M, Schlüter PM, Fer T, Sun B-Y, Kato H. 2006. Anagenetic evolution in island plants. *Journal of Biogeography* **33**:1259–1265.
- Stuessy TF, Takayama K, López-Sepúlveda P, Crawford DJ. 2014. Interpretation of patterns of genetic variation in endemic plant species of oceanic islands. *Botanical Journal of the Linnean Society* **174**:276–288.
- Swenson U, Stuessy TF, Baeza M, Crawford DJ. 1997. New and historical plant introductions, and potential pests in the Juan Fernández Islands, Chile. *Pacific Science* **51**:233–253.
- Takayama K, López P, König C, Kohl G, Novak J, Stuessy TF. 2011. A simple and cost-effective approach for microsatellite isolation in non-model plant species using small-scale 454 pyrosequencing. *Taxon* **60**:1442–1449.
- Takayama K, Sun B-Y, Stuessy TF. 2012. Genetic consequences of anagenetic speciation in *Acer okamotoanum* (Sapindaceae) on Ullung Island, Korea. *Annals of Botany* **109**:321–330.
- Takayama K, Sun B-Y, Stuessy TF. 2013a. Anagenetic speciation in Ullung Island, Korea: genetic diversity and structure in the island endemic species, *Acer takesimensis* (Sapindaceae). *Journal of Plant Research* **126**:323–333.

- Takayama K, López-Sepúlveda P, Kohl G, Novak J, Stuessy TF. 2013b. Development of microsatellite markers in *Robinsonia* (Asteraceae) an endemic genus of the Juan Fernández Archipelago, Chile. *Conservation Genetics Resources* 5:63–67.
- Takayama K, López-Sepúlveda P, Greimler J, Crawford DJ, Peñailillo P, Baeza M, Ruiz E, Kohl G, Tremetsberger K, Gatica A, Letelier L, Novoa P, Novak J, Stuessy TF. 2015. Relationships and genetic consequences of contrasting modes of speciation among endemic species of *Robinsonia* (Asteraceae, Senecioneae) of the Juan Fernández Archipelago, Chile, based on AFLPs and SSRs. *New Phytologist* 205:415–428.
- Tremetsberger K, Stuessy TF, Guo Y-P, Baeza CM, Weiss H, Samuel RM. 2003. Amplified fragment length polymorphism (AFLP) variation within and among populations of *Hypochaeris acaulis* (Asteraceae) of Andean southern South America. *Taxon* 52:237–245.
- Valdebenito H, Stuessy TF, Crawford DJ, Silva M. 1992. Evolution of *Erigeron* (Compositae) in the Juan Fernandez Islands, Chile. *Systematic Botany* 17:470–480.
- van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4:535–538.
- Vargas R, Reif A, Faúndez MJ. 2011. The forests of Robinson Crusoe Island, Chile: an endemism hotspot in danger. *Bosque* 32:155–164.
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Friters A, Pot J, Paleman J, Kuiper M, Zabeau M. 1995. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* 23:4407–4414.
- Wallace AR. 1881. *Island life*. London: Macmillan and Co.
- Wester L. 1991. Invasions and extinctions on Masatierra (Juan Fernández Islands): a review of early historical evidence. *Journal of Historical Geography* 17:18–34.
- Whittaker RJ, Fernández-Palacios JM. 2007. *Island biogeography*, 2nd edn. Oxford: Oxford University Press.
- Whittaker RJ, Triantis KA, Ladle RJ. 2008. A general dynamic theory of oceanic island biogeography. *Journal of Biogeography* 35: 977–994.
- Woodward RL. 1969. Robinson Crusoe's Island: a history of the Juan Fernandez Islands. Chapel Hill: University of North Carolina Press.
- Yim Y-J, Lee E-B, Kim S-H. 1981. *Vegetation of Ulreung and Dogdo Islands*. A Report on the Scientific Survey of the Ulreung and Dogdo Islands. Seoul: The Korean Association for Conservation of Nature.